

Microlocal perverse sheaves

Ingo Waschkes

Abstract

In this paper we give an explicit construction of the stack of microlocal perverse sheaves on the projective cotangent bundle of a complex manifold. Microlocal perverse sheaves will be represented as complexes of analytic ind-sheaves which have recently been studied by Kashiwara-Schapira [KS2]. This description allows us to formulate the microlocal Riemann-Hilbert correspondance in order to establish an equivalence of stacks with the stack of regular holonomic microdifferential modules.

Contents

1	Introduction	3
2	Abelian substacks of a prestack	5
2.1	A criterion for substacks	5
2.2	Limits and colimits in stacks	9
2.3	A criterion for abelian stacks	11
3	Microlocalization of sheaves	12
3.1	Notations	12
3.2	Microlocalization of sheaves	13
3.3	Refined microlocal cut-off	15
3.4	Morphisms in $D^b(k_X, \{x\} \times \dot{\delta})$	18
4	Microlocalization of constructible sheaves	20
4.1	Microlocalization of $\mathbb{R}$ -constructible sheaves	20
4.2	The category $D_{\mathbb{R}\text{-c}}^b(k_X, \{x\} \times \dot{\delta})$	22
4.3	Microlocally $\mathbb{C}$ -constructible sheaves	22
4.4	The category $D_{\mathbb{C}\text{-c}}^b(k_X, \{x\} \times \dot{\delta})$	24
5	Invariance by quantized contact transformations	24
5.1	Quantization of $\mathbb{R}$ -constructible sheaves	25
5.2	Quantization of $\mathbb{C}$ -constructible sheaves	28
6	Microlocally complex constructible sheaves on $\mathbb{C}^{\times p}$	29
6.1	Microlocal complex constructible sheaves in generic position	29
6.2	The generic position theorem	32
7	Microlocal perverse sheaves in $D_{\mathbb{R}\text{-c}}^b(k_X, S)$	33
7.1	Andronikof's prestack of microlocally perverse sheaves	33
7.2	The abelian category $D_{\text{perv}}^b(k_X, \mathbb{C}^{\times p})$	35

8	Microlocal perverse sheaves on P^*X	38
8.1	The stack of microlocal perverse sheaves	38
8.2	Autoduality	42
8.3	Integral transforms for microlocal perverse sheaves	42
9	Microlocal Riemann-Hilbert correspondence	48
9.1	The ind-objects $\mu\mathcal{O}_X$ and $\mu\mathcal{O}_X^t$	49
9.2	$\mathcal{E}_X$ -modules	51
9.3	The microdifferential structure of $\mu\mathcal{O}_X$ and $\mu\mathcal{O}_X^t$	53
9.4	Quantized contact transformations for $\mu\mathcal{O}_X$ and $\mu\mathcal{O}_X^t$	56
9.5	Classical Riemann-Hilbert Theorem and Ind-sheaves	59
9.6	Microlocal Riemann-Hilbert morphism	60
A	Stacks on Topological Spaces	62
A.1	Prestacks	63
A.2	Operations on prestacks	64
A.3	Stalks	64
A.4	Stacks	65
A.5	The stack associated to a prestack	66
A.6	Patching of stacks	68
B	The functor of ind-micolocalization	68
B.1	Ind-sheaves	68
B.2	Microlocalization of ind-sheaves	70

1 Introduction

In [K4] Kashiwara constructed the stack of microdifferential modules on a complex contact manifold, generalizing the stack of modules over the ring of microdifferential operators $\mathcal{E}_X$ on the projective cotangent bundle P^*X of a complex variety X . In his discussion Kashiwara asked for the construction of a stack of microlocal (complex) perverse sheaves that should be equivalent to the stack of regular holonomic modules by a microlocal Riemann-Hilbert correspondance. Such a stack should be defined over any field k , but the Riemann-Hilbert morphism only makes sense over $\mathbb{C}$.

There have been several attempts to construct a local version of such a stack. In [An1] Andronikof defined a prestack on P^*X and announced the microlocal Riemann-Hilbert correspondance on the stalks. However, at this time there did not exist tools to define a global microlocal Riemann-Hilbert morphism. Another topological construction was proposed in [GMV], but to our knowledge this project has neither been completed nor published.

Our approach makes use of the theory of analytic ind-sheaves, recently introduced in [KS2] by Kashiwara and Schapira. Hence, microlocal perverse sheaves on a $\mathbb{C}^\times$ -conic open subset $U \subset T^*X$ will be ind-sheaves (or more precisely objects of the derived category of ind-sheaves) on U contrary to the construction of [An1] in which microlocal perverse sheaves on $U \subset T^*X$ were represented by complexes of sheaves on the base space X . The theory of ind-sheaves provides us with a nice representative of the stack associated to the prestack of [An1] and allows us to use the machinery developped in [KS2]. The essential tool in this description is Kashiwara's functor of ind-microlocalization $\mu : D^b(k_X) \rightarrow D^b(I(k_{T^*X}))$ of [K5]. This functor enables us to define explicitly a global Riemann-Hilbert morphism when $k = \mathbb{C}$.

In the future, we will hopefully show that we can actually patch (a twisted version of) this stack on a complex contact manifold and prove the Riemann-Hilbert theorem in the complex case.

In more detail, the contents of this paper are as follows.

Section 2 gives a criterion for a subprestack of the prestack of derived categories of ind-sheaves on a manifold to be a stack. It is a generalization of a proof of [KS1] showing that the prestack of perverse sheaves is a substack of the derived category of sheaves with $\mathbb{C}$ -constructible cohomology. Then we investigate abelian stacks on a topological spaces. Roughly speaking, an additive stack on a topological space is abelian if and only if its stalks are abelian categories and we have a "lifting property" for kernels and cokernels. This will be applied in Section 7 to construct the abelian stack of microlocal perverse sheaves as a substack of the prestack of derived ind-sheaves on the cotangent bundle of a complex manifold.

Section 3 first recalls the theory of microlocalization of [KS1] on a real manifold X . We do not review in detail the theory of the micro-support of sheaves but concentrate on the definition of the microlocal category $D^b(k_X, S)$ where $S \subset T^*X$ is an arbitrary subset. It is defined as the localization of the category $D^b(k_X)$ by the objects $\mathcal{F} \in D^b(k_X)$ whose micro-support does not intersect S . For any $\mathcal{F}, \mathcal{G} \in D^b(k_X)$ we get a natural morphism

$$\mathrm{Hom}_{D^b(k_X, S)}(\mathcal{F}, \mathcal{G}) \longrightarrow H^0(S, \mu \mathrm{hom}(\mathcal{F}, \mathcal{G})).$$

In the case where $S = \{pt\}$ the category $D^b(k_X, S)$ has been intensively studied in [KS1], and in particular it is proved that the morphism above is an isomorphism.

We will show that this result is still valid in the category $D^b(k_X, \{x\} \times \dot{\delta})$ where x is a point of X , $\delta \subset T_x^*X$ a closed cone and $\dot{\delta} = \delta \setminus \{0\}$. Later we will be mainly interested in the case where δ is a complex line. The main tool is the refined microlocal cut-off lemma

for non-convex sets, which we recall adding a few comments. We will also need the cut-off functor in Section 6.

Section 4 extends the definitions and results of Section 3 first to $\mathbb{R}$ -constructible then to $\mathbb{C}$ -constructible sheaves. There are two natural ways to define the microlocalization of the derived category of $\mathbb{R}$ -constructible sheaves. We either localize the category $D_{\mathbb{R}\text{-c}}^b(k_X)$ by sheaves whose micro-support does not intersect S or we take the full subcategory of $D^b(k_X, S)$ whose objects are represented by $\mathbb{R}$ -constructible sheaves. Following [An2] we will use the first definition. One important question is whether or not the two definitions coincide. The main result of this section is that this is the case when $S = \{x\} \times \hat{\delta}$.

Now suppose that X is a complex manifold. Recall that $\mathbb{C}$ -constructible sheaves may be defined by a microlocal property: an object $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(k_X)$ is $\mathbb{C}$ -constructible if and only if its micro-support is $\mathbb{C}^\times$ -conic. It is then natural to define a microlocally complex constructible sheaf as a microlocal $\mathbb{R}$ -constructible sheaf whose micro-support is $\mathbb{C}^\times$ -conic on S .

In **Section 5** we show that the constructions of Section 4 are locally “invariant under quantized contact transformations”.

Section 6 is devoted to the study of microlocally $\mathbb{C}$ -constructible sheaves in the category $D^b(k_X, \mathbb{C}^\times p)$. In Section 5 we have shown that the category $D_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times p)$ is invariant by quantized contact transformation. Hence we are reduced to study microlocally $\mathbb{C}$ -constructible sheaves in generic position, i.e. complexes of sheaves whose micro-support is contained in T_Z^*X for a complex (not necessarily smooth) hypersurface Z in a neighborhood of p . We give a complete proof that microlocally $\mathbb{C}$ -constructible sheaves in generic position may be represented by $\mathbb{C}$ -constructible sheaves (as announced in [An2]).

Following [An1], we define in **Section 7** the category of microlocal perverse sheaves as a full subcategory of $D_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times p)$. An object $\mathcal{F} \in D_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times p)$ is perverse if for any non-singular point $q \in \text{SS}(\mathcal{F})$ in a neighborhood of $\mathbb{C}^\times p$ the complex $\mathcal{F}$ is isomorphic in $D^b(k_X, \mathbb{C}^\times q)$ to a constant sheaf $M_Y[d_Y]$ supported on a closed submanifold $Y \subset X$. This definition is natural in view of the microlocal characterization of perverse sheaves of [KS1] and also leads to definition of a prestack of microlocal perverse sheaves on P^*X . Then we prove that the category $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$ is abelian as has been announced in [An1]. Our proof gives a refined result which allows us to conclude that the stack associated to this prestack is abelian. This stack is the stack of microlocal perverse sheaves on P^*X . However we will need a more explicit representation in order to define the microlocal Riemann-Hilbert correspondance.

In **Section 8** we finally define microlocal perverse sheaves as objects of the derived category of ind-sheaves on conic open subsets of T^*X . In Section 7 we have constructed the category of microlocal perverse sheaves at any $p \in P^*X$ (or on $\mathbb{C}^\times p \subset \dot{T}^*X$) which will be equivalent to the stalk of the stack μPerv of microlocal perverse sheaves. The idea of the construction of μPerv is to use the fact Kashiwara’s functor μ of ind-microlocalization induces a fully faithful functor from $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$ into the stalk of the prestack of bounded derived categories of ind-sheaves on $\mathbb{C}^\times$ -conic subsets of T^*X . Then we can define a microlocal perverse sheaf on a conic open subset $U \subset T^*X$ as an object of $D^b(I(k_U))$ that is isomorphic to a microlocal perverse sheaf of $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$ at any point of $p \in U$. Finally, we show that the stack of microlocal perverse sheaves is canonically equivalent to the stack associated to the prestack of the last section.

In **Section 9** we establish the microlocal Riemann-Hilbert correspondance:

$$\mu\text{Perv}(\Omega) \begin{array}{c} \xrightarrow{\mu\text{RH}} \\ \xleftarrow{\mu\text{Sol}} \end{array} \mathcal{HolReg}(\mathcal{E}_X|_{\gamma^{-1}\Omega}).$$

It is defined by the formulas

$$\mu\mathrm{Sol}(\mathcal{M}) = \mathrm{R}\mathcal{H}om_{\beta(\mathcal{E}_X|_{\gamma^{-1}\Omega})}(\beta(\mathcal{M}), \mu\mathcal{O}_X|_{\gamma^{-1}\Omega})$$

$$\mu\mathrm{RH}(\mathcal{F}) = \gamma_\Omega^{-1} \mathrm{R}\gamma_{\Omega*}(\mathrm{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}^t|_{\gamma^{-1}(\Omega)}))$$

where γ_Ω is the restriction of the natural map $\gamma : \dot{T}^*X \rightarrow P^*X$ to $\gamma^{-1}\Omega$, $\mathrm{R}\mathcal{H}om, \beta$ are well-known functors from the theory of ind-sheaves (cf. [KS2]) and the “ring” $\mathcal{O}^t \in \mathrm{D}^b(\mathrm{I}(k_X))$ is the complex of tempered holomorphic functions on X of loc. cit.

Appendix A gives a short introduction to stacks with emphasis on the special properties resulting from the fact that we work on a topological space.

In **Appendix B** we give a short summary of the properties of Kashiwara’s functor μ .

First of all, I thank my thesis director P. Schapira both for having suggested this subject to me, and for always having been ready with precious help, guidance and encouragement throughout the last three years. Secondly, my gratitude goes out to M. Kashiwara with whom I had many invaluable conversations. I would particularly like to thank him for having shared with me his unpublished work on the microlocalization of ind-sheaves which provided me with certain key ideas on which this paper is based. It goes without saying, of course, that I could never have been able to complete this work without either of them. Finally, I would like to thank A. D’Agnolo, P. Polesello, F. Ivorra and D.-C. Cisinski for many useful discussions.

2 Abelian substacks of a prestack

Perverse sheaves on a complex manifold X are local objects - they form an abelian stack which is a subprestack of the prestack of (derived) sheaves on X (see [BBD] for the general theory of perverse sheaves, see also [KS1], Chapter X, for a microlocal approach to perverse sheaves). In Section 2.1 we will generalize the method used in [KS1] in order to prove that this subprestack is actually a stack. In particular we will show that a similar method can be applied to find substacks of the prestack of (derived) ind-sheaves. The abelian structure of the stack of perverse sheaves is defined by a t -structure on the triangulated prestack of derived categories of sheaves with $\mathbb{C}$ -constructible cohomology. However, the category of microlocal perverse sheaves will not be defined as the heart of a t -structure. Our strategy is based on the idea that a stack is “almost” abelian, if its stalks are abelian categories. Roughly speaking, an additive stack is abelian if and only if its stalks are abelian and kernels and cokernels can be lifted to small neighborhoods. We will investigate this statement more precisely in Sections 2.2 and 2.3.

2.1 A criterion for substacks

The basic definitions from the theory of stacks (on a topological space) are recalled in Appendix B. The results on proper stacks and ind-sheaves that we will use can be found in [KS2].

Definition 2.1.1. *Consider a prestack $\mathcal{C}$ on a topological space X . We say that a full subprestack $\mathcal{C}' \subset \mathcal{C}$ is **defined by a local property** (with respect to $\mathcal{C}$) if the following conditions are satisfied:*

- (i) *the prestack $\mathcal{C}'$ is stable by isomorphisms, i.e. if $U \subset X$ is open, $A \in \mathrm{Ob} \mathcal{C}'(U)$ then any object $B \in \mathrm{Ob} \mathcal{C}(U)$ isomorphic to A is also an object of $\mathrm{Ob} \mathcal{C}'(U)$,*
- (ii) *if $U \subset X$ is open and $A \in \mathrm{Ob} \mathcal{C}(U)$ then $A \in \mathrm{Ob} \mathcal{C}'(U)$ if and only if there is an open covering $U = \bigcup_{i \in I} U_i$ such that $A|_{U_i} \in \mathrm{Ob} \mathcal{C}'(U_i)$ for all $i \in I$.*

Remark 2.1.2. Consider a full subprestack $\mathcal{C}' \subset \mathcal{C}$ and a point $p \in X$. Then the natural functor $\mathcal{C}'_p \rightarrow \mathcal{C}_p$ is fully faithful. Therefore the (full) subprestack $\mathcal{C}'$ is defined by a local property if and only if for any object $A \in \text{Ob } \mathcal{C}(U)$ the statements (a) and (b) below are equivalent:

- (a) $A \in \text{Ob } \mathcal{C}'(U)$.
- (b) For every $p \in X$ the object A is in the essential image of the functor $\mathcal{C}'_p \rightarrow \mathcal{C}_p$, i.e. there exists an object $B \in \text{Ob } \mathcal{C}'_p$ such that A is isomorphic to B in $\mathcal{C}_p$.

Lemma 2.1.3. Let $\mathcal{C}$ be a triangulated prestack. Assume moreover that

- (1) for any $V \subset U$ the restriction functor i_{VU}^{-1} has a fully faithful left adjoint $i_{VU}!$,
- (2) these functors satisfy the base change theorem, i.e. for any Cartesian square of open subsets

$$\begin{array}{ccc} U_{12} & \longrightarrow & U_1 \\ \downarrow & \square & \downarrow \\ U_2 & \longrightarrow & V \end{array}$$

we have $i_{U_{12}U_2}!i_{U_{12}U_1}^{-1} \simeq i_{U_2V}^{-1}i_{U_1V}!$, where $U_{12} = U_1 \cap U_2$.

Consider the union of two open sets $U = U_1 \cup U_2$ and suppose that we are given

- (i) objects $A_1 \in \text{Ob } \mathcal{C}(U_1)$ and $A_2 \in \text{Ob } \mathcal{C}(U_2)$,
- (ii) an isomorphism $f_{21} : A_1|_{U_{12}} \xrightarrow{\sim} A_2|_{U_{12}}$ in $\mathcal{C}(U_{12})$.

Then there exist an object $A \in \text{Ob } \mathcal{C}(U)$ and isomorphisms $f_1 : A|_{U_1} \xrightarrow{\sim} A_1$, $f_2 : A|_{U_2} \xrightarrow{\sim} A_2$ that are compatible with f_{21} on U_{12} , i.e. the following diagram commutes:

$$\begin{array}{ccc} A_1|_{U_{12}} & \xrightarrow[f_{21}]{\sim} & A_2|_{U_{12}} \\ & \nwarrow \sim \quad \nearrow \sim & \\ & A|_{U_{12}} & \end{array}$$

Proof. Let us simplify the notations for the restriction functors by suppressing U . Hence we have the functors

$$\begin{array}{lll} i_1^{-1} : \mathcal{C}(U) \rightarrow \mathcal{C}(U_1) & i_2^{-1} : \mathcal{C}(U) \rightarrow \mathcal{C}(U_2) & i_{12}^{-1} : \mathcal{C}(U) \rightarrow \mathcal{C}(U_{12}) \\ i_{12,1}^{-1} : \mathcal{C}(U_1) \rightarrow \mathcal{C}(U_{12}) & i_{12,2}^{-1} : \mathcal{C}(U_2) \rightarrow \mathcal{C}(U_{12}) & \end{array}$$

We use a similar notation with lower-case symbol $!$ for the left adjoints of these functors. Now define A by choosing a distinguished triangle

$$i_{12}!(A_1|_{U_{12}}) \xrightarrow{\begin{pmatrix} g_1 \\ g_2 \end{pmatrix}} i_1!A_1 \oplus i_2!A_2 \xrightarrow{(h_1, h_2)} A \xrightarrow{+}$$

where the first morphism defining the triangle is given by

$$\begin{aligned} g_1 : i_{12}!(A_1|_{U_{12}}) &\xrightarrow{\sim} i_{1!}i_{12,1}!(A_1|_{U_{12}}) \longrightarrow i_{1!}(A_1), \\ g_2 : i_{12}!(A_1|_{U_{12}}) &\xrightarrow[\sim]{-f_{21}} i_{12}!(A_2|_{U_{12}}) \xrightarrow{\sim} i_{2!}i_{12,2}!(A_2|_{U_{12}}) \longrightarrow i_{2!}(A_2). \end{aligned}$$

¹Recall that $i_{VU}!$ is fully faithful if and only if the adjunction morphism $\text{Id} \rightarrow i_{VU}!i_{VU}^{-1}$ is an isomorphism. Also note that for any three open subsets $W \subset V \subset U$ the isomorphism $i_{WV}^{-1}i_{VU}^{-1} \simeq i_{WU}^{-1}$ induces an isomorphism $i_{WV}!i_{VU}! \simeq i_{WU}!$.

Define f_1 and f_2 by

$$\begin{aligned} f_1 : A_1 &\simeq i_1^{-1} i_{1!} A_1 \xrightarrow{h_1|_{U_1}} A|_{U_1}, \\ f_2 : A_2 &\simeq i_2^{-1} i_{2!} A_2 \xrightarrow{h_2|_{U_2}} A|_{U_2}. \end{aligned}$$

Let us check that f_1 is an isomorphism. It is sufficient to show that $g_2|_{U_1}$ is an isomorphism. Since by the base change theorem we have

$$i_1^{-1} i_{2!}(A_2) \simeq i_{12,1!}(A_2|_{U_{12}}) \simeq i_1^{-1} i_{12!}(A_2|_{U_{12}}),$$

we get the result. A similar argument shows that f_2 is an isomorphism. To prove that these isomorphisms are compatible with f_{21} one uses the fact that $h_1 g_1 + h_2 g_2 = 0$. $\square$

Remark 2.1.4. In the situation of the preceding Lemma 2.1.3, suppose that we are given a full (but not necessarily triangulated) subprestack $\mathcal{C}' \subset \mathcal{C}$ that is defined by a local property. Then the lemma holds in $\mathcal{C}'$, i.e. if the objects A_1, A_2 are in $\mathcal{C}'$ then the object A lies also in $\mathcal{C}'$. Indeed, we may patch the given objects A_1, A_2 of $\mathcal{C}'$ to an object A in the prestack $\mathcal{C}$ using Lemma 2.1.3. Then the axioms (cf. Definition 2.1.1) immediately imply that A is an object of $\mathcal{C}'$.

Note that if moreover $\mathcal{C}'$ is separated, then the object A is unique up to unique isomorphism².

Let us apply Lemma 2.1.3 to the prestack of bounded derived categories of ind-sheaves. Denote by $D^b(I(k_*))$ the prestack $U \mapsto D^b(I(k_U))$ on a locally compact space X with a countable base of open sets. This prestack has the following properties:

- (i) it is a triangulated prestack,
- (ii) if $V \subset U \subset X$ are open subsets and if we denote as usual by $i_{VU}^{-1} : D^b(I(k_U)) \rightarrow D^b(I(k_V))$ the restriction functor, then this functor has a fully faithful left adjoint $Ri_{VU}!!$,
- (iii) the base change theorem is satisfied (cf. Lemma 2.1.3).

The proof is based on the fact that $I(k_*)$ is a proper stack (cf. [KS2]). Hence we get

Corollary 2.1.5. *Consider the prestack $D^b(I(k_*))$ on a locally compact topological space X and two open subsets $U_1, U_2 \subset X$. Set $U = U_1 \cup U_2$.*

Suppose that we are given the following data

- (1) *two objects $\mathcal{F}_1 \in \text{Ob } D^b(I(k_{U_1}))$, $\mathcal{F}_2 \in \text{Ob } D^b(I(k_{U_2}))$,*
- (2) *an isomorphism $f_{21} : \mathcal{F}_1|_{U_{12}} \xrightarrow{\sim} \mathcal{F}_2|_{U_{12}}$.*

Then there exist an object $\mathcal{F} \in \text{Ob } D^b(I(k_U))$ and isomorphisms $f_1 : \mathcal{F}|_{U_1} \xrightarrow{\sim} \mathcal{F}_1$, $f_2 : \mathcal{F}|_{U_2} \xrightarrow{\sim} \mathcal{F}_2$ compatible with f_{21} on U_{12} .

Now let us state Proposition 10.2.9. of [KS1] in a slightly more general context and change the proof so that we may adapt it later to the case of ind-sheaves.

²More precisely, if A' is another object with isomorphisms $f'_i : A'|_{U_i} \xrightarrow{\sim} A_i$ for $i \in \{1, 2\}$ such that $f_{21} \circ f'_1|_{U_{12}} = f'_2|_{U_{12}}$ then there exists a unique isomorphism $\varphi : A \xrightarrow{\sim} B$ such that $f'_i \circ \varphi|_{U_i} = f_i$ for $i \in \{1, 2\}$.

Proposition 2.1.6. *Let X be a locally compact paracompact space with a countable base of open sets. Consider a proper stack³ $\mathcal{A}$ such that for every open subset $U \subset X$ the category $\mathcal{A}(U)$ has enough injective objects. Denote by $D^b(\mathcal{A})$ the associated prestack of bounded derived categories.*

Let $\mathcal{C} \subset D^b(\mathcal{A})$ be a separated full subprestack that is defined by a local property. Then $\mathcal{C}$ is a stack.

Proof. We have to show that $\mathcal{C}$ satisfies the patching condition (cf. Definition A.4.2 in Appendix B). Since X is paracompact we have to verify the patching condition only for countable coverings.

Since the conditions of Lemma 2.1.3 are satisfied for the prestack $D^b(\mathcal{A})$ and $\mathcal{C}$ is defined by a local property, we know that $\mathcal{C}$ satisfies the patching condition for any covering of type $U = U_1 \cup U_2$.

Using the fact that $\mathcal{C}$ is separated, we can easily verify by induction that the patching condition is satisfied for finite coverings.

Therefore, using again the fact that $\mathcal{C}$ is separated, it is sufficient to prove that objects may be patched in $\mathcal{C}$ for open coverings of type $U = \bigcup_{n \in \mathbb{N}} U_n$ where $U_n \subset U_{n+1}$.

Consider a family of objects $A_n \in \text{Ob } D^b(\mathcal{A}(U_n))$ and isomorphisms $f_{n-1} : A_{n-1} \xrightarrow{\sim} A_n|_{U_{n-1}}$ (the other isomorphisms are uniquely determined by the cocycle condition).

Denote by $i_n : U_n \hookrightarrow U$ the inclusion map. Then we lift the morphisms of the system to

$$g_{n-1} : i_{n-1}!A_{n-1} \xrightarrow[\sim]{i_{n-1}!(f_{n-1})} i_{n-1}!A_n|_{U_{n-1}} \longrightarrow i_n!A_n$$

in $D^b(\mathcal{A}(U))$. Hence we get a family of morphisms $\{g_{n-1} : i_{n-1}!A_{n-1} \rightarrow i_n!A_n\}_{n \geq 1}$ in $D^b(\mathcal{A}(U))$. Note that $g_{n-1}|_{U_{n-1}}$ is an isomorphism by the base change theorem.

By hypothesis $\mathcal{A}(U)$ has enough injective objects and therefore we have an equivalence of categories $K^b(\text{Inj}(\mathcal{A}(U))) \simeq D^b(\mathcal{A}(U))$. Here $\text{Inj}(\mathcal{A}(U))$ denotes the full abelian subcategory of $\mathcal{A}(U)$ whose objects are the injective objects of $\mathcal{A}(U)$ and $K^b(\text{Inj}(\mathcal{A}(U)))$ is the triangulated category of bounded complexes of $\text{Inj}(\mathcal{A}(U))$ where morphisms of complexes are considered up to homotopy. Hence there exist objects I_n and morphisms $h_{n-1} : I_{n-1} \rightarrow I_n$ in $C^b(\text{Inj}(\mathcal{A}(U)))$ such that the diagram $\{h_{n-1} : I_{n-1} \rightarrow I_n\}_{n \geq 1}$ is isomorphic (in $D^b(\mathcal{A}(U))$) to the diagram $\{g_{n-1} : i_{n-1}!A_{n-1} \rightarrow i_n!A_n\}_{n \geq 1}$.

Let $A = \varinjlim I_n$ and consider A as an object in $D^b(\mathcal{A}(U))$. Then $A|_{U_n}$ is quasi-isomorphic in $C^b(\mathcal{A}(U))$ to I_n because for $m \geq n$ the morphism $I_n \rightarrow I_m|_{U_n}$ is a quasi-isomorphism. Hence there are natural isomorphisms $A|_{U_n} \simeq A_n$ in $D^b(\mathcal{A}(U))$, and a simple diagram chase shows that they are compatible with the morphisms f_n . $\square$

Remark 2.1.7. Note that in Proposition 2.1.6 the hypothesis that the categories $D^b(\mathcal{A}(U))$ possess enough injective objects can be weakened. During the proof, we actually only use the fact that any diagram in $D^b(\mathcal{A}(U))$ of type $\{A_n \rightarrow A_{n+1}\}_{n \geq 0}$ can be lifted to a diagram $\{I_n \rightarrow I_{n+1}\}_{n \geq 0}$ in $C^b(\mathcal{A}(U))$. We do not use the fact that the objects I_n are injective.

We need a proposition of [KS2] (part of Theorem 11.2.6).

Proposition 2.1.8. *Let $\mathcal{A}$ be an abelian category with a system of strict generators. Denote by $\text{Ind } \mathcal{A}$ the category of ind-objects of $\mathcal{A}$ and let $S \subset \text{Ob Ind}(\mathcal{A})$ be a small subset. Then there exists an essentially small⁴ full abelian subcategory $\mathcal{B} \subset \mathcal{A}$ such that*

³For the definition of a proper stack see [KS2]. A proper stack $\mathcal{A}$ and the associated prestack of bounded derived categories $D^b(\mathcal{A})$ satisfy the hypothesis of Lemma 2.1.3. Moreover for each open subset $U \subset X$, the abelian category $\mathcal{A}(U)$ admits filtered exact colimits and the restriction functors commute to such colimits.

⁴A category is essentially small if it is equivalent to a small category.

- (i) $\mathcal{B}$ is stable by subobject, quotient and extension in $\mathcal{A}$,
- (ii) $\text{Ind } \mathcal{B} \subset \text{Ind } \mathcal{A}$ is stable by subobject, quotient and extension and contains S ,
- (iii) $\text{Ind } \mathcal{B}$ has enough injectives.

Hence we get the following corollary.

Corollary 2.1.9. *Let $\mathcal{S}$ be a small diagram in $\text{D}^b(\text{Ind}(\mathcal{A}))$, i.e. $\mathcal{S} \subset \text{Mor } \text{D}^b \text{Ind}(\mathcal{A})$ is a set of morphisms. Then there exists an essentially small full abelian subcategory $\mathcal{B} \subset \mathcal{A}$ such that*

- (i) $\mathcal{B}$ is stable by subobject, quotient and extension in $\mathcal{A}$,
- (ii) $\text{Ind } \mathcal{B} \subset \text{Ind } \mathcal{A}$ is stable by subobject, quotient and extension,
- (iii) $\text{Ind } \mathcal{B}$ has enough injectives,
- (iv) $\mathcal{S}$ is contained in the image of the natural functor

$$\text{D}^b(\text{Ind}(\mathcal{B})) \longrightarrow \text{D}^b(\text{Ind}(\mathcal{A})).$$

Proof. Every morphism $(f : A \rightarrow A') \in \mathcal{S}$ may be represented by a diagram

$$A \longrightarrow A'' \longleftarrow A'.$$

Chose such a diagram for every $f \in \mathcal{S}$. Consider the set of objects of $\text{Ind } \mathcal{A}$ appearing in some complex in some diagram and apply Proposition 2.1.8 to this set. $\square$

Combining Proposition 2.1.6 with Remark 2.1.7 and Corollary 2.1.9 we get:

Theorem 2.1.10. *Let X be a paracompact locally compact topological space with a countable base of open sets and consider a separated full subprestack $\mathcal{C} \subset \text{D}^b(I(k_*))$ that is defined by a local property. Then $\mathcal{C}$ is a stack.*

2.2 Limits and colimits in stacks

Recall that if $\mathcal{C}$ is a prestack on a topological space X we denote by ρ_{VU} the restriction functor for two open subsets $V \subset U \subset X$ and $\rho_p^U : \mathcal{C}(U) \rightarrow \mathcal{C}_p$ the canonical functor into the stalk at p .

Definition 2.2.1. *Let $\mathcal{I}$ be a small category. We say that $\mathcal{C}$ admits limits (resp. colimits) indexed by $\mathcal{I}$ if for every open subset $U \subset X$ the category $\mathcal{C}(U)$ admits limits (resp. colimits) indexed by $\mathcal{I}$ such that the restriction functors commute to these limits (resp. colimits).*

Let $\mathcal{I}$ be a finite category. It is easy to see that if $\mathcal{C}$ admits limits (resp. colimits) indexed by $\mathcal{I}$, then for every $p \in X$ the category $\mathcal{C}_p$ admits limits (resp. colimits) indexed by $\mathcal{I}$ and the functor ρ_p^U commutes to such limits (resp. colimits).

However, the converse is not true. We cannot know simply by looking at the stalks whether or not a prestack admits limits or colimits indexed by $\mathcal{I}$ (even if $\mathcal{C}$ is a stack).

If $\mathcal{C}$ is separated we can at least see from the stalks whether or not a given object represents a limit or colimit indexed by a finite category. By duality we only need to consider the case of finite colimits.

Lemma 2.2.2. *Let $\mathcal{C}$ be a separated prestack on a topological space X .*

Consider a finite category $\mathcal{I}$, an open subset $U \subset X$ and a functor $\alpha : \mathcal{I} \rightarrow \mathcal{C}(U)$.

Suppose given an object $L \in \text{Ob } \mathcal{C}(U)$ and morphisms $\sigma_i : \alpha(i) \rightarrow L$ such that for any morphism $s : i \rightarrow j$ of $\text{Mor } \mathcal{I}$ we have $\sigma_j \circ \alpha(s) = \sigma_i$.

Then the two following assertions are equivalent:

(i) *$(L, \{\sigma_i\}_{i \in \mathcal{I}})$ is a colimit of α in $\mathcal{C}(U)$ and for any open subset $V \subset U$ the pair $(L|_V, \{\sigma_i|_V\}_{i \in \mathcal{I}})$ is a colimit of $\rho_{VU}\alpha$ in $\mathcal{C}(V)$.*

(ii) *$(L, \{(\sigma_i)_p\}_{i \in \mathcal{I}})$ is a colimit of $\rho_p^U \alpha$ in $\mathcal{C}_p$ for all $p \in U$.*

Proof. Let $V \subset U$ be an open subset.

The object $L \in \text{Ob } \mathcal{C}(U)$ and the morphisms $\sigma_i|_V$ define a natural morphism of sheaves ($\mathcal{C}$ is separated) for any object $A \in \text{Ob } \mathcal{C}(V)$

$$\mathcal{H}\text{om}_{\mathcal{C}|_V}(L|_V, A) \longrightarrow \varprojlim_{i \in \mathcal{I}} \mathcal{H}\text{om}_{\mathcal{C}|_V}(\alpha(i)|_V, A) \quad (2.2.1)$$

Since $\mathcal{I}$ is a finite category we have for every $p \in V$

$$(\varprojlim_{i \in \mathcal{I}} \mathcal{H}\text{om}_{\mathcal{C}|_V}(\alpha(i)|_V, A))_p \simeq \varprojlim_{i \in \mathcal{I}} \mathcal{H}\text{om}_{\mathcal{C}|_V}(\alpha(i)|_V, A)_p.$$

Hence the morphism (2.2.1) induces in the stalks

$$\mathcal{H}\text{om}_{\mathcal{C}_p}(L, A) \simeq \mathcal{H}\text{om}_{\mathcal{C}|_V}(L, A)_p \longrightarrow \varprojlim_{i \in \mathcal{I}} \mathcal{H}\text{om}_{\mathcal{C}|_V}(\alpha(i)|_V, A)_p \quad (2.2.2)$$

Assertion (i) is clearly equivalent to the fact that the morphism (2.2.1) is an isomorphism for all $V \subset U$ and any $A \in \mathcal{C}(V)$.

Assertion (ii) is equivalent to the fact that the morphism (2.2.2) is an isomorphism for all $V \subset U$, $A \in \mathcal{C}(V)$ and $p \in V$.

Since $\mathcal{C}$ is separated the morphism (2.2.1) is an isomorphism if and only if for every $p \in V$ the morphism (2.2.2) is an isomorphism, which proves the lemma. $\square$

Remark 2.2.3. Consider a prestack $\mathcal{C}$, $\eta^\dagger : \mathcal{C} \rightarrow \mathcal{C}^\dagger$ the natural functor into the associated separated prestack (resp. $\eta^\ddagger : \mathcal{C} \rightarrow \mathcal{C}^\ddagger$ the natural functor into the associated stack), $\mathcal{I}$ a finite category and $\alpha : \mathcal{I} \rightarrow \mathcal{C}(U)$ a functor.

Suppose that there is an object $L \in \mathcal{C}(U)$ and morphisms $\sigma_i : \alpha(i) \rightarrow L$ such that condition (ii) of the Lemma 2.2.2 is verified.

Then Lemma 2.2.2 immediately implies that $(\eta^\dagger(L)|_V, \{\eta^\dagger(\sigma_i)|_V\})$ (resp. $(\eta^\ddagger(L)|_V, \{\eta^\ddagger(\sigma_i)|_V\})$) is a colimit of $\rho_{VU}^\dagger \eta^\dagger \alpha$ (resp. $\rho_{VU}^\ddagger \eta^\ddagger \alpha$) in $\mathcal{C}^\dagger(V)$ (resp. $\mathcal{C}^\ddagger(V)$) for all $V \subset U$.

Now suppose that we are given a stack $\mathcal{C}$, a finite category $\mathcal{I}$ and a functor $\alpha : \mathcal{I} \rightarrow \mathcal{C}(U)$. In order to check that there exists a colimit of α in $\mathcal{C}(U)$ we can apply Lemma 2.2.2. However, in practical situations (as in Section 7.2) it is often difficult to establish the existence of an object L defined on U that verifies condition (ii) of Lemma 2.2.2. Therefore we will use a refinement of Lemma 2.2.2 adapted to stacks which states that it is sufficient to prove the existence of the object L locally on U .

Proposition 2.2.4. *Let $\mathcal{C}$ be a stack on a topological space X and $\mathcal{I}$ be a finite category. Suppose that for every open subset $U \subset X$ and every functor $\alpha : \mathcal{I} \rightarrow \mathcal{C}(U)$ there exists an open covering $U = \bigcup_{j \in J} U_j$, objects $L_j \in \text{Ob } \mathcal{C}(U_j)$ and morphisms $\sigma_i^j : \alpha(i)|_{U_j} \rightarrow \mathcal{C}(U_j)$ verifying condition (ii) of Lemma 2.2.2.*

Then $\mathcal{C}$ admits colimits indexed by $\mathcal{I}$.

Proof. Consider an open subset $U \subset X$ and a functor $\alpha : \mathcal{I} \rightarrow \mathcal{C}(U)$. By hypothesis and Lemma 2.2.2 there exists an open covering $U = \bigcup_{j \in J} U_j$ such that $\rho_{U_j} \alpha$ is representable in $\mathcal{C}(U_j)$ by an object L_j and the restriction to any smaller open subset $W \subset U_j$ commutes to these colimits. The conditions clearly imply that we may patch together the colimits $L_j \in \text{Ob } \mathcal{C}(U_j)$ to an object $L \in \text{Ob } \mathcal{C}(U)$. Now applying again Lemma 2.2.2 we see that L is a colimit of α and that all restrictions commute to this colimit. $\square$

Corollary 2.2.5. *Let $\mathcal{C}$ be a prestack, $\eta : \mathcal{C} \rightarrow \mathcal{C}^\dagger$ the natural functor into the associated stack and $\mathcal{I}$ be a finite category.*

Suppose that the stalks of $\mathcal{C}$ admit colimits indexed by $\mathcal{I}$.

Moreover we assume that for any open subset $U \subset X$ and any functor $\alpha : \mathcal{I} \rightarrow \mathcal{C}(U)$ the following statement holds:

For any point $p \in U$ there exists an open neighborhood $U_p \subset U$, an object $L^p \in \mathcal{C}(U_p)$ and morphisms $\sigma_i^p : \alpha(i)|_{U_p} \rightarrow L^p$ such that condition (ii) of Lemma 2.2.2 is verified.

Then $\mathcal{C}^\dagger$ admits colimits indexed by $\mathcal{I}$.

Proof. Let $\alpha : \mathcal{I} \rightarrow \mathcal{C}^\dagger(U)$ be a functor. Consider the functors $\rho_p^{\dagger U} \alpha$ for all $p \in X$. Since $\mathcal{I}$ is finite there exists an open neighborhood U_p of p such that $\rho_p^{\dagger U} \alpha$ factors through $\mathcal{C}(U_p)$. Hence we can apply Proposition 2.2.4. $\square$

In particular we get the much weaker statement that if a prestack $\mathcal{C}$ admits colimits indexed by a finite category $\mathcal{I}$ then $\mathcal{C}^\dagger$ admits colimits indexed by $\mathcal{I}$.

2.3 A criterion for abelian stacks

We can apply the results of the last paragraph to additive prestacks with abelian stalks. First recall that if $\mathcal{C}$ is an additive prestack then $\mathcal{C}^\dagger$ is additive.

Theorem 2.3.1. *Let $\mathcal{C}$ be an additive prestack with abelian stalks. Suppose that for every $p \in X$ and every morphism $f : A \rightarrow B$ in $\mathcal{C}_p$ there exists an open neighborhood U of p such that f may be represented by a morphism $\tilde{f} : \tilde{A} \rightarrow \tilde{B}$ in $\mathcal{C}(U)$ and there are morphisms $K \rightarrow \tilde{A}$, $\tilde{B} \rightarrow K'$ such that $K \rightarrow \tilde{A}$ is a kernel in $\mathcal{C}_q$ and $\tilde{B} \rightarrow K'$ is a cokernel in $\mathcal{C}_q$ for any $q \in U$.*

Then $\mathcal{C}^\dagger$ is an abelian stack.

Proof. Clearly the conditions of Proposition 2.2.4 and Corollary 2.2.5 are satisfied for cokernels and kernels. Hence $\mathcal{C}^\dagger$ admits cokernels and kernels.

Let $f : A \rightarrow B$ be a morphism of $\mathcal{C}^\dagger(U)$ and consider the natural morphism $\text{coim } f \rightarrow \text{im } f$. Since the categories of germs are abelian this morphism is an isomorphism in the stalks. Since $\mathcal{C}^\dagger$ is separated it is also an isomorphism in $\mathcal{C}^\dagger(U)$. $\square$

Corollary 2.3.2. *Let $\mathcal{C}$ be an additive stack on X such that all stalks are abelian categories. Then $\mathcal{C}$ is an abelian stack if and only if for every morphism $f : A \rightarrow B$ in $\mathcal{C}_p$ there is an open neighborhood U of p such that f may be represented by a morphism $\tilde{f} : \tilde{A} \rightarrow \tilde{B}$ in $\mathcal{C}(U)$ and there are morphisms $K \rightarrow \tilde{A}$, $\tilde{B} \rightarrow K'$ such that $K \rightarrow \tilde{A}$ is a kernel in $\mathcal{C}_q$ and $\tilde{B} \rightarrow K'$ is a cokernel in $\mathcal{C}_q$ for any $q \in U$.*

Remark 2.3.3. Hence in order to verify that an additive stack is abelian it is enough to verify that its stalks are abelian categories and that “kernels and cokernels are constant in some neighborhood”.

3 Microlocalization of sheaves

3.1 Notations

Let $\mathbb{R}^+$ denote the group of strictly positive real numbers and $\mathbb{C}^\times$ the group of non-zero complex numbers.

We will mainly work on a fixed complex manifold⁵ X of complex dimension $\dim_{\mathbb{C}} X = d_X$. Let T^*X be its cotangent bundle and T_X^*X the zero section. Set $\dot{T}^*X = T^*X \setminus T_X^*X$ and let $P^*X = \dot{T}^*X/\mathbb{C}^\times$ be the projective cotangent bundle. We denote by

$$\gamma : \dot{T}^*X \longrightarrow P^*X$$

the natural map.

If $\Lambda \subset \dot{T}^*X$ is a subset, we define the antipodal set Λ^a as

$$\Lambda^a = \{(x; \xi) \mid (x; -\xi) \in \Lambda\},$$

and we set

$$\mathbb{R}^+\Lambda = \{(x; \xi) \in \dot{T}^*X \mid \exists \alpha \in \mathbb{R}^+ (x; \alpha\xi) \in \Lambda\}.$$

We define similarly $\mathbb{C}^\times\Lambda$. Hence $\mathbb{C}^\times\Lambda = \gamma^{-1}\gamma(\Lambda)$. If $\Lambda = \{p\}$ is a point, we will write $\mathbb{C}^\times p$ instead of $\mathbb{C}^\times\{p\}$.

We say that a subset $\Lambda \subset \dot{T}^*X$ is $\mathbb{R}^+$ -conic (resp. $\mathbb{C}^\times$ -conic) if it is stable under the action of $\mathbb{R}^+$ (resp. $\mathbb{C}^\times$), i.e. if $\mathbb{R}^+\Lambda = \Lambda$ (resp. $\mathbb{C}^\times\Lambda = \Lambda$).

In the sequel, we will often deal with $\mathbb{R}^+$ -conic subsets that are only locally $\mathbb{C}^\times$ -conic. More precisely, a subset $\Lambda \subset \dot{T}^*X$ is called $\mathbb{C}^\times$ -conic at $p \in \dot{T}^*X$ if there exists an open neighborhood U of p such that $U \cap \mathbb{C}^\times\Lambda = U \cap \Lambda$. Note that this definition still makes sense if Λ is a germ of a subset at p . An open subset is always $\mathbb{C}^\times$ -conic at each $p \in U$.

Let $S \subset \dot{T}^*X$ be another subset, and suppose that Λ is defined on a germ of a neighborhood of S . Then we say that Λ is $\mathbb{C}^\times$ -conic on S if it is $\mathbb{C}^\times$ -conic at every point of S . Clearly this is equivalent to the statement that there exists an open neighborhood U of S such that $U \cap \mathbb{C}^\times\Lambda = U \cap \Lambda$. In particular, Λ is $\mathbb{C}^\times$ -conic on $\dot{T}^*X$ if and only if it is $\mathbb{C}^\times$ -conic. Finally we call the following easy topological lemma to the reader's attention.

Lemma 3.1.1. *Let $S \subset \dot{T}^*X$ be a $\mathbb{C}^\times$ -conic set and $U \supset S$ an $\mathbb{R}^+$ -conic open neighborhood. Then there exists a $\mathbb{C}^\times$ -conic open set V such that $S \subset V \subset U$.*

Now let us fix the conventions for sheaves. All sheaves considered here are sheaves of vector spaces over a given field k .

We will consider the following categories:

$D^b(k_X)$ is the derived category of bounded complexes of sheaves of k vector spaces,

$D_{\mathbb{R}\text{-c}}^b(k_X)$ is the full subcategory of $D^b(k_X)$ whose objects have $\mathbb{R}$ -constructible cohomology,

$D_{\mathbb{C}\text{-c}}^b(k_X)$ is the full subcategory of $D_{\mathbb{R}\text{-c}}^b(k_X)$ whose objects have $\mathbb{C}$ -constructible cohomology,

$\text{Perv}(k_X)$ is the full abelian subcategory of $D_{\mathbb{C}\text{-c}}^b(k_X)$ whose objects are perverse sheaves.

We will not recall the construction of these categories here, for more details see for instance [KS1].

⁵All manifolds (complex or real) in this paper are supposed to be finite dimensional with a countable base of open sets.

3.2 Microlocalization of sheaves

In this section we recall the construction and some properties of the microlocalization of $D^b(k_X)$ on a subset $S \subset T^*X$ ([KS1], Chapter VI) which we will then discuss from the (pre)stack-theoretical point of view. Note that all definitions and statements below which do not involve $\mathbb{C}^\times$ -conic subsets of T^*X are valid on a real manifold.

Recall that if $\mathcal{F} \in D^b(k_X)$ then one can associate to $\mathcal{F}$ a closed $\mathbb{R}^+$ -conic involutive subset $\text{SS}(\mathcal{F})$ of T^*X called the micro-support of $\mathcal{F}$. The theory of the micro-support can be found in [KS1]. It is the set of codirections in which $\mathcal{F}$ “does not propagate”. More precisely, a point $p \in T^*X$ is not a point of the micro-support if and only if there exists an open neighborhood U of p such that for any $x \in X$ and any real map ψ of class C^1 with $\psi(x) = 0$ and $(d\psi)_x \in U$ we have

$$R\Gamma_{\{x \mid \psi(x) \geq 0\}}(\mathcal{F})_x \simeq 0.$$

Let $S \subset T^*X$ be an arbitrary subset. Set

$$\mathcal{N}_S = \left\{ \mathcal{F} \in D^b(k_X) \mid \text{SS}(\mathcal{F}) \cap S = \emptyset \right\}.$$

The micro-support is invariant under the shift functor. Further, if

$$\mathcal{F}' \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}'' \longrightarrow \mathcal{F}'[1]$$

is a distinguished triangle in which $\mathcal{F}'$ and $\mathcal{F}$ are objects of $\mathcal{N}_S$, then it follows from the inclusion

$$\text{SS}(\mathcal{F}'') \subset \text{SS}(\mathcal{F}) \cup \text{SS}(\mathcal{F}')$$

that $\mathcal{F}''$ is also an object of $\mathcal{N}_S$. Hence $\mathcal{N}_S$ is a full triangulated subcategory of $D^b(k_X)$. Note that if $x \in S \cap T_X^*X$ then $\mathcal{F} \in \mathcal{N}_S$ implies $\mathcal{F} \simeq 0$ in a neighborhood of x .

Definition 3.2.1. *The microlocalization of $D^b(k_X)$ on S is the localization of the triangulated category $D^b(k_X)$ by the full triangulated subcategory $\mathcal{N}_S$, i.e.*

$$D^b(k_X, S) = D^b(k_X) / \mathcal{N}_S.$$

If $S = \{p\}$, we will write $D^b(k_X, p)$ for $D^b(k_X, \{p\})$.

Note that an object $\mathcal{F}$ in $D^b(k_X, S)$ is isomorphic to zero if and only if $\mathcal{F} \oplus \mathcal{F}[1] \in \mathcal{N}_S$ and since $\text{SS}(\mathcal{F} \oplus \mathcal{F}[1]) = \text{SS}(\mathcal{F})$ this is equivalent to $\text{SS}(\mathcal{F}) \cap S = \emptyset$.

A morphism $\mathcal{F} \rightarrow \mathcal{G}$ of $D^b(k_X)$ is called an isomorphism on S if it is an isomorphism in $D^b(k_X, S)$. This is equivalent to the existence of a distinguished triangle in $D^b(k_X)$

$$\mathcal{F} \longrightarrow \mathcal{G} \longrightarrow \mathcal{H} \xrightarrow{+}$$

with $\text{SS}(\mathcal{H}) \cap S = \emptyset$.

From this it follows easily that if $\mathcal{F} \xrightarrow{\sim} \mathcal{G}$ is an isomorphism in $D^b(k_X, S)$, then $\text{SS}(\mathcal{F}) \cap S = \text{SS}(\mathcal{G}) \cap S$. Hence the micro-support of $\mathcal{F} \in D^b(k_X, S)$ is well-defined in a germ of a neighborhood of S .

Let $\mathcal{F}, \mathcal{G} \in D^b(k_X, S)$. By definition we have

$$\text{Hom}_{D^b(k_X, S)}(\mathcal{F}, \mathcal{G}) \simeq \varinjlim_{\substack{\mathcal{F}' \xrightarrow{\sim} \mathcal{F} \\ \text{on } S}} \text{Hom}_{D^b(k_X)}(\mathcal{F}', \mathcal{G}).$$

We will have constant recourse to the following easy lemma.

Lemma 3.2.2. *Let $S \subset \dot{T}^*X$ be any subset. Consider a morphism $\mathcal{F} \rightarrow \mathcal{G}$ of $D^b(k_X)$ that is an isomorphism on S . Then there exists an $\mathbb{R}^+$ -conic open neighborhood U of S such that $\mathcal{F} \rightarrow \mathcal{G}$ is an isomorphism on U . In particular $\text{SS}(\mathcal{F}) \cap U = \text{SS}(\mathcal{G}) \cap U$. If moreover S is $\mathbb{C}^\times$ -conic, then we can choose U to be $\mathbb{C}^\times$ -conic.*

Proof. By hypothesis there exists a distinguished triangle in $D^b(k_X)$

$$\mathcal{F} \longrightarrow \mathcal{G} \longrightarrow \mathcal{H} \xrightarrow{+}$$

such that $\text{SS}(\mathcal{H}) \cap S = \emptyset$. Since $\text{SS}(\mathcal{H})$ is a closed $\mathbb{R}^+$ -conic subset of $\dot{T}^*X$, the set $U = \mathbb{C}\text{SS}(\mathcal{H})$ is an open and $\mathbb{R}^+$ -conic neighborhood of S such that $\text{SS}(\mathcal{H}) \cap U = \emptyset$. Now suppose that S is $\mathbb{C}^\times$ -conic. To prove the last statement we use the fact that every $\mathbb{R}^+$ -conic open neighborhood V of S contains a $\mathbb{C}^\times$ -conic open neighborhood of S . $\square$

Recall that to any $\mathcal{F}, \mathcal{G} \in D^b(k_X)$ we can associate the object $\mu\text{hom}(\mathcal{F}, \mathcal{G}) \in D^b(k_{T^*X})$ (see [KS1], Chapter IV). This complex satisfies

$$\text{supp}(\mu\text{hom}(\mathcal{F}, \mathcal{G})) \subset \text{SS}(\mathcal{F}) \cap \text{SS}(\mathcal{G}).$$

Therefore $\mu\text{hom}(\mathcal{F}, \mathcal{G})|_S$ is well-defined for $\mathcal{F}, \mathcal{G} \in D^b(k_X, S)$.

For an arbitrary subset S there is a natural morphism

$$\text{Hom}_{D^b(k_X, S)}(\mathcal{F}, \mathcal{G}) \longrightarrow H^0(S, \mu\text{hom}(\mathcal{F}, \mathcal{G})). \quad (3.2.1)$$

Let us recall its construction. For any two objects $\mathcal{F}_1, \mathcal{F}_2 \in D^b(k_X)$ we have a canonical isomorphism

$$\text{Hom}_{D^b(k_X)}(\mathcal{F}_1, \mathcal{F}_2) \simeq H^0(T^*X, \mu\text{hom}(\mathcal{F}_1, \mathcal{F}_2))$$

which defines a morphism

$$\text{Hom}_{D^b(k_X)}(\mathcal{F}_1, \mathcal{F}_2) \longrightarrow H^0(S, \mu\text{hom}(\mathcal{F}_1, \mathcal{F}_2)).$$

Now if $\mathcal{F}' \rightarrow \mathcal{F}$ is an isomorphism on S we get an induced isomorphism

$$H^0(S, \mu\text{hom}(\mathcal{F}, \mathcal{G})) \xrightarrow{\sim} H^0(S, \mu\text{hom}(\mathcal{F}', \mathcal{G})).$$

Thus we get morphisms

$$\text{Hom}_{D^b(k_X)}(\mathcal{F}', \mathcal{G}) \longrightarrow H^0(S, \mu\text{hom}(\mathcal{F}, \mathcal{G})).$$

which induce the morphism (3.2.1).

There is a well-known situation in which this morphism is an isomorphism ([KS1], Theorem 6.1.2).

Proposition 3.2.3. *Let $p \in T^*X$ and $\mathcal{F}, \mathcal{G} \in D^b(X, p)$. Then the morphism (3.2.1)*

$$\text{Hom}_{D^b(X, p)}(\mathcal{F}, \mathcal{G}) \longrightarrow H^0 \mu\text{hom}(\mathcal{F}, \mathcal{G})_p$$

is an isomorphism.

The idea is to calculate both sides by using microlocal cut-off functors. We will show that such a strategy works in the case of a closed cone in $\dot{T}_x^*X$, $x \in X$.

However, the morphism (3.2.1) is not an isomorphism in general (cf. [KS1], Exercise VI.6 which gives a counter-example on an open subset).

Remark 3.2.4. The correspondance

$$T^*X \supset U \mapsto D^b(k_X, U)$$

defines a prestack on T^*X , which we will usually denote by $D^b(k_X, *)$. Therefore $\gamma_* D^b(k_X, *)|_{\dot{T}^*X}$ defines a prestack on P^*X .

Proposition 3.2.5. *Let S be a subset of $\dot{T}^*X$. Then the natural functor*

$$\varinjlim_{\substack{S \subset U \subset \dot{T}^*X \\ U \text{ } \mathbb{R}^+-\text{conic}}} D^b(k_X, U) \longrightarrow D^b(k_X, S)$$

is an equivalence.

Proof. The functor is obviously essentially surjective. Let us show that it is fully faithful. Let $\mathcal{F}, \mathcal{G} \in D^b(k_X)$. By Lemma 3.2.2 we get

$$\begin{aligned} \text{Hom}_{\varinjlim_{\substack{S \subset U \subset \dot{T}^*X \\ U \text{ } \mathbb{R}^+-\text{conic}}} D^b(k_X, U)}(\mathcal{F}, \mathcal{G}) &\simeq \varinjlim_{\substack{S \subset U \subset \dot{T}^*X \\ U \text{ } \mathbb{R}^+-\text{conic}}} \text{Hom}_{D^b(k_X, U)}(\mathcal{F}, \mathcal{G}) \\ &\simeq \varinjlim_{\substack{S \subset U \subset \dot{T}^*X \\ U \text{ } \mathbb{R}^+-\text{conic}}} \varinjlim_{\substack{\mathcal{F}' \simeq \mathcal{F} \\ \text{on } U}} \text{Hom}_{D^b(k_X)}(\mathcal{F}', \mathcal{G}) \\ &\simeq \varinjlim_{S \subset U \subset \dot{T}^*X} \varinjlim_{\substack{\mathcal{F}' \simeq \mathcal{F} \\ \text{on } U}} \text{Hom}_{D^b(k_X)}(\mathcal{F}', \mathcal{G}) \simeq \text{Hom}_{D^b(k_X, S)}(\mathcal{F}, \mathcal{G}). \end{aligned}$$

□

Corollary 3.2.6. *Let S be a $\mathbb{C}^\times$ -conic subset of $\dot{T}^*X$ and $p \in T^*X$.*

(i) *The natural functor*

$$\varinjlim_{\substack{S \subset U \subset \dot{T}^*X \\ U \text{ } \mathbb{C}^\times\text{-conic}}} D^b(k_X, U) \longrightarrow D^b(k_X, S)$$

is an equivalence.

(ii) *The natural functor*

$$D^b(k_X, *)_p \longrightarrow D^b(k_X, p)$$

*is an equivalence. If moreover $p \in \dot{T}^*X$ then*

$$\gamma_*(D^b(k_X, *)|_{\dot{T}^*X})_{\gamma(p)} \simeq D^b(k_X, \gamma^{-1}\gamma(p)) = D^b(k_X, \mathbb{C}^\times p).$$

Proof. Any $\mathbb{R}^+$ -conic neighborhood of a $\mathbb{C}^\times$ -conic subset contains a $\mathbb{C}^\times$ -conic neighborhood. Thus part (i) of the corollary follows from a cofinality argument applied to the result of Proposition 3.2.5 and part (ii) follows from Proposition 3.2.5 and (i). □

3.3 Refined microlocal cut-off

Let us recall the basic idea of a microlocal cut-off functor.

Let X be a finite dimensional real vector space, $U \ni 0$ a relatively compact open neighborhood of 0 and consider an open cone $\gamma \subset X^*$.

Definition 3.3.1. A microlocal cut-off functor on $U \times \gamma$ is a functor

$$\Phi_{U,\gamma} : D^b(k_X) \rightarrow D^b(k_X)$$

such that

- (i) $SS(\Phi_{U,\gamma}\mathcal{F}) \subset X \times \overline{\gamma}$ (the micro-support has been cut off at $\overline{\gamma}$),
- (ii) $SS(\Phi_{U,\gamma}(\mathcal{F})) \cap U \times \gamma = SS(\mathcal{F}) \cap U \times \gamma$,
- (iii) $\Phi_{U,\gamma}$ is equipped with a morphism of functors $\alpha : \Phi_{U,\gamma} \rightarrow \text{Id}$ such that α induces an isomorphism in $D^b(k_X, U \times \gamma)$ which can be visualized by

$$\begin{array}{ccc} D^b(k_X) & \xrightarrow{\Phi_{U,\gamma}} & D^b(k_X) \\ \downarrow & & \downarrow \\ D^b(k_X, U \times \gamma) & \xrightarrow{\Phi_{U,\gamma} \simeq \text{Id}} & D^b(k_X, U \times \gamma). \end{array}$$

Note that condition (iii) implies (ii).

If the cut-off functor $\Phi_{U,\gamma}$ allows us to estimate the micro-support of $\Phi_{U,\gamma}(\mathcal{F})$ in the fiber $\{0\} \times X^*$, we usually call it a refined microlocal cut-off.

A cut-off functor is easily constructed in the case of a convex open cone (see Proposition 5.2.3 of [KS1]). A generalization to non-convex cones is stated in Exercise V.8 of [KS1] (for a proof see [D'A]). These tools will allow us in Section 3.4 to calculate sections of μhom along a complex line (or more generally along closed cones of T_x^*X where $x \in X$). The result will imply that the morphism (3.2.1) is an isomorphism in this case (see Section 3.4).

Later we will need to construct a functor $D_{\mathbb{C}\text{-c},\Lambda}^b(k_X, \mathbb{C}^\times p) \rightarrow D_{\mathbb{C}\text{-c}}^b(k_X, \pi(p))$ if Λ is in generic position at p (Section 6.1). For this purpose we will need the refined microlocal cut-off of [D'A]. It is an extension of the classical “refined microlocal cut-off lemma” (Proposition 6.1.4 of [KS1]) to non-convex cones with a good estimate for the micro-support.

Let us recall the cut-off functor of [KS1], Exercise V.8. Let X be a real, finite dimensional vector space, $\dot{X} = X \setminus \{0\}$, $U \subset X$ an open subset and $\gamma \subset X$ an open cone.

We have the following natural morphisms:

$$\begin{array}{ccccc} & & X \times X & & \\ & q_1 \swarrow & \downarrow \tilde{s} & \searrow q_2 & \\ X & & X & & X \end{array}$$

where q_1 and q_2 are the natural projections and

$$\begin{aligned} s : X \times X &\longrightarrow X & ; & (x, y) \mapsto x + y, \\ \tilde{s} : X \times X &\longrightarrow X & ; & (x, y) \mapsto x - y. \end{aligned}$$

We define the functor $\Phi_{U,\gamma}$ by setting for any $\mathcal{F} \in D^b(k_X)$

$$\Phi_{U,\gamma}(\mathcal{F}) = k_{\gamma^a}^\wedge * \mathcal{F}_U = \text{Rs}!(q_1^{-1}k_{\gamma^a}^\wedge \otimes q_2^{-1}\mathcal{F}_U).$$

It can be shown that $\Phi_{U,\gamma}$ is a microlocal cut-off functor in the sense of Definition 3.3.1.

Let us add two easy lemmas which will be useful in the next section.

Lemma 3.3.2. *Let $\mathcal{F}, \mathcal{G} \in \mathbf{D}^b(k_X)$. Then we have a canonical isomorphism*

$$\mathbf{R} s_!(q_1^{-1}\mathcal{F} \otimes q_2^{-1}\mathcal{G}) \simeq \mathbf{R} q_{1!}(\tilde{s}^{-1}\mathcal{F} \otimes q_2^{-1}\mathcal{G})$$

Proof. Consider the map

$$\varphi : X \times X \longrightarrow X \times X \quad ; \quad (x, y) \mapsto (x + y, y)$$

We obviously have the formulas:

$$\tilde{s} \circ \varphi = q_1 \quad q_1 \circ \varphi = s \quad q_2 \circ \varphi = q_2$$

Hence we get the chain of natural isomorphisms

$$\begin{aligned} \mathbf{R} s_!(q_1^{-1}\mathcal{F} \otimes q_2^{-1}\mathcal{G}) &\simeq \mathbf{R} q_{1!} \mathbf{R} \varphi_!(\varphi^{-1}\tilde{s}^{-1}\mathcal{F} \otimes \varphi^{-1}q_2^{-1}\mathcal{G}) \simeq \mathbf{R} q_{1!}(\mathbf{R} \varphi_!\varphi^{-1}\tilde{s}^{-1}\mathcal{F} \otimes q_2^{-1}\mathcal{G}) \\ &\simeq \mathbf{R} q_{1!}(\tilde{s}^{-1}\mathcal{F} \otimes q_2^{-1}\mathcal{G}). \end{aligned}$$

□

Lemma 3.3.3. *Let $\gamma_1, \gamma_2 \subset X$ be two open cones and $U \subset X$ open. Then there is a natural distinguished triangle*

$$\Phi_{U, \gamma_1 \cap \gamma_2}(\mathcal{F}) \longrightarrow \Phi_{U, \gamma_1}(\mathcal{F}) \oplus \Phi_{U, \gamma_2}(\mathcal{F}) \longrightarrow \Phi_{U, \gamma_1 \cup \gamma_2}(\mathcal{F}) \xrightarrow{+}$$

Proof. Follows immediately from the distinguished triangle

$$k_{\gamma_1^a \cap \gamma_2^a} \longrightarrow k_{\gamma_1^a} \oplus k_{\gamma_2^a} \longrightarrow k_{\gamma_1^a \cup \gamma_2^a} \xrightarrow{+}$$

by applying the triangulated functors $(\cdot)^\wedge$ and $(\cdot) * \mathcal{F}_U$. □

Next, we recall D'Agnolo's condition under which the cut-off functor $\Phi_{U, \gamma}$ is refined. These results will not be needed until Section 6.

Definition 3.3.4. *If $\gamma \subset X^*$ is an open cone, set*

$$\partial^\circ \gamma = \pi \chi(\mathrm{SS}(\mathbb{C}_\gamma) \setminus \{0; 0\})$$

where $\chi : T^*X^* \rightarrow T^*X$ is defined by $\chi(\xi; x) = (x; -\xi)$.

One says that (U, γ) is a refined cutting pair at 0, if $U \subset X$ is a relatively compact open neighborhood of 0 and for any $x \in \partial U \cap \partial^\circ \gamma$ there exists $\xi \in \dot{X}$ such that $N_x^*(U) = \mathbb{R}_{\geq 0}\xi$ and $\chi(\mathrm{SS}(k_\gamma)) \cap \pi^{-1}(x) = \mathbb{R}_{\leq 0}\xi$.

This allows one to give a good estimate of the micro-support of $\Phi_{U, \gamma}(\mathcal{F})$ at 0.

Proposition 3.3.5. *Let (U, γ) be a refined cutting pair at 0. Then*

$$\mathrm{SS}(\Phi_{U, \gamma}(\mathcal{F})) \cap \pi^{-1}(0) \subset \left\{ \xi \in \gamma \mid (0; \xi) \in \mathrm{SS}(\mathcal{F}) \right\} \cup \left\{ \xi \in \partial \gamma \mid \exists x \in \overline{U} : (x, \xi) \in \mathrm{SS}(\mathcal{F}) \right\}.$$

Let us add a useful corollary:

Corollary 3.3.6. *Let (U, γ) be a refined cutting pair at 0 and suppose that $\mathrm{SS}(\mathcal{F}) \cap (\overline{U} \times \partial \gamma) = \emptyset$. Then there exists an open neighborhood V of 0 such that*

$$\mathrm{SS}(\Phi_{U, \gamma}(\mathcal{F})) \cap \pi^{-1}(V) = \mathrm{SS}(\mathcal{F}) \cap (V \times \gamma).$$

Proof. Since $\text{SS}(\Phi_{U,\gamma}(\mathcal{F})) \subset X \times \overline{\gamma}$ and $\text{SS}(\Phi_{U,\gamma}(\mathcal{F})) \cap (U \times \gamma) = \text{SS}(\mathcal{F}) \cap (U \times \gamma)$, it is enough to show that $\text{SS}(\Phi_{U,\gamma}(\mathcal{F})) \cap V \times \partial\gamma = \emptyset$ for some neighborhood V of 0. D'Agnolo's estimate of the microsupport implies that this is at least true at 0.

Now suppose that such a neighborhood V does not exist. Then we can construct a sequence (x_n, ξ_n) such that $x_n \rightarrow 0$ and $\xi_n \in \text{SS}_{x_n}(\Phi_{U,\gamma}(\mathcal{F})) \cap \partial\gamma$. Since both sets are invariant by $\mathbb{R}_{\geq 0}$ we can assume that $|\xi_n| = 1$, hence by extracting a subsequence we can suppose that $\xi_n \rightarrow \xi$. Since $\partial\gamma$ and $\text{SS}(\Phi_{U,\gamma}(\mathcal{F}))$ are closed we get by the estimate of the micro-support that there exists $x \in \overline{U}$ such that $(x, \xi) \in \text{SS}(\mathcal{F})$ which is impossible by hypothesis. $\square$

Finally, let us state an existence lemma for refined cutting pairs.

Lemma 3.3.7. *Let X be a real vector space and $L \subset X$ a subspace of X . Then there exists a fundamental system of open conic neighborhoods γ of $(\dot{T}_L^*X)_0$ such that for each γ there exists a fundamental system of open neighborhoods U of 0 in X such that (U, γ) is a refined cutting pair.*

Proof. This lemma is shown during the proof of Corollary 3.4 using Lemma 3.3 in [D'A]. $\square$

We are now ready to give the refined version of the microlocal cut-off lemma along a linear subspace.

Proposition 3.3.8 (Refined microlocal cut-off).

*Let X be a real vector space and $L \subset X$ a subspace. Let $\mathcal{F} \in \text{Ob } D^b(k_X)$. Then there exists a fundamental system of open conic neighborhoods γ of $(T_L^*X)_0$ in $T_0^*X = X^*$ and for each γ there exists a fundamental system of open neighborhoods U of 0, such that*

- (i) *the natural morphism $u : \Phi_{U,\gamma}(\mathcal{F}) \rightarrow \mathcal{F}$ induces an isomorphism in $D^b(k_X, U \times \gamma)$.*
- (ii) $\text{SS}(\Phi_{U,\gamma}(\mathcal{F})) \subset X \times \overline{\gamma}$
- (iii) $\dot{\text{SS}}(\Phi_{U,\gamma}(\mathcal{F})) \cap \partial\gamma \cap \pi^{-1}(0) \subset \left\{ \xi \in \partial\gamma \mid \exists x \in \overline{U} : (x, \xi) \in \text{SS}(\mathcal{F}) \right\}$ where we have set $\dot{\text{SS}}(\Phi_{U,\gamma}(\mathcal{F})) = \text{SS}(\Phi_{U,\gamma}(\mathcal{F})) \cap \dot{T}^*X$.

3.4 Morphisms in $D^b(k_X, \{x\} \times \dot{\delta})$

In this section we will calculate the morphisms in the category $D^b(k_X, \{x\} \times \dot{\delta})$ where $x \in X$ and $\delta \subset T_x^*X$ is a closed cone. More precisely we will show that the natural morphism

$$\text{Hom}_{D^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \longrightarrow H^0(\{x\} \times \dot{\delta}, \mu\text{hom}(\mathcal{F}, \mathcal{G}))$$

is an isomorphism. In order to prove this, we will consider the composition

$$\varinjlim_{U, \gamma} H^0 \text{RHom}(\Phi_{U,\gamma}(\mathcal{F})_U, \mathcal{G}) \longrightarrow \text{Hom}_{D^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \longrightarrow H^0(\{x\} \times \dot{\delta}, \mu\text{hom}(\mathcal{F}, \mathcal{G})) \quad (*)_{\delta}$$

and show that it is an isomorphism. Here U runs through the family of relatively compact open neighborhoods of 0 and γ through the set of open cones containing $\dot{\delta}$.

Proposition 3.4.1. *Let X be a real vector space and consider a closed convex proper cone $\delta \subset X$. Then the natural morphism*

$$\varinjlim_{U, \gamma} H^n \text{RHom}(\Phi_{U,\gamma}(\mathcal{F})_U, \mathcal{G}) \longrightarrow H^n(\{0\} \times \dot{\delta}, \mu\text{hom}(\mathcal{F}, \mathcal{G})) \quad (*)_{\delta}$$

is an isomorphism. Here U runs through the family of relatively compact open neighborhoods of 0 and γ through the set of open cones containing $\dot{\delta}$.

Proof. First note that since $\mu\text{hom}(\mathcal{F}, \mathcal{G})$ is conic, we have

$$H^n(\{0\} \times \dot{\delta}, \mu\text{hom}(\mathcal{F}, \mathcal{G})) \simeq \varinjlim_{U, \gamma} H^n(U \times \gamma, \mu\text{hom}(\mathcal{F}, \mathcal{G})).$$

Since γ is open, convex and proper, we have by [KS1], Theorem 4.3.2

$$H^n(U \times \gamma, \mu\text{hom}(\mathcal{F}, \mathcal{G})) \simeq \varinjlim_{V, Z} H^n_{Z \cap V}(V, R\mathcal{H}om(q_2^{-1}\mathcal{F}, q_1^!\mathcal{G}))$$

where V runs through the family of open subsets of $T_{\Delta_X}(k_X \times X) \simeq X \times X$ such that $V \cap \Delta_X = U$ (i.e. $q_1(V) = U$) and Z through the family of closed subsets such that the inclusion $C_{\Delta_X} Z \subset U \times \gamma^\circ$ holds ⁶.

We have the following chain of isomorphisms

$$\begin{aligned} R\Gamma_{Z \cap V}(V, R\mathcal{H}om(q_2^{-1}\mathcal{F}, q_1^!\mathcal{G})) &\simeq R\Gamma(V, R\Gamma_{Z \cap V} R\mathcal{H}om(q_2^{-1}\mathcal{F}, q_1^!\mathcal{G})) \\ &\simeq R\Gamma(V, R\mathcal{H}om((q_2^{-1}\mathcal{F})_{Z \cap V}, q_1^!\mathcal{G})) \simeq R\Gamma(U \times X, R\mathcal{H}om((q_2^{-1}\mathcal{F})_{Z \cap V}, q_1^!\mathcal{G})) \\ &\simeq R\Gamma(U, R\mathcal{H}om(Rq_{1!}(q_2^{-1}\mathcal{F})_{Z \cap V}, \mathcal{G})) \simeq R\text{Hom}((Rq_{1!}(k_{Z \cap V} \otimes q_2^{-1}\mathcal{F})_U, \mathcal{G})). \end{aligned}$$

Hence we have

$$H^n(\{0\} \times \dot{\delta}, \mu\text{hom}(\mathcal{F}, \mathcal{G})) \simeq \varinjlim_{U, \gamma} \varinjlim_{V, Z} H^n R\text{Hom}((Rq_{1!}(k_{Z \cap V} \otimes q_2^{-1}\mathcal{F})_U, \mathcal{G})).$$

Now fix U, γ and V, Z . Then V contains a small relatively compact open neighborhood of 0 of type $U' \times U'$. Moreover we may assume by cofinality that Z is of the form $\tilde{s}^{-1}\gamma^\circ$ in a neighborhood of 0. Hence $Z \cap V$ contains $\tilde{s}^{-1}\gamma^\circ \cap U' \times U'$. We can therefore remove the second limit by replacing $V \cap Z$ with $\tilde{s}^{-1}\gamma^\circ \cap U \times U$. Then we get

$$\begin{aligned} R\text{Hom}((Rq_{1!}(k_{Z \cap V} \otimes q_2^{-1}\mathcal{F})_U, \mathcal{G})) &\simeq R\text{Hom}((Rq_{1!}(\tilde{s}^{-1}k_{\gamma^\circ} \otimes q_2^{-1}\mathcal{F}_U))_U, \mathcal{G}) \\ &\simeq R\text{Hom}((R s_!(q_1^{-1}k_{\gamma^\circ}^\wedge \otimes q_2^{-1}\mathcal{F}_U))_U, \mathcal{G}) \simeq R\text{Hom}(\Phi_{U, \gamma}(\mathcal{F})_U, \mathcal{G}). \end{aligned}$$

Therefore

$$H^n(\{0\} \times \dot{\delta}, \mu\text{hom}(\mathcal{F}, \mathcal{G})) \simeq \varinjlim_{U, \gamma} H^n R\text{Hom}(\Phi_{U, \gamma}(\mathcal{F})_U, \mathcal{G}).$$

□

Lemma 3.4.2. *For any closed cone $\delta \subset X$ consider the morphism $(*)_\delta$. Let $\delta_1, \delta_2 \subset X$ be two closed cones such that the morphisms $(*)_{\delta_1}, (*)_{\delta_2}$ and $(*)_{\delta_1 \cap \delta_2}$ are isomorphisms.*

Then $()_{\delta_1 \cup \delta_2}$ is an isomorphism.*

Proof. By Lemma 3.3.3 we get a morphism of distinguished triangles such that the vertical morphisms are given by $(*)_{\delta_1 \cap \delta_2}, (*)_{\delta_1} \oplus (*)_{\delta_2}$ and $(*)_{\delta_1 \cup \delta_2}$. Then the lemma follows from the Five Lemma. □

Proposition 3.4.3. *Let X be a real vector space and consider a closed cone $\delta \subset X$. Then the natural morphism*

$$\varinjlim_{U, \gamma} H^n R\text{Hom}(\Phi_{U, \gamma}(\mathcal{F})_U, \mathcal{G}) \longrightarrow H^n(\{0\} \times \dot{\delta}, \mu\text{hom}(\mathcal{F}, \mathcal{G})) \quad (*)_\delta$$

is an isomorphism. Here U runs through the family of relatively compact open subsets of 0 and γ through the set of open cones containing $\dot{\delta}$.

⁶Here $C_{\Delta_X} Z$ denotes the normal cone to Z along the diagonal Δ_X (see [KS1], Definition 4.1.1) and γ° denotes the polar cone of γ , i.e.

$$\gamma^\circ = \{x \in X^* \mid \langle y, x \rangle \geq 0 \text{ for all } y \in \gamma.\}$$

Proof. First suppose that δ can be written as a finite union of closed convex proper cones. Note that the intersection of two proper, closed, convex cones is again proper, closed and convex. Therefore, if δ' is the union of n closed, convex, proper cones then the intersection of a closed, convex, proper cone with δ' can be written as a union of n closed convex proper cones. Using Proposition 3.4.1 and the previous lemma we can then easily show the proposition by induction on the number of closed, convex, proper cones that cover δ . Now let us consider the general case. Every closed cone is a decreasing intersection of closed cones δ_i that can be covered by a finite number of closed convex proper cones⁷. Then $(*)_\delta = \varinjlim (*)_{\delta_i}$ is an isomorphism. $\square$

Theorem 3.4.4. *Let X be a real manifold, $x \in X$ and $\delta \subset T_x^*X$ a closed cone. Then the natural morphism*

$$\mathrm{Hom}_{\mathrm{D}^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \longrightarrow \mathrm{H}^0(\{x\} \times \dot{\delta}, \mu\mathrm{hom}(\mathcal{F}, \mathcal{G}))$$

is an isomorphism.

Proof. By the last proposition we know that the morphism $(*)_\delta$ is an isomorphism. Therefore the morphism of the theorem is surjective.

Let us prove that it is injective.

Let $\mathcal{F} \rightarrow \mathcal{G}$ be a morphism of $\mathrm{D}^b(k_X, \{x\} \times \dot{\delta})$ that is zero in $\mathrm{H}^0(\{x\} \times \dot{\delta}, \mu\mathrm{hom}(\mathcal{F}, \mathcal{G}))$. Then we may represent this morphism by a morphism $\mathcal{F}' \rightarrow \mathcal{G}$ in $\mathrm{D}^b(k_X)$ and a morphism $\mathcal{F}' \rightarrow \mathcal{F}$ that is an isomorphism on $\{x\} \times \dot{\delta}$. We get a commutative diagram

$$\begin{array}{ccccc} \mathrm{Hom}_{\mathrm{D}^b(k_X)}(\mathcal{F}', \mathcal{G}) & \longrightarrow & \mathrm{Hom}_{\mathrm{D}^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) & \longrightarrow & \varinjlim_{U, \gamma} \mathrm{R}^0 \mathrm{Hom}(\Phi_{U, \gamma}(\mathcal{F})_U, \mathcal{G}) \\ \downarrow \mathrm{Id} & & \downarrow \sim & & \downarrow \sim \\ \mathrm{Hom}_{\mathrm{D}^b(k_X)}(\mathcal{F}', \mathcal{G}) & \longrightarrow & \mathrm{Hom}_{\mathrm{D}^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}', \mathcal{G}) & \longrightarrow & \varinjlim_{U, \gamma} \mathrm{R}^0 \mathrm{Hom}(\Phi_{U, \gamma}(\mathcal{F}')_U, \mathcal{G}). \end{array}$$

Using the diagram, we see that there exists (U, γ) such that $\Phi_{U, \gamma}(\mathcal{F}')_U \rightarrow \mathcal{F}' \rightarrow \mathcal{G}$ is the zero map in $\mathrm{D}^b(k_X)$. But $\Phi_{U, \gamma}(\mathcal{F}')_U \rightarrow \mathcal{F}'$ is an isomorphism on $\{x\} \times \dot{\delta}$, and therefore $\mathcal{F}' \rightarrow \mathcal{G}$ represents the zero morphism in $\mathrm{D}^b(k_X, \{x\} \times \dot{\delta})$. $\square$

4 Microlocalization of constructible sheaves

4.1 Microlocalization of $\mathbb{R}$ -constructible sheaves

Consider the full triangulated subcategory $\mathrm{D}_{\mathbb{R}\text{-c}}^b(k_X) \subset \mathrm{D}^b(k_X)$ and a subset $S \subset T^*X$. There are two obvious ways to define the microlocalization of the derived category of $\mathbb{R}$ -constructible sheaves on S that we recall now. Set

$$\mathcal{N}_{\mathbb{R}\text{-c}, S} = \mathcal{N}_S \cap \mathrm{D}_{\mathbb{R}\text{-c}}^b(k_X).$$

Then the inclusion $\mathrm{D}_{\mathbb{R}\text{-c}}^b(k_X) \subset \mathrm{D}^b(k_X)$ induces a functor

$$\mathrm{D}_{\mathbb{R}\text{-c}}^b(k_X) / \mathcal{N}_{\mathbb{R}\text{-c}, S} \longrightarrow \mathrm{D}^b(k_X) / \mathcal{N}_S = \mathrm{D}^b(k_X, S).$$

⁷The proof is done by a simple compactness argument in $\dot{X}/\mathbb{R}^+$. For every $p \in \delta$ choose a closed convex proper cone with angle ε that contains p . Then a finite number of these cones, say $\gamma_1, \gamma_2, \dots, \gamma_n$, will cover δ and we set $\delta_1 = \bigcup_{i=1, \dots, n} \gamma_i$. The next cone δ_2 is constructed by choosing for each point a closed

convex proper cone with angle $\varepsilon/2$. Again a finite number will cover δ , say $\gamma'_1, \dots, \gamma'_m$. Then we define δ_2 as the union of all intersections $\gamma_i \cap \gamma'_j$ and we proceed by induction. It is clear by construction that the intersection of the δ_i is the cone δ .

Clearly the objects of the image of this functors are complexes in $D^b(k_X, S)$ with $\mathbb{R}$ -constructible cohomology. But the functor is not fully faithful in general. For our purpose it will be convenient to work with the category $D_{\mathbb{R}\text{-c}}^b(k_X)/\mathcal{N}_{\mathbb{R}\text{-c}, S}$ as does Andronikof in [An1],[An2].

Definition 4.1.1. *We set*

$$D_{\mathbb{R}\text{-c}}^b(k_X, S) = D_{\mathbb{R}\text{-c}}^b(k_X)/\mathcal{N}_{\mathbb{R}\text{-c}, S}.$$

Remark 4.1.2. Let us emphasize again that the natural functor

$$D_{\mathbb{R}\text{-c}}^b(k_X, S) \longrightarrow D^b(k_X, S) \quad (4.1.1)$$

is not fully faithful. Hence although the objects of $D_{\mathbb{R}\text{-c}}^b(k_X, S)$ have $\mathbb{R}$ -constructible cohomology, the category can not be identified to the full subcategory of $D^b(k_X, S)$ whose objects have $\mathbb{R}$ -constructible cohomology. More precisely, let $\mathcal{F}, \mathcal{G} \in D_{\mathbb{R}\text{-c}}^b(k_X)$.

- (i) A morphism $\mathcal{F} \rightarrow \mathcal{G}$ in $D_{\mathbb{R}\text{-c}}^b(k_X, S)$ is represented by a morphism $\mathcal{F}' \rightarrow \mathcal{G}$ where $\mathcal{F}'$ has $\mathbb{R}$ -constructible cohomology sheaves and is isomorphic to $\mathcal{F}$ on S .
- (ii) A morphism $\mathcal{F} \rightarrow \mathcal{G}$ in $D^b(k_X, S)$ is represented by a morphism $\mathcal{F}' \rightarrow \mathcal{G}$ where $\mathcal{F}' \in D^b(k_X)$ is isomorphic to $\mathcal{F}$ on S .

Therefore it is not obvious to compare morphisms in these two categories.

Nevertheless note that for any $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(k_X, S)$ we have by definition $\mathcal{F} \simeq 0$ if and only if $\text{SS}(\mathcal{F}) \cap S = \emptyset$. Hence if $\mathcal{F} \rightarrow \mathcal{G}$ is a morphism in $D_{\mathbb{R}\text{-c}}^b(k_X)$ we get that $\mathcal{F} \rightarrow \mathcal{G}$ is an isomorphism in $D_{\mathbb{R}\text{-c}}^b(k_X, S)$ if and only if it is an isomorphism in $D^b(k_X, S)$, hence if and only if there is a distinguished triangle in $D_{\mathbb{R}\text{-c}}^b(k_X)$

$$\mathcal{F} \longrightarrow \mathcal{G} \longrightarrow \mathcal{H} \xrightarrow{+}$$

such that $\text{SS}(\mathcal{H}) \cap S = \emptyset$. More generally we get

Proposition 4.1.3. *The natural functor*

$$D_{\mathbb{R}\text{-c}}^b(k_X, S) \longrightarrow D^b(k_X, S)$$

is conservative, i.e. a morphism $\mathcal{F} \rightarrow \mathcal{G}$ of $D_{\mathbb{R}\text{-c}}^b(k_X, S)$ is an isomorphism in $D_{\mathbb{R}\text{-c}}^b(k_X, S)$ if and only if it is an isomorphism in $D^b(k_X, S)$.

Proof. We embed $\mathcal{F} \rightarrow \mathcal{G}$ in a distinguished triangle

$$\mathcal{F} \longrightarrow \mathcal{G} \longrightarrow \mathcal{H} \xrightarrow{+}$$

in $D_{\mathbb{R}\text{-c}}^b(k_X, S)$. If $\mathcal{F} \rightarrow \mathcal{G}$ is an isomorphism in $D^b(k_X, S)$ then $\mathcal{H} \simeq 0$ in $D^b(k_X, S)$. Hence $\text{SS}(\mathcal{H}) \cap S = \emptyset$. Therefore $\mathcal{H} \simeq 0$ in $D_{\mathbb{R}\text{-c}}^b(k_X, S)$ and $\mathcal{F} \rightarrow \mathcal{G}$ is an isomorphism in $D_{\mathbb{R}\text{-c}}^b(k_X, S)$. $\square$

However there are some situations when the functor (4.1.1) is fully faithful. For instance, Andronikof remarked that the proof of Proposition 3.2.3 holds in the constructible case, hence

Proposition 4.1.4. *Let $\mathcal{F}, \mathcal{G} \in D_{\mathbb{R}\text{-c}}^b(k_X, p)$. Then there is a canonical isomorphism*

$$\text{Hom}_{D_{\mathbb{R}\text{-c}}^b(k_X, p)}(\mathcal{F}, \mathcal{G}) \xrightarrow{\sim} H^0 \mu \text{hom}(\mathcal{F}, \mathcal{G})_p$$

and the natural functor

$$D_{\mathbb{R}\text{-c}}^b(k_X, p) \longrightarrow D^b(k_X, p)$$

is fully faithful.

4.2 The category $D_{\mathbb{R}\text{-c}}^b(k_X, \{x\} \times \dot{\delta})$

We will see that the natural morphism (4.1.1) is an isomorphism for orbits of $\mathbb{C}^\times$ and more generally for closed cones $\dot{\delta}$ in $\dot{T}_x^*X$ for some $x \in X$.

Proposition 4.2.1. *Let $\mathcal{F}, \mathcal{G} \in D_{\mathbb{R}\text{-c}}^b(k_X)$. Then the natural morphism*

$$\mathrm{Hom}_{D_{\mathbb{R}\text{-c}}^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \longrightarrow \mathrm{Hom}_{D^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G})$$

is an isomorphism.

Proof. We may assume that X is a vector space. Note that if $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(k_X)$, then $\Phi_{U, \gamma}(\mathcal{F})_U$ is $\mathbb{R}$ -constructible for any relatively compact subanalytic open subset $U \ni 0$ and any subanalytic open cone $\gamma \subset X$.

Then the results of Section 3.4 hold in the $\mathbb{R}$ -constructible case. More precisely, we see first (as in Proposition 3.4.3) that the composition

$$\begin{aligned} \varinjlim_{U, \gamma} H^0 \mathrm{RHom}(\Phi_{U, \gamma}(\mathcal{F})_U, \mathcal{G}) &\longrightarrow \mathrm{Hom}_{D_{\mathbb{R}\text{-c}}^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \longrightarrow \mathrm{Hom}_{D^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \\ &\longrightarrow H^0(\{x\} \times \dot{\delta}, \mu\mathrm{hom}(\mathcal{F}, \mathcal{G})) \end{aligned}$$

is an isomorphism and then (as in Theorem 3.4.4) that the composition

$$\mathrm{Hom}_{D_{\mathbb{R}\text{-c}}^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \longrightarrow \mathrm{Hom}_{D^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \longrightarrow H^0(\{x\} \times \dot{\delta}, \mu\mathrm{hom}(\mathcal{F}, \mathcal{G}))$$

is an isomorphism. Since the second morphism of the last composition is an isomorphism, we get the result. $\square$

Combining Proposition 4.2.1 and Theorem 3.4.4 we get the following theorem.

Theorem 4.2.2. *The natural functor*

$$D_{\mathbb{R}\text{-c}}^b(k_X, \{x\} \times \dot{\delta}) \longrightarrow D^b(k_X, \{x\} \times \dot{\delta})$$

is fully faithful. Moreover for every $\mathcal{F}, \mathcal{G} \in D_{\mathbb{R}\text{-c}}^b(k_X, \{x\} \times \dot{\delta})$ we have

$$\mathrm{Hom}_{D_{\mathbb{R}\text{-c}}^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \xrightarrow{\sim} H^0(\{x\} \times \dot{\delta}, \mu\mathrm{hom}(\mathcal{F}, \mathcal{G})).$$

4.3 Microlocally $\mathbb{C}$ -constructible sheaves

Recall the microlocal characterization of complexes with $\mathbb{C}$ -constructible cohomology sheaves given in [KS1] (Theorem 8.5.5).

Proposition 4.3.1. *A complex $\mathcal{F}$ in $D^b(k_X)$ has $\mathbb{C}$ -constructible cohomology if and only if $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(k_X)$ and $\mathrm{SS}(\mathcal{F})$ is a $\mathbb{C}^\times$ -conic subset of T^*X .*

If $S \subset T^*X$ is a not necessarily $\mathbb{C}^\times$ -conic subset, then this suggests the definition of a microlocally $\mathbb{C}$ -constructible sheaf on S in as follows:

Definition 4.3.2. (i) *An object $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(k_X)$ (or $D_{\mathbb{R}\text{-c}}^b(k_X, S')$ for $S' \supset S$) is called microlocally $\mathbb{C}$ -constructible on S if $\mathrm{SS}(\mathcal{F})$ is $\mathbb{C}^\times$ -conic on S .*

(ii) *We denote by $D_{\mathbb{C}\text{-c}}^b(k_X, S)$ be the full subcategory of $D_{\mathbb{R}\text{-c}}^b(k_X, S)$ consisting of microlocally $\mathbb{C}$ -constructible sheaves (on S).*

Remark 4.3.3. Note that the category of microlocally $\mathbb{C}$ -constructible sheaves (on S) is different from the category $D_{\mathbb{C}\text{-c}}^b(k_X)/(\mathcal{N}_S \cap D_{\mathbb{C}\text{-c}}^b(k_X))$ (i.e. the microlocalization of $\mathbb{C}$ -constructible sheaves). There is a natural functor

$$D_{\mathbb{C}\text{-c}}^b(k_X)/(\mathcal{N}_S \cap D_{\mathbb{C}\text{-c}}^b(k_X)) \longrightarrow D_{\mathbb{C}\text{-c}}^b(k_X, S),$$

but in general, an object in $D_{\mathbb{C}\text{-c}}^b(k_X, S)$ cannot be represented by a complex with $\mathbb{C}$ -constructible cohomology sheaves. One shall keep in mind that by definition an object in $D_{\mathbb{R}\text{-c}}^b(k_X, S)$ (the microlocalization of $\mathbb{R}$ -constructible sheaves) is represented by an $\mathbb{R}$ -constructible sheaf on X . Since $\mathbb{R}$ -constructible sheaves can be defined by a microlocal property⁸, there also exists a natural definition of a microlocally $\mathbb{R}$ -constructible sheaf which we do not consider in this paper.

Remark 4.3.4. Of course Definition 4.3.2, (i) is equivalent to the statement

- (i) A sheaf $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(k_X)$ (resp. $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(k_X, S')$ for $S' \supset S$) is microlocally $\mathbb{C}$ -constructible on S if for every point p of S there exists an open neighborhood U of p such that $U \cap \text{SS}(\mathcal{F}) = U \cap \mathbb{C}^\times \text{SS}(\mathcal{F})$.

Obviously $D_{\mathbb{C}\text{-c}}^b(k_X, T^*X) = D_{\mathbb{C}\text{-c}}^b(k_X)$ and if $x \in X$, then $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(k_X)$ defines an object of $D_{\mathbb{C}\text{-c}}^b(k_X, x)$ if and only if $\mathcal{F}|_V$ is $\mathbb{C}$ -constructible for some neighborhood V of x .

However the category $D_{\mathbb{C}\text{-c}}^b(k_X, S)$ is not very easy to understand in general, especially if S is not $\mathbb{C}^\times$ -conic.

Lemma 4.3.5. *Let $\mathcal{F} \in D^b(k_X)$ and $S \subset \dot{T}^*X$*

- (i) *The object $\mathcal{F}$ is microlocally $\mathbb{C}$ -constructible on S if and only if $\mathcal{F}$ is microlocally $\mathbb{C}$ -constructible on $\mathbb{R}^+S$.*
- (ii) *Suppose that $S \subset \dot{T}^*X$ is $\mathbb{C}^\times$ -conic. Then $\mathcal{F}$ is microlocally $\mathbb{C}$ -constructible on S if and only if $\mathcal{F}$ is microlocally $\mathbb{C}$ -constructible in $\gamma^{-1}(U)$ where U is a germ of a neighborhood of $\gamma(S)$ in P^*X .*

Proof. Statement (i) is a consequence of Lemma 3.2.2 and (ii) follows from (i) and the fact that any $\mathbb{R}^+$ -conic neighborhood of S contains a $\mathbb{C}^\times$ -conic neighborhood. $\square$

Remark 4.3.6. There is an obvious functor $D_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times S) \rightarrow D_{\mathbb{C}\text{-c}}^b(k_X, S)$. One might ask the question whether or not a sheaf $\mathcal{F}$ of $D_{\mathbb{C}\text{-c}}^b(k_X, S)$ can be lifted to $D_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times S)$ and if there is a tool to produce an object of $D_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times S)$ that is isomorphic to $\mathcal{F}$ on S . There does not seem to be an obvious answer as the following example shows:

Consider $X = \mathbb{C}^2$ and the sheaf $\mathcal{F} = \mathbb{C}_{\mathbb{C}^\times \times \{(0,0)\}} \oplus \mathbb{C}_{\{0\} \times \mathbb{R} \times \mathbb{R} \times \{0\}}$. Then

$$\text{SS}(\mathcal{F}) = T_{\mathbb{C}^\times \times \{(0,0)\}}^* X \cup T_{\{0\} \times \mathbb{R} \times \mathbb{R} \times \{0\}}^* X.$$

Take $p = ((0, 0, 0, 0); (0, 0, 1, 1)) \in \dot{T}^*X$. Then $\text{SS}(\mathcal{F}) = \mathbb{C}^\times p$ in a neighborhood of p . But if $U \supset \mathbb{C}^\times p$ is an arbitrary neighborhood of $\mathbb{C}^\times p$, $\text{SS}(\mathcal{F})$ is not $\mathbb{C}^\times$ -conic on U , hence $\mathcal{F}$ is microlocally $\mathbb{C}$ -constructible at p but not on $\mathbb{C}^\times p$. However $\mathcal{F}$ is isomorphic in $D_{\mathbb{R}\text{-c}}^b(k_X, p)$ to the sheaf $\mathbb{C}_{\mathbb{C}^\times \times \{(0,0)\}}$ which is globally $\mathbb{C}$ -constructible. The problem is how to construct $\mathbb{C}_{\mathbb{C}^\times \times \{(0,0)\}}$ functorially from $\mathcal{F}$. It cannot be done by a cut-off functor which will always preserve the micro-support in a neighborhood of $\mathbb{C}^\times p$.

Hence microlocally $\mathbb{C}$ -constructible sheaves should be defined on P^*X rather than T^*X and Definition 4.3.2 will mostly be used for a $\mathbb{C}^\times$ -conic subset X .

⁸An object $\mathcal{F} \in D^b(k_X)$ is weakly $\mathbb{R}$ -constructible if and only if its micro-support is Lagrangian.

4.4 The category $D_{\mathbb{C}\text{-}c}^b(k_X, \{x\} \times \dot{\delta})$

Proposition 4.4.1. *The natural functors*

$$D_{\mathbb{C}\text{-}c}^b(k_X, \{x\} \times \dot{\delta}) \longrightarrow D_{\mathbb{R}\text{-}c}^b(k_X, \{x\} \times \dot{\delta}) \longrightarrow D^b(k_X, \{x\} \times \dot{\delta})$$

are fully faithful. Moreover for every $\mathcal{F}, \mathcal{G} \in D_{\mathbb{R}\text{-}c}^b(k_X, \{x\} \times \dot{\delta})$ we have

$$\mathrm{Hom}_{D_{\mathbb{C}\text{-}c}^b(k_X, \{x\} \times \dot{\delta})}(\mathcal{F}, \mathcal{G}) \xrightarrow{\sim} H^0(\mathbb{C}^{\times p}, \mu\mathrm{hom}(\mathcal{F}, \mathcal{G})).$$

Proof. The first functor is fully faithful by definition, the second by Theorem 4.2.2. The second part follows again from Theorem 4.2.2. $\square$

5 Invariance by quantized contact transformations

Let $\Omega_X \subset T^*X$ be an open subset of a real manifold X . In [KS1], Kashiwara-Schapira showed that the category $D^b(k_X, \Omega_X)$ (or more generally the prestack $D^b(k_X, *)|_{\Omega_X}$) is invariant under “quantized contact transformations”. Let us briefly explain this statement. Consider real manifolds X, Y of the same dimension and open subsets $\Omega_X \subset T^*X$, $\Omega_Y \subset T^*Y$. An $\mathbb{R}^+$ -homogenous symplectic isomorphism

$$\chi : \Omega_X \xrightarrow{\sim} \Omega_Y$$

is often called a contact transformation (although strictly speaking, the contact structures are defined on the projective bundles). Invariance under “quantized contact transformations” means that locally we can construct from χ an equivalence of categories

$$\Phi_{\mathcal{K}} : D^b(k_X, \Omega_X) \xrightarrow{\sim} D^b(k_Y, \Omega_Y).$$

The equivalence $\Phi_{\mathcal{K}}$ is explicitly given by an integral transform and depends on the choice of a kernel $\mathcal{K} \in D^b(k_{Y \times X})$. The main result ([KS1], Corollary 7.2.2) is:

Theorem 5.0.2. *Let X, Y be two real manifolds, $\Omega_X \subset T^*X$, $\Omega_Y \subset T^*Y$ open subsets and*

$$\chi : \Omega_X \longrightarrow \Omega_Y$$

a real contact transformation. Set

$$\Lambda = \left\{ ((y; \eta), (x; \xi)) \in \Omega_Y \times \Omega_X^a \mid (y, \eta) = \chi(x, -\xi) \right\}.$$

Let $p_X \in \Omega_X$ and $p_Y = \chi(p_X)$.

*There exist open neighborhoods X' of $\pi(p_X)$, Y' of $\pi(p_Y)$, Ω'_X of p_X , Ω'_Y of p_Y with $\Omega'_X \subset T^*X' \cap \Omega_X$, $\Omega'_Y \subset T^*Y' \cap \Omega_Y$ and a kernel $\mathcal{K} \in D^b(k_{Y' \times X'})$ such that:*

(1) χ induces a contact transformation $\Omega'_X \xrightarrow{\sim} \Omega'_Y$,

(2)

$$\left((\Omega_Y'' \times T^*X') \cup (T^*Y' \times \Omega_X''^a) \right) \cap \mathrm{SS}(\mathcal{K}) \subset \Lambda \cap (\Omega_Y'' \times \Omega_X''^a)$$

for every open subsets $\Omega_X'' \subset \Omega'_X$ and $\Omega_Y'' = \chi(\Omega_X'')$,

(3) *composition with $\mathcal{K}$ induces an equivalence of prestacks*

$$\Phi_{\mathcal{K}} = \mathcal{K} \circ : \chi_* D^b(k'_X, *)|_{\Omega'_X} \longrightarrow D^b(k_{Y'}, *)|_{\Omega'_Y},$$

a quasi-inverse being given by $\Phi_{\mathcal{K}^}$ with $\mathcal{K}^* = r_* R\mathcal{H}om(\mathcal{K}, \omega_{Y \times X|X})$ where $r : Y \times X \rightarrow X \times Y$ switches the factors.*

$$(4) \text{ SS}(\Phi_{\mathcal{K}}(\mathcal{F})) \cap \Omega_Y'' = \chi(\text{SS}(\mathcal{F}) \cap \Omega_X''),$$

$$(5) \chi_* \mu\text{hom}(\mathcal{F}, \mathcal{G})|_{\Omega_X'} \simeq \mu\text{hom}(\Phi_{\mathcal{K}}(\mathcal{F}), \Phi_{\mathcal{K}}(\mathcal{G}))|_{\Omega_Y'}.$$

Remark 5.0.3. There is a slight difference to [KS1], Corollary 7.2.2 where the smaller open subset Ω_X'' is not introduced. For instance, statement (3) is stated as an equivalence of categories

$$\Phi_{\mathcal{K}} : D^b(k_{X'}, \Omega_X') \longrightarrow D^b(k_{Y'}, \Omega_Y').$$

However, it is obvious that if we have the theorem for fixed $X', Y', \Omega_X', \Omega_Y'$, then (1), (2), (4) and (5) are valid for any open subset $\Omega_X'' \subset \Omega_X'$ and $\Omega_Y'' = \chi(\Omega_X'')$. The fact that (3) is valid on Ω_X'' is based on the observation that (3) is essentially a consequence of (2) (plus some independent conditions on $\mathcal{K}$, cf. [KS1], Theorem 7.2.1).

Now let us consider constructible sheaves. It is not immediately obvious that the equivalence (3) of Theorem 5.2.1 should induce an equivalence on the microlocalization of $\mathbb{R}$ -constructible (resp. on microlocally $\mathbb{C}$ -constructible or later on microlocally perverse) sheaves

$$\chi_* D_{\mathbb{R}\text{-c}}^b(k_X, *)|_{\Omega_X} \xrightarrow{?} D_{\mathbb{R}\text{-c}}^b(k_Y, *)|_{\Omega_Y}$$

since the functor $\Phi_{\mathcal{K}}$ is not well defined on $\mathbb{R}$ -constructible sheaves. This problem can be solved at a point $p \in T^*X$ by using the microlocal composition of [KS1]. In [An2], Andronikof uses this tool to construct the functor $\Phi_{\mathcal{K}}^\mu$. Then one can treat a variety of kernels $\mathcal{K}$ but a priori, one can no longer work in an open neighborhood.

However, under the hypothesis of Theorem 5.0.2, we do not need to use microlocal composition of kernels. We can always define the functor

$$\chi_* D_{\mathbb{R}\text{-c}}^b(k_X, *)|_{\Omega_X} \longrightarrow \chi_* D^b(k_X, *)|_{\Omega_X} \xrightarrow{\Phi_{\mathcal{K}}} D^b(k_Y, *)|_{\Omega_Y}$$

and hope that it factors through $D_{\mathbb{R}\text{-c}}^b(k_Y, *)|_{\Omega_Y}$. However, one has to beware that $D_{\mathbb{R}\text{-c}}^b(k_Y, *)|_{\Omega_Y}$ is not a full subprestack of $D^b(k_Y, *)|_{\Omega_Y}$, hence it is not sufficient to show that $\Phi_{\mathcal{K}}(\mathcal{F})$ is isomorphic to an $\mathbb{R}$ -constructible sheaf in $D^b(k_Y, *)|_{\Omega_Y}$. We encounter this problem in Section 5.1 (see Theorem 5.1.2).

5.1 Quantization of $\mathbb{R}$ -constructible sheaves

Recall the definition of the full subcategory $N(Y, X, \Omega_Y, \Omega_X)$ of $D^b(k_{Y \times X}, \Omega_Y \times T^*X)$. Its objects are kernels $\mathcal{K}$ on $Y \times X$ such that

- (i) $\text{SS}(\mathcal{K}) \cap (\Omega_Y \times T^*X) \subset \Omega_Y \times \Omega_X^a$
- (ii) The projection $p_1 : \text{SS}(\mathcal{K}) \cap (\Omega_Y \times T^*X) \rightarrow \Omega_Y$ is proper.

If $V = \pi_X(\Omega_X)$ is a subanalytic relatively compact open subset of X we set

$$N_{\mathbb{R}\text{-c}}(Y, X, \Omega_Y, \Omega_X) = N(Y, X, \Omega_Y, \Omega_X) \cap D_{\mathbb{R}\text{-c}}^b(k_{Y \times X}, \Omega_Y \times T^*X).$$

Definition 5.1.1. Let $\mathcal{K} \in N_{\mathbb{R}\text{-c}}(Y, X, \Omega_Y, \Omega_X)$. We define the functor $\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}}$ as

$$D_{\mathbb{R}\text{-c}}^b(k_X, \Omega_X) \longrightarrow D_{\mathbb{R}\text{-c}}^b(k_Y, \Omega_Y) \quad ; \quad \mathcal{F} \mapsto \mathcal{K} \circ \mathcal{F}_{\overline{V}}.$$

Note that $\mathcal{K} \circ \mathcal{F}_{\overline{V}}$ is $\mathbb{R}$ -constructible since V is subanalytic and relatively compact. We may visualize the situation by the following diagram

$$\begin{array}{ccc} D_{\mathbb{R}\text{-c}}^b(k_X, \Omega_X) & \xrightarrow{\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}}} & D_{\mathbb{R}\text{-c}}^b(k_Y, \Omega_Y) \\ \downarrow & & \downarrow \\ D^b(k_X, \Omega_X) & \xrightarrow{\mathcal{K} \circ} & D^b(k_Y, \Omega_Y). \end{array}$$

The square is commutative up to natural isomorphism since the natural morphism $\mathcal{F} \rightarrow \mathcal{F}_{\overline{V}}$ induces an isomorphism $\mathcal{K} \circ \mathcal{F} \xrightarrow{\sim} \mathcal{K} \circ \mathcal{F}_{\overline{V}}$ in $D^b(k_Y, \Omega_Y)$.

Now suppose that $\mathcal{K} \in N_{\mathbb{R}\text{-c}}(Z, Y, \Omega_Z, \Omega_Y)$ and $\mathcal{L} \in N_{\mathbb{R}\text{-c}}(Y, X, \Omega_Y, \Omega_X)$. Then their composition $\mathcal{K} \circ \mathcal{L}$ is well defined in $N(Z, X, \Omega_Z, \Omega_X)$ but not necessarily in $N_{\mathbb{R}\text{-c}}(Z, X, \Omega_Z, \Omega_X)$. Note however that by definition $\mathcal{L} \in D^b(k_{Y \times X}, \Omega_Y \times T^*X)$. Hence, if we set $W = \pi_Y(\Omega_Y)$, we do not distinguish between $\mathcal{L}$ and $\mathcal{L}_{\overline{W} \times X}$ in $\mathcal{L} \in N(Y, X, \Omega_Y, \Omega_X)$. In other words we get a natural isomorphism

$$\mathcal{K} \circ \mathcal{L} \xrightarrow{\sim} \mathcal{K} \circ \mathcal{L}_{\overline{W} \times X}$$

in $D^b(k_{Z \times X}, \Omega_Z \times T^*X)$. Then we get natural isomorphisms

$$\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}} \Phi_{\mathcal{L}}^{\mathbb{R}\text{-c}}(\mathcal{F}) \simeq \mathcal{K} \circ (\mathcal{L} \circ \mathcal{F}_{\overline{V}})_{\overline{W}} \simeq (\mathcal{K} \circ \mathcal{L}_{\overline{W} \times X}) \circ \mathcal{F}_{\overline{V}} \simeq \Phi_{\mathcal{K} \circ \mathcal{L}_{\overline{W} \times X}}^{\mathbb{R}\text{-c}}(\mathcal{F}).$$

Hence, the theory of microlocal kernels (Section 7.1. of [KS1]) works well in the $\mathbb{R}$ -constructible case if we restrict ourselves to relatively compact subanalytic open sets.

Finally suppose that $\Omega'_X \subset \Omega_X$, $\Omega'_Y \subset \Omega_Y$ and that $\mathcal{K} \in N_{\mathbb{R}\text{-c}}(Y, X, \Omega_Y, \Omega_X) \cap N_{\mathbb{R}\text{-c}}(Y, X, \Omega'_Y, \Omega'_X)$. Then we get a diagram

$$\begin{array}{ccc} D_{\mathbb{R}\text{-c}}^b(k_X, \Omega_X) & \xrightarrow{\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}}} & D_{\mathbb{R}\text{-c}}^b(k_Y, \Omega_Y) \\ \downarrow & & \downarrow \\ D_{\mathbb{R}\text{-c}}^b(k_X, \Omega'_X) & \xrightarrow{\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}}} & D_{\mathbb{R}\text{-c}}^b(k_Y, \Omega'_Y) \end{array}$$

that is commutative up to natural isomorphism induced by $(\mathcal{F}_{\overline{V}})_{\overline{V'}} \simeq \mathcal{F}_{\overline{V'}}$ where $V' = \pi(\Omega'_Y)$.

Now suppose that we are given a contact transformation

$$\chi : \Omega_X \xrightarrow{\sim} \Omega_Y$$

where we assume that $\pi(\Omega_X), \pi(\Omega_Y)$ are relatively compact subanalytic open sets. If there exists an object $\mathcal{K}$ such that $\mathcal{K} \in N_{\mathbb{R}\text{-c}}(Y, X, \Omega'_Y, \Omega'_X)$ for all open subsets $\Omega'_X \subset \Omega_X$ and $\Omega'_Y = \chi(\Omega'_X)$ then we get a commutative diagram of functors of prestacks

$$\begin{array}{ccc} D_{\mathbb{R}\text{-c}}^b(k_X, *)|_{\Omega_X} & \xrightarrow{\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}}} & D_{\mathbb{R}\text{-c}}^b(k_Y, *)|_{\Omega_Y} \\ \downarrow & & \downarrow \\ D^b(k_X, *)|_{\Omega_X} & \xrightarrow{\Phi_{\mathcal{K}}} & D^b(k_Y, *)|_{\Omega_Y} \end{array}$$

All compatibility conditions are easily verified by diagram chases.

We are now ready to quantize $\mathbb{R}$ -constructible sheaves. All we need to know is that the kernel produced in Theorem 5.0.2 can be taken $\mathbb{R}$ -constructible.

Theorem 5.1.2. *Let X, Y be two real manifolds, $\Omega_X \subset T^*X$, $\Omega_Y \subset T^*Y$ open subsets and*

$$\chi : \Omega_X \xrightarrow{\sim} \Omega_Y$$

a contact transformation. Set

$$\Lambda = \left\{ ((y; \eta), (x; \xi)) \in \Omega_Y \times \Omega_X^a \mid (y, \eta) = \chi(x, -\xi) \right\}.$$

Let $p_X \in \Omega_X$ and $p_Y = \chi(p_X)$.

*There exist open neighborhoods X' of $\pi(p_X)$, Y' of $\pi(p_Y)$, Ω'_X of p_X , Ω'_Y of p_Y with $\Omega'_X \subset T^*X' \cap \Omega_X$, $\Omega'_Y \subset T^*Y' \cap \Omega_Y$, and there exists $\mathcal{K} \in D_{\mathbb{R}\text{-c}}^b(k_{Y' \times X'})$ such*

(1) χ induces a contact transformation $\Omega'_X \xrightarrow{\sim} \Omega'_Y$,

(2)

$$\left((\Omega''_Y \times T^*X') \cup (T^*Y' \times \Omega''_X{}^a) \right) \cap \text{SS}(\mathcal{K}) \subset \Lambda \cap (\Omega''_Y \times \Omega''_X{}^a)$$

for every open subsets $\Omega''_X \subset \Omega'_X$ and $\Omega''_Y = \chi(\Omega''_X)$,

(3) composition with $\mathcal{K}$ induces an equivalence of prestacks

$$\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}} : \chi_* \text{D}_{\mathbb{R}\text{-c}}^b(k_{X'}, *)|_{\Omega'_X} \longrightarrow \text{D}_{\mathbb{R}\text{-c}}^b(k_{Y'}, *)|_{\Omega'_Y},$$

and $\Phi_{\mathcal{K}^*}^{\mathbb{R}\text{-c}}$ defines a quasi-inverse where $\mathcal{K}^* = r_* \text{R}\mathcal{H}om(\mathcal{K}, \omega_{Y \times X|X})$ and $r : Y \times X \rightarrow X \times Y$ switches the factors.

(4) $\text{SS}(\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}}(\mathcal{F})) \cap \Omega''_Y = \chi(\text{SS}(\mathcal{F}) \cap \Omega''_X)$,

(5) $\chi_* \mu\text{hom}(\mathcal{F}, \mathcal{G})|_{\Omega'_X} \simeq \mu\text{hom}(\Phi_{\mathcal{K}}(\mathcal{F}), \Phi_{\mathcal{K}}(\mathcal{G}))|_{\Omega'_Y}$.

Proof. Let us show that in the situation of Theorem 5.0.2 we can choose the kernel to be $\mathcal{K}$ $\mathbb{R}$ -constructible.

First assume that Λ is the conormal bundle of a smooth hypersurface $S \subset Y \times X$. Then one can take the kernel $\mathcal{K} = k_S$ (cf. [KS1], Corollary 7.2.2). Recall that locally χ may be decomposed as

$$\chi = \chi_1 \circ \chi_2 : \Omega_X \longrightarrow \Omega_Z \longrightarrow \Omega_Y$$

where the Lagrangian manifold Λ_i associated to the contact transformation χ_i is the conormal bundle to a smooth hypersurface S_i . By shrinking Ω_X and Ω_Y we may assume that $\pi_Z(\Omega_Z)$ is subanalytic and relatively compact. Then $k_{S_1} \circ k_{S_2 \cap (\pi_Z(\Omega_Z) \times X)}$ is $\mathbb{R}$ -constructible and satisfies (1), (2), (4) and (5) of the theorem (cf. [KS1], Corollary 7.2.2).

Moreover we know that $\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}}$ and $\Phi_{\mathcal{K}^*}^{\mathbb{R}\text{-c}}$ are well-defined and that they are quasi-inverse functors in the non-constructible case.

By definition, we have

$$\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}} \circ \Phi_{\mathcal{K}^*}^{\mathbb{R}\text{-c}} \simeq \Phi_{\mathcal{K} \circ \mathcal{K}_{\overline{V \times Y}}^*}^{\mathbb{R}\text{-c}}.$$

Recall that there is a natural isomorphism

$$k_{\Delta_X} \longrightarrow \mathcal{K} \circ \mathcal{K}^*$$

in $\text{D}^b(k_X, \Omega_X)$. Hence we get an isomorphism

$$k_{\Delta_X} \longrightarrow \mathcal{K} \circ \mathcal{K}^* \longrightarrow \mathcal{K} \circ \mathcal{K}_{\overline{V \times Y}}^* \quad (5.1.1)$$

in $\text{D}^b(k_X, \Omega_X)$. It is sufficient to prove that this morphism is well defined in $\text{D}_{\mathbb{R}\text{-c}}^b(k_X, \Omega_X)$. Denote by q_{12}, q_{13}, q_{23} the obvious projections from $X \times X \times Y$. We get a commutative diagram

$$\begin{array}{ccccc} k_{\Delta_X} & \longrightarrow & \text{R}q_{12*} \text{R}\mathcal{H}om(q_{13}^{-1} \mathcal{K}, q_{23}^! \mathcal{K}) & \longleftarrow & \mathcal{K} \circ \mathcal{K}^* \\ & \searrow & \downarrow & & \downarrow \\ & & \text{R}q_{12*} \text{R}\mathcal{H}om(q_{13}^{-1} \mathcal{K}, q_{23}^! \mathcal{K}_{\overline{V \times Y}}) & \longleftarrow & \mathcal{K} \circ \mathcal{K}_{\overline{V \times Y}}^* \end{array}$$

that is defined in $\text{D}^b(k_X)$. Note that the lower part is well-defined in $\text{D}_{\mathbb{R}\text{-c}}^b(k_X)$. All morphisms become isomorphisms in $\text{D}^b(k_X, \Omega_X)$ (cf. [KS1], Theorem 7.2.1). Since the natural functor $\text{D}_{\mathbb{R}\text{-c}}^b(k_X, \Omega_X) \rightarrow \text{D}^b(k_X, \Omega_X)$ is conservative, this shows that (5.1.1) is well-defined in $\text{D}_{\mathbb{R}\text{-c}}^b(k_X, \Omega_X)$.

Similarly one shows that the kernel $r^{-1} \text{R}\mathcal{H}om(\mathcal{K}, \omega_{X \times Y|X})$ defines a right inverse of $\mathcal{K}$ which proves that $\Phi_{\mathcal{K}}^{\mathbb{R}\text{-c}}$ is an equivalence. Then $\Phi_{\mathcal{K}^*}^{\mathbb{R}\text{-c}}$ is actually a quasi-inverse since it is a left inverse of an equivalence. $\square$

5.2 Quantization of $\mathbb{C}$ -constructible sheaves

It is now easy to transfer the results of the last section to microlocally $\mathbb{C}$ -constructible sheaves.

Consider complex manifolds X, Y of the same dimension and open $\mathbb{C}^\times$ -conic subsets $\Omega_X \subset \dot{T}^*X$, $\Omega_Y \subset \dot{T}^*Y$. We will call a $\mathbb{C}^\times$ -homogenous symplectic isomorphism

$$\chi : \Omega_X \xrightarrow{\sim} \Omega_Y$$

a contact transformation (omitting “complex” since we will never consider real contact transformations when dealing with microlocally $\mathbb{C}$ -constructible sheaves). Then we get the analogous statements of Theorems 5.0.2 and 5.1.2 by replacing open sets with $\mathbb{C}^\times$ -conic open sets. Let us give the precise formulation only in the case of microlocally $\mathbb{C}$ -constructible sheaves.

Theorem 5.2.1. *Let X, Y be two complex manifolds, $\Omega_X \subset \dot{T}^*X$, $\Omega_Y \subset \dot{T}^*Y$ open subsets and*

$$\chi : \Omega_X \longrightarrow \Omega_Y$$

a homogenous complex contact transformation. Set

$$\Lambda = \left\{ ((y; \eta), (x; \xi)) \in \Omega_Y \times \Omega_X^a \mid (y, \eta) = \chi(x, -\xi) \right\}.$$

Then Λ is $\mathbb{C}^\times$ -conic. Let $p_X \in \Omega_X$ and $p_Y = \chi(p_X)$.

*There exist open neighborhoods X' of $\pi(p_X)$, Y' of $\pi(p_Y)$, $\mathbb{C}^\times$ -conic open neighborhoods Ω'_X of $\mathbb{C}^\times p_X$, Ω'_Y of $\mathbb{C}^\times p_Y$ with $\Omega'_X \subset T^*X' \cap \Omega_X$, $\Omega'_Y \subset T^*Y' \cap \Omega_Y$ and a kernel $\mathcal{K} \in D_{\mathbb{R}-c}^b(k_{Y' \times X'})$ satisfying*

$$(1) \quad \chi \text{ induces a homogenous contact transformation } \chi : \Omega'_X \xrightarrow{\sim} \Omega'_Y,$$

(2)

$$\left((\Omega_Y'' \times T^*X') \cup (T^*Y' \times \Omega_X''^a) \right) \cap \text{SS}(\mathcal{K}) \subset \Lambda \cap (\Omega_Y'' \times \Omega_X''^a)$$

for every $\mathbb{C}^\times$ -conic open subset $\Omega_X'' \subset \Omega'_X$ and $\Omega_Y'' = \chi(\Omega_X'')$,

(3) *the functor $\Phi_{\mathcal{K}}^{\mathbb{R}-c}$ induces an equivalence of prestacks*

$$\Phi_{\mathcal{K}}^{\mathbb{C}-c} : \chi_* D_{\mathbb{C}-c}^b(k_{X'}, *)|_{\Omega'_X} \longrightarrow D_{\mathbb{C}-c}^b(k_{Y'}, *)|_{\Omega'_Y}.$$

$$(4) \quad \text{SS}(\Phi_{\mathcal{K}}^{\mathbb{C}-c}(\mathcal{F})) \cap \Omega_Y'' = \chi(\text{SS}(\mathcal{F}) \cap \Omega_X'')$$

$$(5) \quad \chi_* \mu \text{hom}(\mathcal{F}, \mathcal{G})|_{\Omega'_X} \simeq \mu \text{hom}(\Phi_{\mathcal{K}}^{\mathbb{C}-c}(\mathcal{F}), \Phi_{\mathcal{K}}^{\mathbb{C}-c}(\mathcal{G}))|_{\Omega'_Y}.$$

Proof. Since by definition $D_{\mathbb{C}-c}^b(k_X, \Omega_X'')$ is a full subcategory of $D_{\mathbb{R}-c}^b(k_X, \Omega_X'')$, it is enough to show that for any $\mathcal{F} \in D_{\mathbb{C}-c}^b(k_X, \Omega_X)$ the object $\Phi_{\mathcal{K}}^{\mathbb{R}-c}(\mathcal{F})$ is an object of $D_{\mathbb{C}-c}^b(k_Y, \Omega_Y)$. Hence we have to show that $\text{SS}(\Phi(\mathcal{F}))$ is $\mathbb{C}^\times$ -conic on Ω_Y . Since χ is $\mathbb{C}^\times$ -homogenous this is easily verified by the formula

$$\text{SS}(\Phi_{\mathcal{K}}^{\mathbb{R}-c}(\mathcal{F})) \cap \Omega_Y'' = \chi(\text{SS}(\mathcal{F}) \cap \Omega_X'').$$

□

6 Microlocally complex constructible sheaves on $\mathbb{C}^\times p$

Let $p \in \dot{T}^*X$. As a special case of Proposition 4.4.1 we get that the natural functor

$$D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times p) \longrightarrow D^b(k_X, \mathbb{C}^\times p)$$

is fully faithful. Moreover, morphisms in $D^b(k_X, \mathbb{C}^\times p)$ between microlocally $\mathbb{C}$ -constructible sheaves $\mathcal{F}, \mathcal{G} \in D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times p)$ are given by sections of $\mu\text{hom}(\mathcal{F}, \mathcal{G})$ on $\mathbb{C}^\times p$.

In this section we will give a description of the objects of $D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times p)$ using quantized contact transformation and the generic position theorem. More precisely we will show in Section 6.1 that if $\mathcal{F}$ is microlocally $\mathbb{C}$ -constructible on $\mathbb{C}^\times p$ and $\text{SS}(\mathcal{F})$ is in generic position (i.e. $\text{SS}(\mathcal{F}) \cap \mathbb{C}^\times p$ is isolated in $\pi^{-1}\pi(p)$) then $\mathcal{F}$ is isomorphic in $D^b(k_X, \mathbb{C}^\times p)$ to an object of $D_{\mathbb{C}\text{-}c}^b(k_X)$.

It will often be convenient to fix an $\mathbb{R}^+$ -conic Lagrangian variety Λ in T^*X and to consider only sheaves whose micro-support is contained in Λ . We introduce the following categories:

Definition 6.0.2. *Let $\Lambda \subset T^*X$ be an $\mathbb{R}^+$ -conic Lagrangian variety defined in a neighborhood of a subset $S \subset T^*X$. Then we define the following two categories:*

- (1) $D_{\mathbb{R}\text{-}c, \Lambda}^b(k_X, S) \subset D_{\mathbb{R}\text{-}c}^b(k_X, S)$ is the full subcategory of $D_{\mathbb{R}\text{-}c}^b(k_X, S)$ whose objects $\mathcal{F}$ satisfy $\text{SS}(\mathcal{F}) \subset \Lambda$ in a neighborhood of S .
- (2) $D_{\mathbb{C}\text{-}c, \Lambda}^b(k_X, S) = D_{\mathbb{C}\text{-}c}^b(k_X, S) \cap D_{\mathbb{R}\text{-}c, \Lambda}^b(k_X, S)$.

Of course, the second definition is only of interest when S is $\mathbb{C}^\times$ -conic and Λ is $\mathbb{C}^\times$ -conic on the subset S .

6.1 Microlocal complex constructible sheaves in generic position

Definition 6.1.1. *Let $p \in \dot{T}^*X$ and $\Lambda \subset T^*X$ be an $\mathbb{R}^+$ -conic Lagrangian subset such that Λ is $\mathbb{C}^\times$ -conic in a neighborhood of $\mathbb{C}^\times p$. We say that Λ is in generic position at p if*

$$\Lambda \cap \pi^{-1}(\pi(p)) \subset \mathbb{C}^\times p$$

in a neighborhood of $\mathbb{C}^\times p$.

Remark 6.1.2. If Λ is in generic position at p , then Λ is $\mathbb{C}^\times$ -conic in a neighborhood of $\mathbb{C}^\times p$ by definition. Hence either $\Lambda \cap \pi^{-1}(\pi(p)) \subset \mathbb{C}^\times p$ or $\Lambda \cap \pi^{-1}(\pi(p)) = \emptyset$. Moreover, being in generic position is an open property. Also note that if Λ is in generic position then it is in generic position in the sense of [KK], Chapter I.6 where Λ is supposed to be only locally $\mathbb{C}^\times$ -conic in a neighborhood of p .

Proposition 6.1.3. *Let $\Lambda \subset T^*X$ be an $\mathbb{R}^+$ -conic Lagrangian variety that is $\mathbb{C}^\times$ -conic in a neighborhood of $\mathbb{C}^\times p$. Suppose that Λ is in generic position at p . Then there exists a fundamental system of conic open subanalytic neighborhoods γ of $\mathbb{C}^\times p$ in $\dot{T}_{\pi(p)}^*X$ and for each γ a fundamental system of open relatively compact subanalytic neighborhoods U of $\pi(p)$ such that*

- (1) *the microlocal cut-off functor $\Phi_{U, \gamma} : D^b(k_X) \rightarrow D^b(k_X)$ induces a functor*

$$\Phi_{U, \gamma} : D_{\mathbb{C}\text{-}c, \Lambda}^b(k_X, U \times \gamma) \longrightarrow D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times p),$$

and this functor factors as

$$\Phi_{U, \gamma} : D_{\mathbb{C}\text{-}c, \Lambda}^b(k_X, U \times \gamma) \longrightarrow D_{\mathbb{C}\text{-}c}^b(k_X, \pi(p)) \longrightarrow D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times p).$$

- (2) There is an isomorphism of functors $\Phi_{U,\gamma} \xrightarrow{\sim} \iota$ where $\iota : D_{\mathbb{C}\text{-c},\Lambda}^b(k_X, U \times \gamma) \rightarrow D_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times p)$ is the natural functor.
- (3) $\text{SS}(\Phi_{U,\gamma}(\mathcal{F})) \cap \pi^{-1}(V) = \text{SS}(\mathcal{F}) \cap (V \times \gamma)$ for sufficiently small open neighborhoods V of $\pi(p)$.
- (4) $\text{SS}(\Phi_{U,\gamma}(\mathcal{F})) \cap \pi^{-1}\pi(p) \subset \mathbb{C}^\times p$.

Proof. Since the functor $\Phi_{U,\gamma}$ sends $D_{\mathbb{R}\text{-c}}^b(k_X)$ to $D_{\mathbb{R}\text{-c}}^b(k_X)$ (cf. Proposition 4.2.1), it induces

$$\Phi_{U,\gamma} : D_{\mathbb{R}\text{-c}}^b(k_X, U \times \gamma) \longrightarrow D_{\mathbb{R}\text{-c}}^b(k_X, U \times \gamma).$$

Since $\text{SS}(\Phi_{U,\gamma}(\mathcal{F})) \cap U \times \gamma = \text{SS}(\mathcal{F}) \cap U \times \gamma$ the functor $\Phi_{U,\gamma}$ preserves microlocally $\mathbb{C}$ -constructible sheaves and we get the functor of (1) as

$$\Phi_{U,\gamma} : D_{\mathbb{C}\text{-c}}^b(k_X, U \times \gamma) \longrightarrow D_{\mathbb{C}\text{-c}}^b(k_X, U \times \gamma) \longrightarrow D_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times p).$$

Recall that there exist a fundamental system of conic subanalytic open neighborhoods γ of $\mathbb{C}^\times p$ and for each γ a fundamental system of relatively compact subanalytic open neighborhoods U of $\pi(p)$ such that (U, γ) is a refined cutting pair. If γ is sufficiently small then $\gamma \cap \pi^{-1}\pi(p) \cap \Lambda = \mathbb{C}^\times p$. Next we choose U sufficiently small such that $U \cap \partial\gamma \cap \Lambda = \emptyset$. Then (2) follows from the refined micolocal cut-off lemma, (3) from Corollary 3.3.6 and (4) from (3).

Finally let us prove the factorization of (1). For this purpose let us first write $D_{\mathbb{C}\text{-c},\Lambda}^b(k_X, U \times \gamma)$ as a localization of a full subcategory of $D_{\mathbb{R}\text{-c}}^b(k_X)$.

Denote by $D_{\mathbb{R}\text{-c},\Lambda,U \times \gamma}^b(k_X)$ the full subcategory of $D_{\mathbb{R}\text{-c}}^b(k_X)$ such that

$$\text{SS}(\mathcal{F}) \cap U \times \gamma \subset \Lambda.$$

The category $D_{\mathbb{R}\text{-c},\Lambda,U \times \gamma}^b(k_X)$ is obviously a full triangulated subcategory. Now we localize $D_{\mathbb{R}\text{-c},\Lambda,U \times \gamma}^b(k_X)$ by complexes whose micro-support is disjoint from $U \times \gamma$ (hence by $\mathcal{N}_{\mathbb{R}\text{-c},U \times \gamma}$). Then we get a natural functor

$$D_{\mathbb{R}\text{-c},\Lambda,U \times \gamma}^b(k_X) / \mathcal{N}_{\mathbb{R}\text{-c},U \times \gamma} \longrightarrow D_{\mathbb{R}\text{-c},\Lambda}^b(k_X, U \times \gamma). \quad (6.1.1)$$

If $\mathcal{F} \in D_{\mathbb{R}\text{-c},\Lambda,U \times \gamma}^b(k_X)$ and

$$(\text{SS}(\mathcal{F}) \cap U \times \gamma) \subset (\Lambda \cap U \times \gamma),$$

then any object $\mathcal{F}'$ that is isomorphic to $\mathcal{F}$ on $U \times \gamma$ is also an object of $D_{\mathbb{R}\text{-c},\Lambda,U \times \gamma}^b(k_X)$. Hence we get that (6.1.1) is an equivalence.

By assumption Λ is $\mathbb{C}^\times$ -conic in a neighborhood of $\mathbb{C}^\times p$. Hence we may assume that Λ is $\mathbb{C}^\times$ -conic on $U \times \gamma$. One can show that if $\text{SS}(\mathcal{F}) \subset \Lambda$ then $\text{SS}(\mathcal{F})$ is $\mathbb{C}^\times$ -conic on $U \times \gamma$ (cf. Theorem 8.5.5 of [KS1]). Let us recall the idea of the proof. First one shows that $\text{SS}(\mathcal{F})$ is open in Λ (on $U \times \gamma$) and therefore locally $\mathbb{C}^\times$ -conic, i.e. for every $\mathbb{C}^\times$ -orbit S the set $\text{SS}(\mathcal{F}) \cap S \cap U \times \gamma$ is open in $S \cap U \times \gamma$ (Lemma 8.3.14 of [KS1]). Then, by Proposition 8.5.2 of [KS1], one gets that Λ is $\mathbb{C}$ -analytic on $U \times \gamma$. Hence Λ is $\mathbb{C}$ -analytic and $\mathbb{R}^+$ -conic and therefore $\mathbb{C}^\times$ -conic.

Thus, we get the equivalence

$$D_{\mathbb{R}\text{-c},\Lambda,U \times \gamma}^b(k_X) / \mathcal{N}_{\mathbb{R}\text{-c},U \times \gamma} \simeq D_{\mathbb{R}\text{-c},\Lambda}^b(k_X, U \times \gamma) \simeq D_{\mathbb{C}\text{-c},\Lambda}^b(k_X, U \times \gamma).$$

By (3) and the assumption that Λ is $\mathbb{C}^\times$ -conic in a neighborhood of $\mathbb{C}^\times p$ we get the functor

$$D_{\mathbb{R}\text{-c},\Lambda,U \times \gamma}^b(k_X) \xrightarrow{\Phi_{U,\gamma}} D_{\mathbb{C}\text{-c}}^b(k_X, \pi(p))$$

for sufficiently small (U, γ) . If $\mathcal{F} \in D_{\mathbb{R}\text{-c},\Lambda,U \times \gamma}^b(k_X) \cap \mathcal{N}_{\mathbb{R}\text{-c},U \times \gamma}$ then again by (3) $\mathcal{F}$ is zero in $D_{\mathbb{C}\text{-c}}^b(k_X, \pi(p))$. Hence we get the factorization of (1). $\square$

Lemma 6.1.4. *In the situation of Proposition 6.1.3, suppose that we have two pairs $(V, \delta) \subset (U, \gamma)$ such that (1), (2), (3), (4) are satisfied. Then the natural morphism*

$$\Phi_{V, \delta}(\mathcal{F}) \longrightarrow \Phi_{U, \gamma}(\mathcal{F})$$

is an isomorphism in $D_{\mathbb{C}-c}^b(k_X, \pi(p))$.

Proof. Embed the morphism in a distinguished triangle

$$\Phi_{V, \delta}(\mathcal{F}) \longrightarrow \Phi_{U, \gamma}(\mathcal{F}) \longrightarrow \mathcal{H} \longrightarrow$$

in $D_{\mathbb{R}-c}^b(k_X)$. We want to show that $\mathcal{H}$ is isomorphic to zero in a neighborhood of $\pi(p)$. First note that we have $\text{SS}(\mathcal{H}) \cap V \times \delta = \emptyset$ because $\Phi_{V, \delta}(\mathcal{F}) \longrightarrow \Phi_{U, \gamma}(\mathcal{F})$ is an isomorphism on $V \times \delta$. By (3), there exists an open neighborhood $V' \subset V$ of $\pi(p)$ such that

$$\text{SS}(\Phi_{V, \delta}(\mathcal{F})) \cap \dot{\pi}^{-1}(V') = \text{SS}(\mathcal{F}) \cap (V' \times \delta)$$

and

$$\text{SS}(\Phi_{U, \gamma}(\mathcal{F})) \cap \dot{\pi}^{-1}(V') = \text{SS}(\mathcal{F}) \cap (V' \times \gamma).$$

However if we take V' sufficiently small then we will show that

$$\text{SS}(\Phi_{U, \gamma}(\mathcal{F})) \cap \dot{\pi}^{-1}(V') = \text{SS}(\mathcal{F}) \cap (V' \times \delta).$$

Indeed suppose that this is not possible. Then we can construct a sequence $(x_n, \xi_n) \in \dot{T}^*X$ such that $x_n \rightarrow \pi(p)$ and $\xi_n \in \gamma \setminus \delta$ and $|\xi_n| = 1$. By extracting a convergent subsequence, we get a point $(\pi(p), \xi) \in \text{SS}(\Phi_{U, \gamma}(\mathcal{F})) \cap \{\pi(p)\} \times (\gamma \setminus \delta)$. But this is impossible because by (4) we have $\text{SS}(\Phi_{U, \gamma}(\mathcal{F})) \cap \pi^{-1}\pi(p) = \mathbb{C}^\times p$.

Hence

$$\text{SS}(\mathcal{H}) \cap \dot{\pi}^{-1}(V') \subset (\text{SS}(\Phi_{V, \delta}(\mathcal{F}) \cup \text{SS}(\Phi_{U, \gamma}(\mathcal{F}))) \cap \dot{\pi}^{-1}\pi(V') \subset V' \times \delta$$

and therefore $\text{SS}(\mathcal{H}) \cap \pi^{-1}(V') \subset T_X^*X$. But if V' is sufficiently small then

$$\text{SS}(\mathcal{H}) \cap V' \subset (\text{SS}(\Phi_{V, \delta}(\mathcal{F}) \cup \text{SS}(\Phi_{U, \gamma}(\mathcal{F}))) \cap V' \subset \pi(\Lambda) \cap V'$$

where $\pi(\Lambda)$ is a closed hypersurface and we can conclude that $\mathcal{H} \simeq 0$ by involutivity of the micro-support. $\square$

Theorem 6.1.5. *Suppose that Λ is in generic position at p . Then the microlocal cut-off functors induce a fully faithful functor*

$$\Phi : D_{\mathbb{C}-c, \Lambda}^b(k_X, \mathbb{C}^\times p) \longrightarrow D_{\mathbb{C}-c}^b(k_X, \pi(p))$$

Proof. We obviously have the equivalence

$$\varinjlim_{\mathbb{C}^\times p \in U \times \gamma} D_{\mathbb{C}-c, \Lambda}^b(k_X, U \times \gamma) \xrightarrow{\sim} D_{\mathbb{C}-c, \Lambda}^b(k_X, \mathbb{C}^\times p).$$

We may assume by cofinality that (U, γ) is a sufficiently small refined cutting pair such that Proposition 6.1.3 holds. By Lemma 6.1.4 the functors $\Phi_{U, \gamma}$ of Proposition 6.1.3 induce a functor

$$\Phi : D_{\mathbb{C}-c, \Lambda}^b(k_X, \mathbb{C}^\times p) \longrightarrow D_{\mathbb{C}-c}^b(k_X, \pi(p)).$$

Let us show that Φ is fully faithful.

Let $\mathcal{F}, \mathcal{G} \in D_{\mathbb{C}-c, \Lambda}^b(k_X, \mathbb{C}^\times p)$. Note that by Proposition 6.1.3 if $\mathcal{H} \rightarrow \mathcal{F}$ is an isomorphism

on $\mathbb{C}^\times p$ then $\Phi_{U,\gamma}(\mathcal{H}) \rightarrow \Phi_{U,\gamma}(\mathcal{F})$ is an isomorphism at $\pi(p)$ for sufficiently small (U, γ) . Consider the following chain of morphisms:

$$\begin{aligned} \mathrm{Hom}_{\mathrm{D}_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times p)}(\mathcal{F}, \mathcal{G}) &= \varinjlim_{\substack{\mathcal{H} \rightarrow \mathcal{F} \\ \text{iso on } \mathbb{C}^\times p}} \mathrm{Hom}_{\mathrm{D}^b(k_X)}(\mathcal{H}, \mathcal{G}) \\ &\longrightarrow \varinjlim_{\substack{\Phi_{U,\gamma}(\mathcal{H}) \rightarrow \Phi_{U,\gamma}(\mathcal{F}) \\ \text{iso on } \pi(p)}} \mathrm{Hom}_{\mathrm{D}^b(k_X)}(\Phi_{U,\gamma}(\mathcal{H}), \Phi_{U,\gamma}(\mathcal{G})) \\ &\longrightarrow \varinjlim_{\substack{\mathcal{H}' \rightarrow \Phi_{U,\gamma}(\mathcal{F}) \\ \text{iso on } \pi(p)}} \mathrm{Hom}_{\mathrm{D}^b(k_X)}(\mathcal{H}', \Phi_{U,\gamma}(\mathcal{G})) = \mathrm{Hom}_{\mathrm{D}_{\mathbb{C}\text{-c}}^b(k_X, \pi(p))}(\Phi_{U,\gamma}(\mathcal{F}), \Phi_{U,\gamma}(\mathcal{G})). \end{aligned}$$

We have to check that the composition is an isomorphism for a sufficiently small cutting pair (U, γ) .

Consider a morphism $\mathcal{H}' \rightarrow \Phi_{U,\gamma}(\mathcal{G})$ and an isomorphism $\mathcal{H}' \rightarrow \Phi_{U,\gamma}(\mathcal{F})$ on $\pi(p)$. Then $\mathrm{SS}(\mathcal{H}') = \mathrm{SS}(\Phi_{U,\gamma}(\mathcal{F}))$ in $\pi^{-1}(V)$ for some neighborhood V of $\pi(p)$ and in particular $\mathcal{H}' \in \mathrm{D}_{\mathbb{C}\text{-c}, \Lambda}^b(k_X, \mathbb{C}^\times p)$, hence $\Phi_{U,\gamma}(\mathcal{H}') \rightarrow \mathcal{H}'$ is an isomorphism in $\pi(p)$. Then $\mathcal{H}' \rightarrow \Phi_{U,\gamma}(\mathcal{G}) \rightarrow \mathcal{G}$ is sent to $\Phi(\mathcal{H}') \rightarrow \Phi(\mathcal{G})$ which is the same morphism in $\mathrm{D}_{\mathbb{C}\text{-c}}^b(k_X, \pi(p))$ as $\mathcal{H}' \rightarrow \Phi_{U,\gamma}(\mathcal{G})$. Hence the map is surjective.

Let us show that it is injective. If $\Phi_{U,\gamma}(\mathcal{H}) \rightarrow \Phi_{U,\gamma}(\mathcal{G})$ is zero in $\mathrm{D}_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times p)$ then there exists $\mathcal{K}$ and an isomorphism $\mathcal{K} \rightarrow \Phi_{U,\gamma}(\mathcal{H}) \rightarrow \Phi_{U,\gamma}(\mathcal{F})$ on $\pi(p)$ such that $\mathcal{K} \rightarrow \Phi(\mathcal{H}) \rightarrow \Phi(\mathcal{G})$ is the zero morphism. Then $\mathcal{K} \rightarrow \Phi(\mathcal{H}) \rightarrow \mathcal{H} \rightarrow \mathcal{F}$ is an isomorphism on $\mathbb{C}^\times p$ and $\mathcal{K} \rightarrow \Phi(\mathcal{H}) \rightarrow \mathcal{H} \rightarrow \mathcal{G}$ is zero hence $\mathcal{H} \rightarrow \mathcal{G}$ is zero in $\mathrm{D}_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times p)$. $\square$

Remark 6.1.6. Note that

$$\mathrm{D}_{\mathbb{C}\text{-c}}^b(k_*)_{\pi(p)} \xrightarrow{\sim} \mathrm{D}_{\mathbb{C}\text{-c}}^b(k_X, \pi(p)),$$

where $\mathrm{D}_{\mathbb{C}\text{-c}}^b(k_*)$ denotes the prestack $U \mapsto \mathrm{D}_{\mathbb{C}\text{-c}}^b(k_U)$. In particular we get the fully faithful functor

$$\mathrm{D}_{\mathbb{C}\text{-c}, \Lambda}^b(k_X, \mathbb{C}^\times p) \longrightarrow \mathrm{D}_{\mathbb{C}\text{-c}}^b(k_*)_{\pi(p)}.$$

Also recall that $\mathrm{D}_{\mathbb{C}\text{-c}, \Lambda}^b(k_X, \mathbb{C}^\times p)$ is additive (because of the triangular inequality for the micro-support). Hence $\mathrm{D}_{\mathbb{C}\text{-c}, \Lambda}^b(k_X, \mathbb{C}^\times p)$ is a full additive subcategory of $\mathrm{D}_{\mathbb{C}\text{-c}}^b(k_*)_{\pi(p)}$.

Therefore we have shown that if we are given $\mathcal{F} \in \mathrm{D}_{\mathbb{C}\text{-c}}^b(k_X, \mathbb{C}^\times p)$ such that $\mathrm{SS}(\mathcal{F})$ is in generic position at p then there exists an object $\tilde{\mathcal{F}} \in \mathrm{D}_{\mathbb{R}\text{-c}}^b(k_X)$ that is $\mathbb{C}$ -constructible in a neighborhood of $\pi(p)$. Moreover we can construct $\tilde{\mathcal{F}}$ in a functorial way.

6.2 The generic position theorem

Let us recall Kashiwara-Kawai's generic position theorem (cf. [KK], Chapter I.6).

Proposition 6.2.1. *Let $p \in \dot{T}^*X$ and $\Lambda \subset T^*X$ be an $\mathbb{R}^+$ -conic Lagrangian subset such that Λ is $\mathbb{C}^\times$ -conic in a neighborhood of $\mathbb{C}^\times p$. Then there exists a complex contact transformation*

$$\chi : T^*X \rightarrow T^*X,$$

defined in a neighborhood of $\mathbb{C}^\times p$, such that $\chi(\Lambda)$ is in generic position at $q = \chi(p)$.

Proof. Kashiwara-Kawai show this result on a neighborhood of p for locally $\mathbb{C}^\times$ -conic Lagrangian varieties assuming only that $\Lambda \cap \pi^{-1}\pi(p) = \mathbb{C}^\times p$ in a neighborhood of p . Hence if we suppose that Λ is $\mathbb{C}^\times$ -conic in a neighborhood of $\mathbb{C}^\times p$, we get that $\chi(\Lambda)$ is in generic position at q since χ is $\mathbb{C}^\times$ -homogenous. $\square$

Remark 6.2.2. Consider a microlocal complex constructible sheaf $\mathcal{F} \in D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times p)$. By the generic position theorem and invariance under quantized contact transformations we can find a contact transformation $\chi : T^*X \rightarrow T^*X$ defined in a neighborhood of $\mathbb{C}^\times p$ and an equivalence of categories

$$\Phi_{\mathcal{X}}^{\mathbb{C}\text{-}c} : D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times p) \xrightarrow{\sim} D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times q)$$

such that $\text{SS}(\Phi_{\mathcal{X}}^{\mathbb{C}\text{-}c}(\mathcal{F})) = \chi(\text{SS}(\mathcal{F}))$ is in generic position at q . Then $\Phi_{\mathcal{X}}^{\mathbb{C}\text{-}c}(\mathcal{F})$ is functorially isomorphic in $D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times q)$ to an object of $D_{\mathbb{C}\text{-}c}^b(*)_{\pi(q)}$. Hence many problems in $D_{\mathbb{C}\text{-}c}^b(k_X, \mathbb{C}^\times p)$ can be reduced to the study of germs of complex constructible sheaves in generic position.

7 Microlocal perverse sheaves in $D_{\mathbb{R}\text{-}c}^b(k_X, S)$

7.1 Andronikof's prestack of microlocally perverse sheaves

In this section we will first recall Andronikof's definition of the prestack of microlocal perverse sheaves. In [An2] he suggests a definition of a microlocal perverse sheaf on an arbitrary subset S of T^*X which is based on the microlocal characterisation of perverse sheaves (see Proposition 7.1.1 below). However he is not very precise concerning $\mathbb{C}^\times$ -conicity. In particular the proof of his main tool (Proposition 3.2) is incomplete under the given assumptions.

Hence we will recall a slightly more precise version of his prestack and we will restrict ourselves from the beginning to $\mathbb{C}^\times$ -conic sets. Moreover we will add complete proofs to the statements of [An2] which hold in this case. Also while Andronikof restricts his studies to points and $\mathbb{C}^\times$ -orbits we will work from the beginning with the entire prestack.

In [KS2] we find the following microlocal characterization of perverse sheaves.

Proposition 7.1.1. *Let $\mathcal{F} \in D_{\mathbb{C}\text{-}c}^b(k_X)$. Then we have equivalence between*

- (P1) $\mathcal{F}$ is a perverse sheaf.
- (P2) For every non-singular point p of $\text{SS}(\mathcal{F})$ such that the projection $\pi : \text{SS}(\mathcal{F}) \rightarrow X$ has constant rank on a neighborhood of p , there exist a submanifold $Y \subset X$ and an object $M \in \text{Mod}(k)$ such that $\mathcal{F} \simeq M_Y[\dim Y]$ in $D^b(k_X, p)$.
- (P3) The assertion (P2) is true for some point p of any irreducible component of $\text{SS}(\mathcal{F})$.

This naturally leads to the following

Definition 7.1.2. *Let $S \subset T^*X$ be a $\mathbb{C}^\times$ -conic subset.*

- (i) *A sheaf $\mathcal{F} \in D_{\mathbb{R}\text{-}c}^b(k_X)$ is called microlocally perverse on S , if it is microlocally $\mathbb{C}$ -constructible on S and there exists an open neighborhood U of S such that for every non-singular point p of $\text{SS}(\mathcal{F}) \cap U$ such that the projection $\pi : \text{SS}(\mathcal{F}) \rightarrow X$ has constant rank on a neighborhood of p , there exists a submanifold $Y \subset X$ and an object $M \in \text{Mod}(k)$ such that $\mathcal{F} \simeq M_Y[\dim Y]$ in $D^b(k_X, p)$.*
- (ii) *We will denote by $D_{\text{perv}}^b(k_X, S)$ the full subcategory of $D_{\mathbb{C}\text{-}c}^b(k_X, S)$ whose objects are perverse on S (i.e. which may be represented by a microlocally perverse sheaf on S).*
- (iii) *In particular if $\Omega \subset T^*X$ is an open $\mathbb{C}^\times$ -conic subset, we get the category $D_{\text{perv}}^b(k_X, \Omega)$. Clearly this defines a prestack of categories on P^*X , denoted by $D_{\text{perv}}^b(k_X, *)$. This is Andronikof's prestack of microlocal perverse sheaves.*

(iv) Let $\Lambda \subset T^*X$ be an $\mathbb{R}^+$ -conic Lagrangian variety that is $\mathbb{C}^\times$ -conic in a neighborhood of $\mathbb{C}^\times p$. Then we set

$$D_{\text{perv}, \Lambda}^b(k_X, \mathbb{C}^\times p) = D_{\mathbb{C}^\times, \Lambda}^b(k_X, \mathbb{C}^\times p) \cap D_{\text{perv}}^b(k_X, \mathbb{C}^\times p).$$

Proposition 7.1.3. *Let $p \in \dot{T}^*X$. The stalk of Andronikof's prestack at $\gamma(p)$ is precisely the category $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$.*

Proof. Consider the functor

$$\varinjlim_{\substack{\mathbb{C}^\times p \subset U \subset \dot{T}^*X \\ U \text{ } \mathbb{C}^\times\text{-conic}}} D_{\text{perv}}^b(k_X, U) \longrightarrow D_{\text{perv}}^b(k_X, \mathbb{C}^\times p).$$

It is essentially surjective by definition of $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$.

The proof that it is fully faithful is analogous to Proposition 3.2.5 and Corollary 3.2.6. $\square$

Proposition 7.1.4. *The duality functor*

$$D : D_{\mathbb{R}\text{-}c}^b(k_X) \longrightarrow D_{\mathbb{R}\text{-}c}^b(k_X) \quad ; \quad \mathcal{F} \mapsto R\mathcal{H}om(\mathcal{F}, \omega_X)$$

induces contravariant equivalences of prestacks

$$D : D_{\mathbb{C}\text{-}c}^b(k_X, *) \longrightarrow D_{\mathbb{C}\text{-}c}^b(k_X, *),$$

$$D : D_{\text{perv}}^b(k_X, *) \longrightarrow D_{\text{perv}}^b(k_X, *).$$

Proof. The first functor is well-defined because $\text{SS}(D\mathcal{F}) = \text{SS}(\mathcal{F})$.

Let $p \in \text{SS}(\mathcal{F})$ such that $\mathcal{F} \simeq M_Y[\dim X]$ in $D^b(k_X, p)$. Then $D\mathcal{F} \simeq D M_Y[\dim Y]$ in $D^b(k_X, p)$ since D is an anti-equivalence in $D^b(k_X, p)$. But $D M_Y[\dim Y] \simeq M_Y[\dim_Y]$. Hence the second functor is well-defined.

The two functors are equivalences because $D^2 = \text{Id}$ in $D_{\mathbb{R}\text{-}c}^b(k_X)$. $\square$

Andronikof's prestack is invariant by quantized contact transformations:

Proposition 7.1.5. *Let X, Y be two complex manifolds, $\Omega_X \subset \dot{T}^*X$, $\Omega_Y \subset \dot{T}^*Y$ open $\mathbb{C}^\times$ -conic subsets and*

$$\chi : \Omega_X \xrightarrow{\sim} \Omega_Y$$

a contact transformation. Set

$$\Lambda = \left\{ ((y; \eta), (x; \xi)) \in \Omega_Y \times \Omega_X^a \mid (y, \eta) = \chi(x, -\xi) \right\}.$$

Then Λ is $\mathbb{C}^\times$ -conic. Let $p_X \in \Omega_X$ and $p_Y = \chi(p_X)$.

*There exist open neighborhoods X' of $\pi(p_X)$, Y' of $\pi(p_Y)$, $\mathbb{C}^\times$ -conic open neighborhoods Ω'_X of $\mathbb{C}^\times p_X$, Ω'_Y of $\mathbb{C}^\times p_Y$ with $\Omega'_X \subset T^*X' \cap \Omega_X$, $\Omega'_Y \subset T^*Y' \cap \Omega_Y$ and a kernel $\mathcal{K} \in D_{\mathbb{R}\text{-}c}^b(k_{Y' \times X'})$ satisfying*

(1) χ induces a homogenous contact transformation $\Omega'_X \xrightarrow{\sim} \Omega'_Y$,

(2)

$$\left((\Omega_Y'' \times T^*X') \cup (T^*Y' \times \Omega_X''^a) \right) \cap \text{SS}(\mathcal{K}) \subset \Lambda \cap (\Omega_Y'' \times \Omega_X''^a)$$

for every $\mathbb{C}^\times$ -conic open subset $\Omega_X'' \subset \Omega'_X$ and $\Omega_Y'' = \chi(\Omega_X'')$,

(3) the functor $\Phi_{\mathcal{K}}^{\mathbb{C}^c}$ induces an equivalence of prestacks

$$\Phi_{\mathcal{K}}^{\text{perv}} : \chi_* D_{\text{perv}}^b(k_{X'}, *)|_{\Omega'_X} \longrightarrow D_{\text{perv}}^b(k_{Y'}, *)|_{\Omega'_Y}.$$

(4) $\text{SS}(\Phi_{\mathcal{K}}^{\text{perv}}(\mathcal{F})) \cap \Omega''_Y = \chi(\text{SS}(\mathcal{F}) \cap \Omega''_X)$

(5) $\chi_* \mu\text{hom}(\mathcal{F}, \mathcal{G})|_{\Omega'_X} \simeq \mu\text{hom}(\Phi_{\mathcal{K}}^{\text{perv}}(\mathcal{F}), \Phi_{\mathcal{K}}^{\text{perv}}(\mathcal{G}))|_{\Omega'_Y}.$

Proof. It is enough to show the proposition in the case in which Λ is the conormal bundle to a smooth hypersurface S and we can choose $\mathcal{K} \simeq k_S$. Then the fact that $\Phi_{\mathcal{K}}^{\mathbb{C}^c}$ preserves microlocal perverse sheaves follows from Proposition 7.4.6 of [KS1]. $\square$

7.2 The abelian category $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$

In general, one does not know much about the category $D_{\text{perv}}^b(k_X, S)$ even if S is $\mathbb{C}^\times$ -conic. In [An2], it is announced that if $p \in \dot{T}^*X$, then $D_{\text{perv}}^b(X, p)$ and $D_{\text{perv}}^b(X, \mathbb{C}^\times p)$ are abelian. While we do not know if this is true for $D_{\text{perv}}^b(X, p)$, we will give a proof here for the category $D_{\text{perv}}^b(X, \mathbb{C}^\times p)$. The main tools for the proof have been developped in Section 6.

We will fix a point $p \in \dot{T}^*X$ and an $\mathbb{R}^+$ -conic Lagrangian subvariety Λ which is $\mathbb{C}^\times$ -conic in a neighborhood of $\mathbb{C}^\times p$.

Proposition 7.2.1. *Suppose that Λ is in generic position at p . Then the functor Φ of Theorem 6.1.5 induces a commutative diagram of fully faithful functors*

$$\begin{array}{ccc} D_{\mathbb{C}^c, \Lambda}^b(k_X, \mathbb{C}^\times p) & \xrightarrow{\Phi} & D_{\mathbb{C}^c}^b(k_X)_{\pi(p)} \\ \uparrow & & \uparrow \\ D_{\text{perv}, \Lambda}^b(k_X, \mathbb{C}^\times p) & \longrightarrow & \text{Perv}(X)_{\pi(p)}. \end{array}$$

Proof. It is sufficient to prove that if $\mathcal{F} \in D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$ then $\Phi(\mathcal{F})$ is perverse in a neighborhood of $\pi(p)$. Since we have

$$\text{SS}(\Phi(\mathcal{F})) \cap \dot{\pi}^{-1}(V) = \text{SS}(\mathcal{F}) \cap (V \times \gamma)$$

for some small neighborhood $V \times \gamma$ of $\mathbb{C}^\times p$, we get the result by the characterization of perverse sheaves (cf. Proposition 7.1.1). $\square$

Also note that $D_{\text{perv}, \Lambda}^b(k_X, \mathbb{C}^\times p)$ is additive. Hence $D_{\text{perv}, \Lambda}^b(k_X, \mathbb{C}^\times p)$ is a full additive subcategory of $\text{Perv}(X)_{\pi(p)}$. In order to prove that it is actually a full abelian subcategory, we will need two lemmas:

Lemma 7.2.2. *Let $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of perverse sheaves such that $\text{SS}(\mathcal{F}) \cup \text{SS}(\mathcal{G}) \subset \Lambda$ in a neighborhood of p . Then $\text{SS}(\ker(\varphi)) \cup \text{SS}(\text{coker}(\varphi)) \subset \Lambda$ in a neighborhood at p .*

Proof. Recall ([KS1], Exercise X.6) that the micro-support of a $\mathbb{C}$ -constructible sheaf $\mathcal{F} \in D_{\mathbb{C}^c}^b(k_X)$ can be calculated as

$$\text{SS}(\mathcal{F}) = \bigcup_{i \in \mathbb{Z}} {}^p\text{H}^i(\mathcal{F})$$

where ${}^p\text{H}^i(\mathcal{F})$ denotes the i^{th} perverse cohomology sheaf of $\mathcal{F}$.

Let $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of perverse sheaves. We embed it into a distinguished triangle

$$\mathcal{F} \longrightarrow \mathcal{G} \longrightarrow \mathcal{H} \xrightarrow{+}$$

Then we consider the canonical distinguished triangle

$${}^{\mathbf{p}\tau \leq -1}(\mathcal{H}) \longrightarrow \mathcal{H} \longrightarrow {}^{\mathbf{p}\tau \geq 0}(\mathcal{H}) \xrightarrow{+}$$

where ${}^{\mathbf{p}\tau \leq -1}, {}^{\mathbf{p}\tau \geq 0}$ denote the perverse truncation functors. Then ${}^{\mathbf{p}\tau \leq -1}(\mathcal{H})[-1]$ is the kernel and ${}^{\mathbf{p}\tau \geq 0}(\mathcal{H})$ is the cokernel of φ . Therefore

$$\mathrm{SS}(\ker \varphi) \cup \mathrm{SS}(\mathrm{coker} \varphi) = \mathrm{SS}(\mathcal{H}) \subset \mathrm{SS}(\mathcal{F}) \cup \mathrm{SS}(\mathcal{G}).$$

□

Lemma 7.2.3. *Let Λ be in generic position at p .*

Consider a morphism $\mathcal{F} \rightarrow \mathcal{G}$ of $D_{\mathrm{per}, \Lambda}^b(k_X, \mathbb{C}^\times p)$. Let $\Phi(\mathcal{F}) \rightarrow \Phi(\mathcal{G})$ be the corresponding morphism in $\mathrm{Perv}_{\pi(p)}$ and $\mathcal{L} \rightarrow \Phi(\mathcal{F})$ a kernel (resp. $\Phi(\mathcal{G}) \rightarrow \mathcal{L}'$ a cokernel) in $\mathrm{Perv}_{\pi(p)}$. Then $\Phi(\mathcal{L}) \rightarrow \mathcal{L}$ (resp. $\Phi(\mathcal{L}') \rightarrow \mathcal{L}'$) is an isomorphism in $\mathrm{Perv}_{\pi(p)}$.

Proof. The proof follows the same idea as the proof of Lemma 6.1.4.

Fix U, γ such that $\Phi(\mathcal{F}) \simeq \Phi_{U, \gamma}(\mathcal{F})$, $\Phi(\mathcal{G}) \simeq \Phi_{U, \gamma}(\mathcal{G})$ and $\Phi(\mathcal{L}) \simeq \Phi_{U, \gamma}(\mathcal{L})$. Embed $\Phi_{U, \gamma}(\mathcal{L}) \rightarrow \mathcal{L}$ in a distinguished triangle

$$\Phi_{U, \gamma}(\mathcal{L}) \longrightarrow \mathcal{L} \longrightarrow \mathcal{H} \xrightarrow{+}$$

Then there is an open neighborhood $V \subset U$ of $\pi(p)$ such that

$$\mathrm{SS}(\Phi_{U, \gamma}(\mathcal{L})) \cap \dot{\pi}^{-1}(V) = \mathrm{SS}(\mathcal{L}) \cap V \times \gamma.$$

Moreover, by Lemma 7.2.2, we have

$$\mathrm{SS}(\mathcal{L}) \cap \dot{\pi}^{-1}(V) \subset \mathrm{SS}(\Phi_{U, \gamma}(\mathcal{F})) \cup \mathrm{SS}(\Phi_{U, \gamma}(\mathcal{G})) \cap \dot{\pi}^{-1}(V) \subset \Lambda \cap V \times \gamma.$$

Hence if V is sufficiently small we get

$$\mathrm{SS}(\mathcal{L}) \cap \dot{\pi}^{-1}(V) \subset \Lambda \cap V \times \gamma.$$

Since $\Phi_{U, \gamma}(\mathcal{L}) \rightarrow \mathcal{L}$ is an isomorphism on $U \times \gamma$ we get that $\mathrm{SS}(\mathcal{H}) \cap V \times \gamma = \emptyset$ and therefore $\mathrm{SS}(\mathcal{H}) \subset T_X^* X$ on V . Then

$$\mathrm{SS}(\mathcal{H}) \cap V \subset (\mathrm{SS}(\mathcal{L}) \cup \mathrm{SS}(\Phi_{U, \gamma}(\mathcal{L}))) \cap V \subset \pi(\Lambda) \cap V.$$

Since Λ is in generic position, locally at $\pi(p)$ the set $\pi(\Lambda)$ is a closed hypersurface. Therefore the involutivity of the micro-support implies that $\mathcal{H} \simeq 0$ in a neighborhood of $\pi(p)$.

The proof for the cokernel is similar. □

Proposition 7.2.4. *The additive category $D_{\mathrm{per}, \Lambda}^b(k_X, \mathbb{C}^\times p)$ is equivalent to an abelian subcategory of $\mathrm{Perv}_{\pi(p)}$.*

Proof. By Lemma 7.2.3 the full additive subcategory $D_{\mathrm{per}, \Lambda}^b(k_X, \mathbb{C}^\times p)$ is stable by kernels and cokernels. Since it is a full subcategory, it is abelian. □

Lemma 7.2.5. *Let Λ be a $\mathbb{C}^\times$ -conic Lagrangian variety (we do not ask Λ to be in generic position at p). Then $D_{\mathrm{per}, \Lambda}^b(k_X, \mathbb{C}^\times p)$ is abelian.*

Moreover if Λ' is another $\mathbb{C}^\times$ -conic Lagrangian variety with $\Lambda \subset \Lambda'$, then the natural functor

$$D_{\mathrm{per}, \Lambda}^b(k_X, \mathbb{C}^\times p) \longrightarrow D_{\mathrm{per}, \Lambda'}^b(k_X, \mathbb{C}^\times p)$$

is exact.

Proof. Consider $\Lambda \subset \Lambda'$. Let χ be a canonical transformation such that $\chi(\Lambda')$ is in generic position at $q = \chi(p)$. Then $\chi(\Lambda)$ is also at generic position at q and we get a diagram

$$\begin{array}{ccccc} D_{\text{perv}, \Lambda}^b(k_X, \mathbb{C}^\times p) & \xrightarrow{\sim} & D_{\text{perv}, \chi(\Lambda)}^b(k_X, \mathbb{C}^\times q) & \longrightarrow & \text{Perv}(X)_{\pi(q)} \\ \alpha \downarrow & & \downarrow \beta & & \downarrow = \\ D_{\text{perv}, \Lambda'}^b(k_X, \mathbb{C}^\times p) & \xrightarrow{\sim} & D_{\text{perv}, \chi(\Lambda')}^b(k_X, \mathbb{C}^\times q) & \longrightarrow & \text{Perv}(X)_{\pi(q)}. \end{array}$$

This diagram is commutative up to isomorphism. The horizontal functors are exact and fully faithful. By Proposition 7.2.4 the categories $D_{\text{perv}, \chi(\Lambda)}^b(k_X, \mathbb{C}^\times q)$ and $D_{\text{perv}, \chi(\Lambda')}^b(k_X, \mathbb{C}^\times q)$ are abelian subcategories of $\text{Perv}(X)_{\pi(q)}$. Hence β (and therefore α) is exact. $\square$

Proposition 7.2.6. *The category $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$ is abelian. Moreover, for every germ of a $\mathbb{C}^\times$ -conic Lagrangian variety $\Lambda \subset T^*X$ defined in a neighborhood of $\mathbb{C}^\times p$, the inclusion functor*

$$D_{\text{perv}, \Lambda}^b(k_X, \mathbb{C}^\times p) \longrightarrow D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$$

is exact.

Proof. We have

$$\varinjlim_{\Lambda \supset \mathbb{C}^\times p} D_{\text{perv}, \Lambda}^b(k_X, \mathbb{C}^\times p) \xrightarrow{\sim} D_{\text{perv}}^b(k_X, \mathbb{C}^\times p).$$

Filtered 2-colimits of abelian categories with exact restriction functors are abelian. $\square$

Now let us prove the following “lifting property” for kernels and cokernels of microlocal perverse sheaves:

Proposition 7.2.7. *Let $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism in $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$. Then there exists a neighborhood V of $\mathbb{C}^\times p$, objects $\mathcal{K}, \mathcal{K}' \in D_{\text{perv}}^b(k_X, V)$ and morphisms $\mathcal{K} \rightarrow \mathcal{F}$, $\mathcal{G} \rightarrow \mathcal{K}'$ in $D_{\text{perv}}^b(k_X, V)$ such that these morphisms induce kernel and cokernel of φ in $D_{\text{perv}}^b(k_X, \mathbb{C}^\times q)$ for all $q \in V$.*

Proof. Choose a Lagrangian variety Λ such that $\text{SS}(\mathcal{F}) \cup \text{SS}(\mathcal{G}) \subset \Lambda$ in a neighborhood of $\mathbb{C}^\times p$.

By the generic position theorem, we can find a contact transformation defined in a $\mathbb{C}^\times$ -conic open neighborhood of $\mathbb{C}^\times p$

$$\chi : (T^*X, \mathbb{C}^\times p) \longrightarrow (T^*X, \mathbb{C}^\times p')$$

such that $\chi(\Lambda)$ is in generic position at p' . Then $\chi(\Lambda)$ is isomorphic to the conormal bundle to a closed hypersurface in a neighborhood of $\mathbb{C}^\times p'$. In particular, there exists a $\mathbb{C}^\times$ -conic open neighborhood V' of $\chi(\Lambda)$ such that $\chi(\Lambda)$ is in generic position at any point of $\chi(\Lambda) \cap V'$.

By Proposition 7.1.5, we can assume that Λ is in generic position at any point in a $\mathbb{C}^\times$ -conic neighborhood Ω of $\mathbb{C}^\times p$.

Consider the functor

$$\Phi : D_{\text{perv}, \Lambda}^b(k_X, \mathbb{C}^\times p) \longrightarrow \text{Perv}_{\pi(p)}.$$

Choose (U, γ) such that the morphism $\Phi(\mathcal{F}) \rightarrow \Phi(\mathcal{G})$ is defined as $\Phi_{U, \gamma}(\mathcal{F}) \rightarrow \Phi_{U, \gamma}(\mathcal{G})$ in $\text{Perv}(V)$ for some small neighborhood V of $\pi(p)$. Let $\mathcal{L} \rightarrow \Phi_{U, \gamma}(\mathcal{F})$ (resp. $\Phi_{U, \gamma}(\mathcal{G}) \rightarrow \mathcal{L}'$) be a kernel (resp. a cokernel) in $\text{Perv}(V)$. Choose a $\mathbb{C}^\times$ -conic open neighborhood of $\mathbb{C}^\times p$ such that $\Omega' \subset \Omega \cap V \times \gamma$. We will show that $\mathcal{L} \rightarrow \Phi_{U, \gamma}(\mathcal{F})$ (resp. $\Phi_{U, \gamma}(\mathcal{G}) \rightarrow \mathcal{L}'$) is a

kernel (resp. a cokernel) in $D_{\text{perv},\Lambda}^b(k_X, \mathbb{C}^\times q)$ for all q in Ω' .
We have distinguished triangles in $D^b(k_X)$

$$\Phi_{U,\gamma}(\mathcal{F}) \longrightarrow \Phi_{U,\gamma}(\mathcal{G}) \longrightarrow \mathcal{H} \xrightarrow{+}$$

and

$$\mathcal{L}[1] \longrightarrow \mathcal{H} \longrightarrow \mathcal{L}' \xrightarrow{+}$$

where $\mathcal{L}$ is the kernel (resp. $\mathcal{L}'$ the cokernel) of $\Phi_{U,\gamma}(\mathcal{F}) \rightarrow \Phi_{U,\gamma}(\mathcal{G})$ in $\text{Perv}(V)$. Now let q be a point of $\Lambda \cap \Omega'$. Since Λ is in generic position at q we get the functor

$$\Phi^q : D_{\text{perv},\Lambda}^b(k_X, \mathbb{C}^\times q) \longrightarrow \text{Perv}_{\pi(q)}.$$

Applying Φ^q to the two triangles, we get distinguished triangles

$$\Phi^q \Phi_{U,\gamma}(\mathcal{F}) \longrightarrow \Phi^q \Phi_{U,\gamma}(\mathcal{G}) \longrightarrow \Phi^q(\mathcal{H}) \xrightarrow{+}$$

and

$$\Phi^q \mathcal{L}[1] \longrightarrow \Phi^q \mathcal{H} \longrightarrow \Phi^q \mathcal{L}' \xrightarrow{+}$$

Since $\Phi^q(\mathcal{L})$ and $\Phi^q(\mathcal{L}')$ are perverse in a neighborhood of $\pi(q)$ we get (by construction of kernels (resp. cokernels) in $\text{Perv}_{\pi(q)}$) that $\Phi^q(\mathcal{L}) \rightarrow \Phi^q \Phi_{U,\gamma}(\mathcal{F})$ (resp. $\Phi^q \Phi_{U,\gamma}(\mathcal{G}) \rightarrow \Phi^q(\mathcal{L}')$) is a kernel (resp. cokernel) in $D_{\text{perv},\Lambda}^b(k_X, \mathbb{C}^\times q)$. Finally since Φ^q is a cut-off functor we have $\Phi^q(\mathcal{L}) \simeq \mathcal{L}$ in $D_{\text{perv},\Lambda}^b(k_X, \mathbb{C}^\times q)$. $\square$

Theorem 7.2.8. *The stack associated to Andronikof's prestack of microlocal perverse sheaves is abelian.*

Proof. We have shown that the stalks of this additive prestack are abelian categories (Proposition 7.2.6). Further we have shown that kernels and cokernels in the stalks may be lifted to small open neighborhoods (Proposition 7.2.7). Therefore the conditions of Proposition 2.3.1 are satisfied and the stack of microlocal perverse sheaves is abelian. $\square$

8 Microlocal perverse sheaves on P^*X

8.1 The stack of microlocal perverse sheaves

We are now ready to give a first definition of the stack of microlocal perverse sheaves.

Definition 8.1.1. *The stack of microlocal perverse sheaves on P^*X is the stack associated to Andronikof's prestack.*

By Theorem 7.2.8 we know that the stack of microlocal perverse sheaves is abelian. Furthermore since the underlying prestack is invariant by quantized contact transformations (by Proposition 7.1.5) we easily get that the stack of microlocal perverse sheaves is invariant by quantized contact transformations.

We will now give an explicit description of microlocal perverse sheaves in terms of ind-sheaves. For this purpose we will construct a subprestack

$$\mu\text{Perv} \subset \gamma_*(D^b(I(k_*))|_{T^*X}).$$

Here, $\gamma_*(D^b(I(k_*))|_{T^*X})$ is the prestack of bounded derived categories of ind-sheaves on $\mathbb{C}^\times$ -conic open subsets of T^*X . Then we will show that μPerv is actually a stack and

construct a morphism $\mu : D_{\text{perv}}^b(k_X, *) \rightarrow \mu\text{Perv}$ and prove that it induces equivalences in the stalks. Hence μPerv can be identified to the stack associated to $D_{\text{perv}}^b(k_X, *)$. In particular it is an abelian stack. Then we will be able to define the microlocal Riemann-Hilbert functors using the theory of ind-sheaves.

First note that the functor μ induces a functor of prestacks

$$\mu : D_{\text{perv}}^b(k_X, *) \longrightarrow \gamma_* D^b(I(k_*)) \quad (8.1.1)$$

where we consider $D^b(I(k_*))$ as a prestack on $\dot{T}^*X$.

Definition 8.1.2. *Let $\Omega \subset P^*X$ be an open subset.*

- (1) *An object $\mathcal{F} \in D^b(I(k_{\gamma^{-1}\Omega}))$ is microlocally perverse (on Ω) if it is locally in the image of the functor (8.1.1), i.e. if for all $p \in \gamma^{-1}(\Omega)$ there exists a $\mathbb{C}^\times$ -conic neighborhood $V \supset \mathbb{C}^\times p$ and an object $\mathcal{G} \in D_{\text{perv}}^b(k_X, V)$ such that $\mu\mathcal{G}|_V \simeq \mathcal{F}|_V$.*
- (2) *We denote by $\mu\text{Perv}(\Omega)$ the full subcategory of $D^b(I(k_{\gamma^{-1}\Omega}))$ whose objects are microlocally perverse.*

Remark 8.1.3. The functor (8.1.1) induces a functor in the stalks

$$\mu : D_{\text{perv}}^b(k_X, \mathbb{C}^\times p) \longrightarrow \gamma_* D^b(I(k_*))_{\gamma(p)}$$

and the definition of a microlocal perverse sheaf is clearly equivalent to

- (1') An object $\mathcal{F} \in D^b(I(k_{\gamma^{-1}\Omega}))$ is microlocally perverse (on Ω) if for all $p \in \gamma^{-1}(\Omega)$ there exists an object $\mathcal{G} \in D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$ such that $\mu\mathcal{G} \simeq \mathcal{F}$ in $\gamma_* D^b(I(k_*))_{\gamma(p)}$.

Here, one shall keep in mind that the natural functor

$$\gamma_* D^b(I(k_*))_{\gamma(p)} \longrightarrow D^b(I(k_{\mathbb{C}^\times p}))$$

is not fully faithful. Therefore germs of microlocal perverse sheaves should not be interpreted as complexes of ind-sheaves on $\mathbb{C}^\times p$.

Remark 8.1.4. Note that it is possible (but less obvious) to define the functor

$$\mu : D_{\text{perv}}^b(k_X, \mathbb{C}^\times p) \longrightarrow \gamma_* D^b(I(k_*))_{\gamma(p)}$$

directly without using the prestack $D_{\text{perv}}^b(k_X, *)$. Hence, we see by definition (1') that in order to define microlocal perverse sheaves, it is not necessary to construct the categories $D_{\text{perv}}^b(k_X, S)$ for any $\mathbb{C}^\times$ -conic subset $S \subset T^*X$ but only the categories $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$ for any $p \in \dot{T}^*X$. Recall that the categories $D_{\mathbb{R}\text{-c}}^b(k_X, \mathbb{C}^\times p)$ can be identified with the full subcategory of $D^b(k_X, \mathbb{C}^\times p)$ whose objects may be represented by $\mathbb{R}$ -constructible sheaves. Hence the construction of microlocal perverse sheaves is independent of the conceptual question of the “correct” definition of the microlocalization of $\mathbb{R}$ -constructible sheaves (cf. Section 4).

However, the prestack $D_{\text{perv}}^b(k_X, *)$ can sometimes be useful to define functors on microlocal perverse sheaves and therefore we decided to include it in our presentation.

Clearly $\mu\text{Perv}(\Omega)$ is an additive subcategory of $D^b(I(k_{\gamma^{-1}(\Omega)}))$, and the correspondance

$$P^*X \supset \Omega \mapsto \mu\text{Perv}(\Omega)$$

defines an additive prestack on P^*X . Hence $\mu Perv$ is a full additive subprestack of $\gamma_* D^b(I(k_*))$, and we can see directly from the construction that it is defined by a local property. Moreover, by definition, for any open subset $\Omega \subset P^*X$ the functor μ induces a natural functor

$$D_{\text{perv}}^b(k_X, \gamma^{-1}(\Omega)) \longrightarrow \mu Perv(\Omega).$$

These functors define a functor of prestacks

$$\mu : D_{\text{perv}}^b(k_X, *) \longrightarrow \mu Perv. \quad (8.1.2)$$

We want to show that the functor (8.1.2) induces equivalences of categories in the stalks. Let $p \in T^*X$. The definition of $\mu Perv$ immediately implies that

$$\mu : D_{\text{perv}}^b(k_X, \mathbb{C}^\times p) \longrightarrow \mu Perv_{\gamma(p)}$$

is essentially surjective. In order to prove that it is an equivalence of categories, we first have to calculate the morphisms in $\mu Perv_{\gamma(p)}$.

Proposition 8.1.5.

- (i) Let $\mathcal{F}, \mathcal{G} \in \mu Perv(\Omega)$ and $p \in \gamma^{-1}(\Omega)$. Let $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{G}}$ be two objects of $D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$ such that $\mu\tilde{\mathcal{F}} \simeq \mathcal{F}$ and $\mu\tilde{\mathcal{G}} \simeq \mathcal{G}$ in $\mu Perv_{\gamma(p)}$. Then we have

$$\text{Hom}_{\mu Perv_{\gamma(p)}}(\mathcal{F}, \mathcal{G}) \simeq H^0(\mathbb{C}^\times p, \mu\text{hom}(\tilde{\mathcal{F}}, \tilde{\mathcal{G}})).$$

- (ii) Let $\mathcal{F}, \mathcal{G} \in D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$. Then

$$\text{Hom}_{\mu Perv_{\gamma(p)}}(\mu\mathcal{F}, \mu\mathcal{G}) \simeq \text{Hom}_{D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)}(\mathcal{F}, \mathcal{G}).$$

- (iii) The functor μ induces a canonical equivalence of categories

$$D_{\text{perv}}^b(k_X, \mathbb{C}^\times p) \xrightarrow{\sim} \mu Perv_{\gamma(p)}.$$

Proof. Note that the functor of (iii) is obviously essentially surjective. Hence (iii) follows from (ii). Moreover (ii) follows from (i) and Proposition 3.4.4.

Let us prove (i). We have

$$\begin{aligned} \text{Hom}_{\mu Perv_{\gamma(p)}}(\mathcal{F}, \mathcal{G}) &\simeq \varinjlim_{\gamma(p) \in V \subset P^*X} \text{Hom}_{D^b(I(k_{\gamma^{-1}(V)}))}(\mathcal{F}|_{\gamma^{-1}(V)}, \mathcal{G}|_{\gamma^{-1}(V)}) \\ &\simeq \varinjlim_{\gamma(p) \in V \subset P^*X} \text{Hom}_{D^b(I(k_{\gamma^{-1}(V)}))}(\mu\tilde{\mathcal{F}}|_{\gamma^{-1}(V)}, \mu\tilde{\mathcal{G}}|_{\gamma^{-1}(V)}) \\ &\simeq \varinjlim_{\gamma(p) \in V \subset P^*X} H^0(V, \text{RHom}(\mu\tilde{\mathcal{F}}, \mu\tilde{\mathcal{G}})) \\ &\simeq \varinjlim_{\gamma(p) \in V \subset P^*X} H^0\Gamma(V, \mu\text{hom}(\tilde{\mathcal{F}}, \tilde{\mathcal{G}})) \\ &\simeq H^0(\mathbb{C}^\times p, \mu\text{hom}(\tilde{\mathcal{F}}, \tilde{\mathcal{G}})). \end{aligned}$$

□

Proposition 8.1.5 implies that if $\mu Perv$ is a stack then it is equivalent to the stack associated to $D_{\text{perv}}^b(k_X, *)$. In order to prove that $\mu Perv$ is a stack we will apply the general results of Section 2.

Let us recall the following well-known proposition with proof.

Proposition 8.1.6. *Let $\mathcal{F}, \mathcal{G}$ be two perverse sheaves on X . Then $\mu\text{hom}(\mathcal{F}, \mathcal{G})[d_X]$ is a perverse sheaf. In particular it is concentrated in positive degrees.*

Proof. According to [KS1], Corollary 10.3.20, $\mu\text{hom}(\mathcal{F}, \mathcal{G})[d_X]$ is a perverse sheaf on T^*X . Hence for any complex analytic subset S of T^*X we have

$$H_S^{j+d_X}(\mu\text{hom}(\mathcal{F}, \mathcal{G}))|_S \simeq 0$$

if $j < -\dim S$. Now recall that

$$\text{supp}(\mu\text{hom}(\mathcal{F}, \mathcal{G})) \subset \text{SS}(\mathcal{F}) \cup \text{SS}(\mathcal{G}).$$

Since $\mathcal{F}, \mathcal{G}$ are perverse sheaves their micro-supports are Lagrangian subsets of T^*X , hence are of dimension d_X . Therefore

$$H^j(\mu\text{hom}(\mathcal{F}, \mathcal{G})) \simeq 0$$

for $j < 0$. □

Proposition 8.1.7. *The prestack μPerv of microlocal perverse sheaves is separated.*

Proof. Let $\mathcal{F}, \mathcal{G}$ be two microlocal perverse sheaves of $\mu\text{Perv}(\Omega)$. Recall that $\mathcal{F}, \mathcal{G} \in D^b(I(k_{\gamma^{-1}(\Omega)}))$ and that we have by definition

$$\mathcal{H}om_{\mu\text{Perv}}(\mathcal{F}, \mathcal{G}) \simeq \gamma_* \mathcal{H}om_{D^b(I(k_*))}(\mathcal{F}, \mathcal{G}).$$

Hence it is sufficient to prove that $\mathcal{H}om_{D^b(I(k_*))}(\mathcal{F}, \mathcal{G})$ is a sheaf.

We will first show that the complex $R\mathcal{H}om(\mathcal{F}, \mathcal{G})$ is concentrated in positive degrees. This is a local question, hence we may assume that $\mathcal{F} \simeq \mu\tilde{\mathcal{F}}, \mathcal{G} \simeq \mu\tilde{\mathcal{G}}$ for two objects $\tilde{\mathcal{F}}, \tilde{\mathcal{G}}$ of $D_{\text{perv}}^b(k_X, \gamma^{-1}(\Omega))$. Since

$$R\mathcal{H}om(\mu(\tilde{\mathcal{F}}), \mu(\tilde{\mathcal{G}})) \simeq \mu\text{hom}(\tilde{\mathcal{F}}, \tilde{\mathcal{G}})$$

we are reduced to study $\mu\text{hom}(\tilde{\mathcal{F}}, \tilde{\mathcal{G}})_p$ for any $p \in \gamma^{-1}(\Omega)$. By invariance of quantized contact transformation (cf. Theorem 7.1.5) we may assume that $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{G}}$ are perverse sheaves on a neighborhood of $\pi(p)$. Hence $\mu\text{hom}(\tilde{\mathcal{F}}, \tilde{\mathcal{G}})_p$ is concentrated in positive degrees by the last proposition.

Therefore $R\mathcal{H}om(\mathcal{F}, \mathcal{G})$ is concentrated in positive degrees. Then $\mathcal{H}om_{D^b(I(k_*))}(\mathcal{F}, \mathcal{G})$ is a sheaf since

$$\mathcal{H}om_{D^b(I(k_*))}(\mathcal{F}, \mathcal{G})(V) \simeq H^0(V, R\mathcal{H}om(\mathcal{F}, \mathcal{G})) \simeq \Gamma(V, H^0 R\mathcal{H}om(\mathcal{F}, \mathcal{G})).$$

□

Theorem 8.1.8. *The prestack μPerv on P^*X is an abelian stack. Moreover the functor μ induces an equivalence of abelian stacks*

$$D_{\text{perv}}^b(k_X, *)^{\ddagger} \xrightarrow{\sim} \mu\text{Perv},$$

where $D_{\text{perv}}^b(k_X, *)^{\ddagger}$ denotes the stack associated to $D_{\text{perv}}^b(k_X, *)$.

Proof. Since microlocal perverse sheaves form a separated subprestack of the prestack of ind-sheaves and are obviously defined by a local property, they form a stack.

Moreover, Proposition (8.1.5) states that the functor of prestacks

$$\mu : D_{\text{perv}}^b(k_X, *) \longrightarrow \mu\text{Perv}$$

induces equivalences of categories in the stalks. Hence μ identifies μPerv with the stack associated to $D_{\text{perv}}^b(k_X, *)$. □

8.2 Autoduality

Proposition 8.2.1. *Let $\mathcal{F}, \mathcal{G} \in \mu Perv(\Omega)$. Then*

$$R\mathcal{H}om(\mathcal{F}, \mathcal{G})[d_X]$$

is a perverse sheaf on $\gamma^{-1}(\Omega)$.

Proof. Locally we can find $\tilde{\mathcal{F}}, \tilde{\mathcal{G}} \in D_{\mathbb{R}\text{-c}}^b(k_X)$ such that

$$R\mathcal{H}om(\mathcal{F}, \mathcal{G}) \simeq \mu hom(\tilde{\mathcal{F}}, \tilde{\mathcal{G}})$$

By invariance of quantized contact transformations we may assume that $\tilde{\mathcal{F}}, \tilde{\mathcal{G}}$ are perverse sheaves. Then the result follows from the fact that $\mu hom(\tilde{\mathcal{F}}, \tilde{\mathcal{G}})[d_X]$ is a perverse sheaf. $\square$

Proposition 8.2.2. *There is a contravariant functor*

$$D : \mu Perv \longrightarrow \mu Perv$$

such that

$$(i) \quad D \circ D \simeq \text{Id}.$$

$$(ii) \quad R\mathcal{H}om(\mathcal{F}, \mathcal{G}) \simeq R\mathcal{H}om(D\mathcal{G}, D\mathcal{F}).$$

(iii) *If $\mathcal{F} \in \mu Perv(\Omega)$ is isomorphic to $\mu\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}} \in D_{\text{perv}}^b(k_X, \gamma^{-1}(\Omega))$ then we have a natural isomorphism $D\mathcal{F} \simeq \mu R\mathcal{H}om(\mathcal{F}, \omega_X)$.*

(iv) *The stack $\mu Perv$ is autodual, i.e. it is equivalent to its opposite stack.*

Proof. Recall that the functor $D = R\mathcal{H}om(\cdot, \omega_X)$ induces a contravariant equivalence of prestacks

$$D : D_{\text{perv}}^b(k_X, *) \longrightarrow D_{\text{perv}}^b(k_X, *).$$

Hence we get a contravariant equivalence D on the stack associated to $D_{\text{perv}}^b(k_X, *)$ which satisfies by definition (i) and (iii).

Let $\mathcal{F}, \mathcal{G} \in \mu Perv(\Omega)$. It is enough to prove (ii) locally. Hence we may assume that there are objects $\tilde{\mathcal{F}}, \tilde{\mathcal{G}} \in D_{\text{perv}}^b(k_X, \gamma^{-1}\Omega)$ such that $\mu\tilde{\mathcal{F}} \simeq \mathcal{F}$ and $\mu\tilde{\mathcal{G}} \simeq \mathcal{G}$. Recall (cf. Exercise IV.4 of [KS1]) that on $\gamma^{-1}\Omega$ we have

$$\mu hom(\tilde{\mathcal{F}}, \tilde{\mathcal{G}}) \simeq \mu hom(D\tilde{\mathcal{G}}, D\tilde{\mathcal{F}}).$$

Then on $\gamma^{-1}\Omega$ we get

$$R\mathcal{H}om(\mu\tilde{\mathcal{F}}, \mu\tilde{\mathcal{G}}) \simeq \mu hom(\tilde{\mathcal{F}}, \tilde{\mathcal{G}}) \simeq \mu hom(D\tilde{\mathcal{G}}, D\tilde{\mathcal{F}}) \simeq R\mathcal{H}om(\mu D\tilde{\mathcal{G}}, \mu D\tilde{\mathcal{F}}) \simeq R\mathcal{H}om(D\mathcal{G}, D\mathcal{F}).$$

$\square$

8.3 Integral transforms for microlocal perverse sheaves

Let $p \in \Omega_X \subset T^*X$, $q \in \Omega_Y \subset T^*Y$ where Ω_X, Ω_Y are $\mathbb{C}^\times$ -conic open subsets. Suppose that we are given a contact transformation

$$\chi : \Omega_X \xrightarrow{\sim} \Omega_Y$$

such that $\chi(p) = q$. We have seen in Section 6 that after shrinking Ω_X and Ω_Y , we can establish an equivalence of prestacks

$$\Phi_{\mathcal{K}}^{\text{perv}} : \chi_* D_{\text{perv}}^b(k_X, *)|_{\Omega_X} \xrightarrow{\sim} D_{\text{perv}}^b(Y, *)|_{\Omega_Y}.$$

Hence we get an equivalence of the associated stacks of microlocal perverse sheaves:

$$\chi_* \mu Perv^X|_{\gamma(\Omega_X)} \xrightarrow{\sim} \mu Perv^Y|_{\gamma(\Omega_Y)} \quad (8.3.1)$$

together with a commutative diagram

$$\begin{array}{ccc} \chi_* D_{\text{perv}}^b(k_X, *)|_{\Omega_X} & \xrightarrow[\sim]{\Phi_{\mathcal{K}}^{\text{perv}}} & D_{\text{perv}}^b(Y, *)|_{\Omega_Y} \\ \downarrow & & \downarrow \\ \chi_* \mu Perv^X|_{\gamma(\Omega_X)} & \longrightarrow & \mu Perv^Y|_{\gamma(\Omega_Y)} \end{array}$$

The existence of this diagram is sufficient for many applications. However, in Section 9, we need to know that the equivalence (8.3.1) is given by integral transform with the kernel $\mu\mathcal{K}$.

Let us fix some notations first. We will first consider real manifolds X, Y, Z and the diagram

$$\begin{array}{ccccc} & & X \times Y \times Z & & \\ & \swarrow q_{12} & \downarrow q_{13} & \searrow q_{23} & \\ X \times Y & & X \times Z & & Y \times Z. \end{array}$$

Composition (or integral transforms) will be considered in its ind-version. For $\mathcal{F} \in D^b(I(k_{X \times Y}))$ and $\mathcal{G} \in D^b(I(k_{Y \times Z}))$ we set

$$\mathcal{F} \circ \mathcal{G} = Rq_{13!!}(q_{12}^{-1}\mathcal{F} \otimes q_{23}^{-1}\mathcal{G}),$$

even if $\mathcal{F}, \mathcal{G}$ are actually complexes of classical sheaves. In this case $\alpha(\mathcal{F} \circ \mathcal{G})$ gives the classical composition. Note that there is always a natural morphism

$$\mathcal{F} \circ \mathcal{G} \longrightarrow \alpha(\mathcal{F} \circ \mathcal{G}).$$

Furthermore, we will consider a variant. Consider

$$q_{12}^a : T^*X \times T^*Y \times T^*Z \rightarrow T^*X \times T^*Y \quad ; \quad ((x; \xi), (y; \eta), (z; \zeta)) \mapsto ((x; \xi), (y; \eta)).$$

Then set for $\mathcal{F} \in D^b(I(k_{T^*X \times T^*Y}))$ and $\mathcal{G} \in D^b(I(k_{T^*Y \times T^*Z}))$

$$\mathcal{F} \circ^a \mathcal{G} = Rq_{13!!}(q_{12}^{a-1}\mathcal{F} \otimes q_{23}^{-1}\mathcal{G}).$$

Note that the operation $\circ^a$ is associative up to natural isomorphism.

Recall that the stack $\mu Perv$ is embedded into the prestack $\gamma_* D^b(I(k_*))$.

Hence, for any kernel $\mathcal{K} \in D^b(k_{Y \times X})$ the functor

$$\Phi_{\mu\mathcal{K}}^a = \mu\mathcal{K} \circ^a : \gamma_* D^b(I(k_*))|_{\Omega_X} \longrightarrow \gamma_* D^b(I(k_*))|_{\Omega_Y}$$

is well defined. Our interest in this section is to find kernels $\mathcal{K}$ such that this functor preserves the subprestack $\mu Perv$. For this purpose, we may use the fact that microlocal perverse sheaves are defined by a local property. Hence, if $\mathcal{F} \in \mu Perv^X$ then $\Phi_{\mu\mathcal{K}}^a(\mathcal{F}) \in$

$\mu Perv^Y$ if and only it is microlocally perverse on $\mathbb{C}^\times p$ for every $p \in \dot{T}^*Y$.
Take an object $\mathcal{F} \in \mu Perv^X$. Locally $\mathcal{F} \simeq \mu \tilde{\mathcal{F}}$ for some object $\tilde{\mathcal{F}} \in D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$. Hence locally

$$\mu \mathcal{K} \overset{a}{\circ} \mathcal{F} \simeq \mu \mathcal{K} \overset{a}{\circ} \mu \tilde{\mathcal{F}}.$$

Since $\mathcal{K} \circ \tilde{\mathcal{F}} \in D_{\text{perv}}^b(Y, \mathbb{C}^\times q)$, if we know that there is an isomorphism

$$\mu \mathcal{K} \overset{a}{\circ} \mu \tilde{\mathcal{F}} \simeq \mu(\mathcal{K} \circ \tilde{\mathcal{F}}) \quad (8.3.2)$$

in a neighborhood of $\mathbb{C}^\times q$ then we have shown that the object $\mu \mathcal{K} \overset{a}{\circ} \mathcal{F}$ lies in $\mu Perv^Y$. Kashiwara's composition formula states that the morphism (8.3.2) exists and gives a criterion for it to be an isomorphism. Hence it allows us to a certain extent to translate results from the microlocal study of sheaves from [KS1] to the study of ind-sheaves and in particular to microlocal perverse sheaves. Let us recall Kashiwara's theorem.

Theorem 8.3.1 (Microlocal composition of kernels).

Let $\mathcal{K}_1 \in D^b(I(k_{X \times Y}))$ and $\mathcal{K}_2 \in D^b(I(k_{Y \times Z}))$.

(1) *There is a natural morphism*

$$\mu_{X \times Y} \mathcal{K}_1 \overset{a}{\circ} \mu_{Y \times Z} \mathcal{K}_2 \longrightarrow \mu_{X \times Z}(\mathcal{K}_1 \circ \mathcal{K}_2). \quad (8.3.3)$$

(2) *Assume the non-characteristic condition⁹*

$$\text{SS}_0(\mathcal{K}_1) \overset{a}{\times}_{T^*Y} \text{SS}_0(\mathcal{K}_2) \cap (T_X^*X \times T^*Y \times T_Z^*Z) \subset T_X^*X \times T_Y^*Y \times T_Z^*Z,$$

then the morphism

$$\text{K}_{X \times Z} \circ (\mu_{X \times Y} \mathcal{K}_1 \overset{a}{\circ} \mu_{Y \times Z} \mathcal{K}_2) \longrightarrow \mu_{X \times Z}(\mathcal{K}_1 \circ \mathcal{K}_2)$$

is an isomorphism and

$$\mu_{X \times Y} \mathcal{K}_1 \overset{a}{\circ} \mu_{Y \times Z} \mathcal{K}_2 \longrightarrow \mu_{X \times Z}(\mathcal{K}_1 \circ \mathcal{K}_2)$$

*is an isomorphism outside $p_{13}(\text{SS}_0(\mathcal{K}_1) \overset{a}{\times}_{T^*Y} \text{SS}_0(\mathcal{K}_2) \cap T^*X \times T_Y^*Y \times T_Z^*Z)$.*

Remark 8.3.2. The “ideal situation” to apply Kashiwara's theorem is given by condition (1) or (2) below.

$$(1) (T_X^*X \times T^*Y \cup T^*X \times T_Y^*Y) \cap \text{SS}(\mathcal{K}_1) \subset T_X^*X \times T_Y^*Y,$$

$$(2) (T_Y^*Y \times T^*Z \cup T^*Y \times T_Z^*Z) \cap \text{SS}(\mathcal{K}_2) \subset T_Y^*Y \times T_Z^*Z.$$

Then the natural morphism (8.3.3) is an isomorphism outside the zero section.

In particular, let $\mathcal{F} \in D^b(I(k_X))$ and consider $\mathcal{K}$ such that

$$\text{SS}(\mathcal{K}) \cap (T_Y^*Y \times T^*X \cup T^*Y \cup T_X^*X) \subset T_Y^*Y \times T_X^*X$$

Then the morphism (8.3.3)

$$\mu(\mathcal{K}) \overset{a}{\circ} \mu \mathcal{F} \longrightarrow \mu(\mathcal{K} \circ \mathcal{F})$$

is an isomorphism outside the zero section. For instance we get the following lemma.

⁹For two sets $S_1 \subset T^*X \times T^*Y$ and $S_2 \subset T^*Y \times T^*Z$ we denote by $S_1 \overset{a}{\times}_{T^*Y} S_2$ the cartesian product of $q_2^a|_{S_1} : S_1 \rightarrow T^*Y$ and $q_1|_{S_2} : S_2 \rightarrow T^*Y$, hence

$$S_1 \overset{a}{\times}_{T^*Y} S_2 = \left\{ ((x; \xi_x), (y; \xi_y), (z; \xi_z)) \in T^*X \times T^*Y \times T^*Z \mid ((x; \xi_x), (y; \xi_y)) \in S_1, ((y; -\xi_y), (z; \xi_z)) \in S_2 \right\}.$$

Lemma 8.3.3. *Let $\mathcal{F} \in D^b(I(k_X))$. The morphism*

$$\mu(k_\Delta) \overset{a}{\circ} \mu\mathcal{F} \longrightarrow \mu(k_\Delta \circ \mathcal{F}) \simeq \mu\mathcal{F}$$

is an isomorphism outside the zero section.

We will mostly be interested in the following more general situation:

Definition 8.3.4. *Let us denote by $\tilde{N}(Y, X, \Omega_Y, \Omega_X)$ the full subcategory of $D^b(k_{Y \times X}, \Omega_Y \times T^*X)$ such that*

$$(i) \quad SS(\mathcal{K}) \cap (\Omega_Y \times T^*X \cup T^*Y \times \Omega_X^a) \subset \Omega_Y \times \Omega_X^a$$

$$(ii) \quad p_1 : SS(\mathcal{K}) \cap \Omega_Y \times T^*X \rightarrow \Omega_Y \text{ is proper.}$$

Note that $\tilde{N}(Y, X, \Omega_Y, \Omega_X) \subset N(Y, X, \Omega_Y, \Omega_X)^{10}$ and that the kernels produced in Theorem 5.0.2 are objects of $\tilde{N}(Y, X, \Omega_Y, \Omega_X)$. We get

Proposition 8.3.5. *Suppose that $\Omega_X \cap T_X^*X = \emptyset$ and $\Omega_Y \cap T_Y^*Y = \emptyset$.*

*Let $\mathcal{K} \in \tilde{N}(Y, X, \Omega_Y, \Omega_X)$ and $\mathcal{F} \in D^b(k_X)$ such that $SS(\mathcal{F}) \cap T^*X \subset \Omega_X$. Then the natural morphism*

$$\mu\mathcal{K} \overset{a}{\circ} \mu\mathcal{F} \longrightarrow \mu(\mathcal{K} \circ \mathcal{F})$$

is an isomorphism on Ω_Y .

Proof. We have by hypothesis

$$\begin{aligned} SS(\mathcal{K}) \cap (T^*Y \times SS(\mathcal{F})^a) &\subset (SS(\mathcal{K}) \cap T^*Y \times \Omega_X^a) \cup (SS(\mathcal{K}) \cap T^*Y \times T_X^*X) \\ &\subset \Omega_Y \times \Omega_X^a \cup T^*Y \times T_X^*X. \end{aligned}$$

Intersecting both sides with $T_Y^*Y \times T^*X$ we get

$$SS(\mathcal{K}) \cap (T^*Y \times SS(\mathcal{F})^a) \cap (T_Y^*Y \times T^*X) \subset T_X^*X \times T_Y^*Y.$$

Hence the non-characteristic condition is satisfied.

Moreover by assumption $SS(\mathcal{K}) \cap (\Omega_Y \times T^*X) \subset \Omega_Y \times \Omega_X^a$, hence

$$SS(\mathcal{K}) \cap (T^*Y \times SS(\mathcal{F})^a) \cap \Omega_Y \times T_X^*X = \emptyset$$

and

$$SS(\mathcal{K}) \cap (T^*Y \times SS(\mathcal{F})^a) \cap T^*Y \times T_X^*X \subset \mathbb{L}\Omega_Y \times T_X^*X$$

Therefore the morphism is an isomorphism outside $\mathbb{L}\Omega_Y$, hence on Ω_Y . $\square$

Lemma 8.3.6. *Consider $\mathcal{K} \in N(Y, X, \Omega_Y, \Omega_X)$ and $\mathcal{F} \in D^b(k_X)$. Then we have a chain of natural isomorphisms:*

$$(\mu\mathcal{K})|_{\Omega_Y \times \Omega_X^a} \overset{a}{\circ} (\mu\mathcal{F})|_{\Omega_X} \simeq (\mu\mathcal{K})^{(1,a)}|_{\Omega_Y \times \Omega_X} \circ (\mu\mathcal{F})|_{\Omega_X} \simeq (\mu\mathcal{K} \overset{a}{\circ} \mu\mathcal{F})|_{\Omega_Y}.$$

Proof. We will show that the three terms are isomorphic to

$$(\mu\mathcal{K})|_{\Omega_Y \times T^*X} \overset{a}{\circ} \mathcal{F}.$$

For the right hand side this is trivial.

For the two terms on the left hand side denote by $j : \Omega_Y \times \Omega_X^a \hookrightarrow \Omega_Y \times X$ the inclusion map. Then we have

$$(\mu\mathcal{K})|_{\Omega_Y \times T^*X} \simeq Rj_! j^{-1} \mu\mathcal{K}|_{\Omega_Y \times T^*X}$$

by the hypothesis on $SS(\mathcal{K}) = \text{supp}(\mu\mathcal{K})$ and the result follows easily. $\square$

¹⁰The category $N(Y, X, \Omega_Y, \Omega_X)$ has been recalled at the beginning of Section 5.1.

Remark 8.3.7. Let $\mathcal{K} \in \tilde{\mathcal{N}}(Y, X, \Omega_Y, \Omega_X)$. Consider the morphism

$$(\mu\mathcal{K} \overset{a}{\circ} \mu\mathcal{F})|_{\Omega_Y} \longrightarrow \mu(\mathcal{K} \circ \mathcal{F})|_{\Omega_Y} \longrightarrow \mu(\alpha(\mathcal{K} \circ \mathcal{F}))|_{\Omega_Y} \quad (8.3.4)$$

The term on the right only depends on the image of $\mathcal{F}$ in $D^b(k_X, \Omega_X)$. By the last lemma, the same result holds for the term on the left side¹¹. In the following we will be interested in situations when the composition (8.3.4) is an isomorphism.

Hence, in the situation of Proposition 8.3.5, we may replace $\mathcal{F}$ by any object isomorphic to $\mathcal{F}$ in $D^b(k_X, \Omega_X)$. However, in general, it is not possible to find an object $\mathcal{F}'$ isomorphic to $\mathcal{F}$ in $D^b(k_X, \Omega_X)$ such that $\text{SS}(\mathcal{F}') \subset \Omega_X$. Therefore we will have to assume a stability condition on $\mathcal{K}$ and the existence of a suitable cut-off functor.

Proposition 8.3.8. *Let X, Y be affine and $\mathcal{K} \in \tilde{\mathcal{N}}(Y, X, \Omega_Y, \Omega_X)$.*

Let $\delta \subset X^$ be an open cone and $U \subset X$ be a relatively compact open set such that*

$$(\overline{U} \times \overline{\delta}) \cap \dot{T}^*X \subset \Omega_X.$$

Set $\Omega'_X = U \times \delta$ and suppose that there exists an open subset $\Omega'_Y \subset \Omega_Y$ such that $\mathcal{K} \in \tilde{\mathcal{N}}(Y, X, \Omega'_Y, \Omega'_X)$.

Then the natural morphism (8.3.4)

$$\mu\mathcal{K} \overset{a}{\circ} \mu\mathcal{F} \longrightarrow \mu(\alpha(\mathcal{K} \circ \mathcal{F}))$$

is an isomorphism on Ω'_Y .

Proof. There is a cut-off functor $\Phi_{U,\delta}$ such that $\text{SS}(\Phi_{U,\delta}(\mathcal{F})) \subset \overline{U} \times \overline{\delta}$ and $\Phi_{U,\delta}(\mathcal{F}) \rightarrow \mathcal{F}$ is an isomorphism in $D^b(k_X, U \times \delta)$.

Then the morphism

$$\mu\mathcal{K} \overset{a}{\circ} \mu(\Phi_{U,\delta}(\mathcal{F})) \longrightarrow \mu(\mathcal{K} \circ \Phi_{U,\delta}(\mathcal{F}))$$

is an isomorphism on Ω'_Y . Moreover $\text{supp}(\Phi_{U,\gamma}(\mathcal{F})) \subset \overline{U}$ which is compact. Hence $\mathcal{K} \circ \Phi_{U,\gamma}(\mathcal{F}) \rightarrow \alpha(\mathcal{K} \circ \Phi_{U,\gamma}(\mathcal{F}))$ is an isomorphism. Therefore

$$\mu\mathcal{K} \overset{a}{\circ} \mu(\Phi_{U,\delta}(\mathcal{F})) \longrightarrow \mu(\alpha(\mathcal{K} \circ \Phi_{U,\delta}(\mathcal{F})))$$

is an isomorphism on Ω'_Y and is isomorphic to

$$\mu\mathcal{K} \overset{a}{\circ} \mu\mathcal{F} \longrightarrow \mu(\alpha(\mathcal{K} \circ \mathcal{F})).$$

□

Summarizing the situation, we have the following diagram:

$$\begin{array}{ccc} D^b(k_X, \Omega_X) & \xrightarrow{\mu} & D^b(I(k_{\Omega_X})) \\ \Phi_{\mathcal{K}} \downarrow & \swarrow & \downarrow \mu(\mathcal{K})|_{\Omega_Y \times \Omega_X^a} \overset{a}{\circ} \\ D^b(k_Y, \Omega_Y) & \xrightarrow{\mu} & D^b(I(k_{\Omega_Y})) \end{array}$$

where the diagonal arrow indicates that the diagram is not commutative, but there is a natural transformation which for any $\mathcal{F} \in D^b(k_X, \Omega_X)$ is induced by

$$\mu(\mathcal{K}) \overset{a}{\circ} \mu(\mathcal{F}) \longrightarrow \mu(\mathcal{K} \circ \mathcal{F}) \longrightarrow \mu(\Phi_{\mathcal{K}}(\mathcal{F})).$$

¹¹The assumptions on $\mathcal{K}$ might imply that the second morphism of the composition is an isomorphism, but I cannot prove it.

It is an isomorphism under suitable conditions on $\mathcal{K}$. In particular, consider the situation of Theorem 5.0.2, we get a diagram of functors of prestacks

$$\begin{array}{ccc} \chi_* D^b(k_X, *)|_{\Omega_X} & \xrightarrow{\mu} & D^b(I(k_*))|_{\Omega_X} \\ \Phi_{\mathcal{K}} \downarrow & \swarrow & \downarrow \mu \mathcal{K}_{\circ}^a \\ D^b(k_Y, *)|_{\Omega_Y} & \xrightarrow{\mu} & D^b(I(k_*))|_{\Omega_Y} \end{array}$$

that commutes on small open sets of $\chi(U \times \gamma)^{12}$. In particular the diagram commutes (up to natural isomorphism) in the stalks.

Now let us consider the complex contact transformations and place ourselves in the $\mathbb{C}^\times$ -conic situation of Theorem 5.2.1. Hence all manifolds are complex, the open sets $\mathbb{C}^\times$ -conic, and the kernel $\mathcal{K}$ has $\mathbb{R}$ -constructible cohomology sheaves. We get a diagram of functors of prestacks

$$\begin{array}{ccc} \gamma_* D^b(k_X, *)|_{\Omega_X} & \xrightarrow{\mu} & \gamma_* D^b(I(k_*))|_{\Omega_X} \\ \Phi_{\mathcal{K}} \downarrow & & \downarrow \mu \mathcal{K}_{\circ}^a \\ \gamma_* D^b(k_Y, *)|_{\Omega_Y} & \xrightarrow{\mu} & \gamma_* D^b(I(k_*))|_{\Omega_Y} \end{array}$$

that commutes up to natural isomorphism in the stalks. Hence let us look at this diagram in the stalks, restrict to microlocal perverse sheaves and immediately get that for $p \in \Omega_X$ and $q = \chi(p) \in \Omega_Y$ the diagram

$$\begin{array}{ccc} D_{\text{perv}}^b(k_X, \mathbb{C}^\times p) & \xrightarrow{\mu} & \mu Perv_{\gamma(p)} \\ \Phi_{\mathcal{K}} \downarrow & & \downarrow \mu \mathcal{K}_{\circ}^a \\ D^b(k_Y, \mathbb{C}^\times q) & \xrightarrow{\mu} & D^b(I(k_*))_{\gamma(q)} \end{array} \quad (8.3.5)$$

is well defined and commutes up to natural isomorphism.

Theorem 8.3.9. *The functor of prestacks*

$$\mu \mathcal{K}_{\circ}^a : \gamma_* D^b(I(k_*))|_{\gamma(\Omega_X)} \longrightarrow \gamma_* D^b(I(k_*))|_{\gamma(\Omega_Y)}$$

induces an equivalence of stacks

$$\mu Perv^X|_{\gamma(\Omega_X)} \xrightarrow{\sim} \mu Perv^Y|_{\gamma(\Omega_Y)}$$

such that the diagram

$$\begin{array}{ccc} D_{\text{perv}}^b(k_X, *)|_{\Omega_X} & \xrightarrow{\mu} & \mu Perv^X|_{\Omega_X} \\ \Phi_{\mathcal{K}} \downarrow & & \downarrow \mu \mathcal{K}_{\circ}^a \\ D^b(k_Y, *)|_{\Omega_Y} & \xrightarrow{\mu} & \mu Perv^Y|_{\Omega_Y} \end{array}$$

commutes up to natural isomorphism and induces in the stalks the following diagram:

$$\begin{array}{ccc} D_{\text{perv}}^b(k_X, \mathbb{C}^\times p) & \xrightarrow{\mu} & \mu Perv_{\gamma(p)}^X \\ \Phi_{\mathcal{K}}^{\text{perv}} \downarrow & & \downarrow \mu \mathcal{K}_{\circ}^a \\ D_{\text{perv}}^b(k_Y, \mathbb{C}^\times q) & \xrightarrow{\mu} & \mu Perv_{\gamma(q)}^Y. \end{array}$$

¹²To apply Proposition 8.3.8, we actually have to consider an $\mathbb{R}^+$ -homogenous variant of Theorem 5.0.2 where we replace the open sets by $\mathbb{R}^+$ -conic open sets. The theorem still holds since there is actually no difference between $D^b(k_X, S)$ and $D^b(k_X, \mathbb{R}^+ S)$ by the $\mathbb{R}^+$ -conicity of the micro-support. Hence the diagram should be considered on the sphere bundle rather than on T^*X .

Proof. Consider the diagram (8.3.5). Let $\mathcal{F} \in D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$. Recall that for a sufficiently small relatively compact open neighborhood V of $\pi(p)$ the object $\mathcal{K} \circ \mathcal{F}_{\overline{V}}$ is a well-defined object of $D_{\text{perv}}^b(k_Y, \mathbb{C}^\times q)$ and isomorphic to $\mathcal{K} \circ \mathcal{F}$ in $D^b(k_Y, \mathbb{C}^\times q)$. Since $D_{\text{perv}}^b(k_Y, \mathbb{C}^\times q)$ is a full subcategory of $D^b(k_Y, \mathbb{C}^\times q)$, we get that $\mu\mathcal{K} \overset{a}{\circ} \mu\mathcal{F}$ is an object of $\mu\text{Perv}_{\gamma(q)}^Y$. Hence we get the induced functor of stacks

$$\mu\text{Perv}^X|_{\gamma(\Omega_X)} \longrightarrow \mu\text{Perv}^Y|_{\gamma(\Omega_Y)}.$$

In the same way we verify that $\mu\mathcal{K}^* \overset{a}{\circ}$ is a well defined functor of stacks in the opposite direction and it is easily verified that $\mu\mathcal{K}^* \overset{a}{\circ}$ is a quasi-inverse¹³. $\square$

Now suppose that we have contact transformations

$$\chi_1 : \Omega_X \xrightarrow{\sim} \Omega_Y \quad \chi_2 : \Omega_Y \xrightarrow{\sim} \Omega_Z$$

and kernels $\mathcal{K}_1 \in D_{\mathbb{R}\text{-c}}^b(I(k_{Y \times X}))$, $\mathcal{K}_2 \in D_{\mathbb{R}\text{-c}}^b(I(k_{Z \times Y}))$ such that Theorem 5.0.2 is valid. Then we know that $\Phi_{\mu\mathcal{K}_1}^a = \mu\mathcal{K}_1 \overset{a}{\circ}$ and $\Phi_{\mu\mathcal{K}_2}^a = \mu\mathcal{K}_2 \overset{a}{\circ}$ are well-defined on microlocally perverse sheaves. However note that $\mathcal{K}_2 \circ \mathcal{K}_1$ is not $\mathbb{R}$ -constructible and we do not know if

$$\mu\mathcal{K}_2 \overset{a}{\circ} \mu\mathcal{K}_1 \longrightarrow \mu(\mathcal{K}_2 \circ \mathcal{K}_1)$$

is an isomorphism in $D^b(I(k_{T^*Z \times T^*X}))$. Nevertheless we get

Proposition 8.3.10. *The functor*

$$\mu(\mathcal{K}_2 \circ \mathcal{K}_1) \overset{a}{\circ} : \mu\text{Perv}^X \longrightarrow \mu\text{Perv}^Z$$

is well-defined and naturally isomorphic to $\Phi_{\mu\mathcal{K}_1}^a \circ \Phi_{\mu\mathcal{K}_2}^a$.

Proof. The strategy is similar to the proof of Theorem 8.3.9. First we can show that $\mu(\mathcal{K}_2 \circ \mathcal{K}_1) \overset{a}{\circ} : \mu\text{Perv}^X \longrightarrow \mu\text{Perv}^Z$ is well-defined by looking in the stalks and using the fact that $D_{\text{perv}}^b(Y, \mathbb{C}^\times q)$ is a full subcategory of $D^b(k_Y, \mathbb{C}^\times q)$. The same argument shows the isomorphism. $\square$

Remark 8.3.11. Proposition 8.3.10 is an important step towards the definition of the stack of microlocal perverse sheaves on a complex contact manifold. Such a manifold is locally isomorphic to an open subset $\Omega_X \subset P^*X$ and the transition maps are contact transformations. Using the proposition we can locally associate an equivalence of stacks. But the choice of the kernel $\mathcal{K}$ is neither unique nor canonical.

9 Microlocal Riemann-Hilbert correspondence

The classical Riemann-Hilbert correspondence states that on a complex variety X the solution functor $\text{RHom}_{\mathcal{D}_X}(\cdot, \mathcal{O}_X)$ defines an equivalence between the stack of regular holonomic $\mathcal{D}_X$ -modules and the stack of perverse sheaves. A quasi-inverse of this functor has been constructed explicitly by Kashiwara (cf. [K2]).

We denote by $\text{HolReg}(\mathcal{D}_X)$ the abelian category of regular holonomic $\mathcal{D}_X$ -modules.

Let $\mathcal{M}$ be a holonomic $\mathcal{D}$ -module. We set

$$\text{Sol}(\mathcal{M}) = \text{RHom}_{\mathcal{D}_X}(\mathcal{M}, \mathcal{O}_X).$$

¹³Note that the difficulties encountered in Theorem 5.1.2 to prove the equivalence which were related to the fact that $D_{Rc}^b(k_X, *)$ is not a full subprestack of $D^b(k_X, *)$ do not appear here since all the morphisms are well-defined in $D^b(I(k_*))$ and μPerv is a full substack of $\gamma_* D^b(I(k_*))$.

Let $\mathcal{F}$ be an object of $D_{\mathbb{R}\text{-c}}^b(k_X)$ and set

$$\mathrm{RH}(\mathcal{F}) = T\mathcal{H}om(\mathcal{F}, \mathcal{O}_X).$$

Then the Riemann-Hilbert correspondance can be stated as follows.

Theorem 9.0.12. *The functors RH and Sol define quasi-inverse equivalences of abelian stacks*

$$\mathrm{Perv}(k_X) \xrightleftharpoons[\mathrm{Sol}]{\mathrm{RH}} \mathcal{H}ol\mathcal{R}eg(\mathcal{D}_X).$$

The microlocal Riemann-Hilbert correspondance should therefore establish an equivalence between the stack of microlocal perverse sheaves and the stack of regular holonomic $\mathcal{E}_X$ -modules.

9.1 The ind-objects $\mu\mathcal{O}_X$ and $\mu\mathcal{O}_X^t$

The microlocalization of the ring of holomorphic functions defines an object of $I(\mathbb{C}_X)$. More precisely, we have

Proposition 9.1.1. *The ind-sheaf*

$$\mu\mathcal{O}_X|_{\dot{T}^*X}$$

is concentrated in degree d_X .

Let $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. We will study the microlocal solution complex

$$\mu hom(\mathcal{F}, \mathcal{O}_X) \simeq R\mathcal{H}om(\pi^{-1}\mathcal{F}, \mu\mathcal{O}_X)$$

The stalks of this complex have been studied in [KS1] and we will show that some results can be extended to open neighborhoods.

The “ring” $\mathcal{O}_X^t \in D^b(I(\mathbb{C}_X))$ of temperate holomorphic functions has been defined in [KS2]. It is defined from the ring $\mathcal{D}b_X^t \in D^b(I(\mathbb{C}_X))$ as

$$\mathcal{O}_X^t = R\mathcal{H}om_{\beta(\mathcal{D}_X)}(\beta(\mathcal{O}_X), \mathcal{D}b_X^t).$$

We will not recall the construction of $\mathcal{D}b_X^t$ here. Recall that $\mathcal{O}_X^t$ is only defined in the derived category. It is not concentrated in a single degree.

The link with Kashiwara’s functor $T\mathcal{H}om$ is given by the formula

$$R\mathcal{H}om(\mathcal{F}, \mathcal{O}_X^t) = T\mathcal{H}om(\mathcal{F}, \mathcal{O}_X) \tag{9.1.1}$$

where $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$.

By microlization we get an object $\mu\mathcal{O}_X^t \in D^b(I(\mathbb{C}_{T^*X}))$. It is not known if this object is concentrated in a single degree or not.

In [KS2], the full functoriality of $\mathcal{O}^t$ is established. We will only need the following result:

Proposition 9.1.2. *Let $f : X \rightarrow Y$ be a smooth map between complex varieties. Then there is a canonical isomorphism in $D^b(I(\beta(\mathcal{D}_X)))$:*

$$R\mathcal{H}om_{\beta\mathcal{D}_X}(\beta\mathcal{D}_{X \rightarrow Y}, \mathcal{O}_X^t) \simeq f^{-1}\mathcal{O}_Y^t.$$

First we will prove that the microlocalization of the formula (9.1.1) holds, i.e.

Proposition 9.1.3. *Let $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$. Then*

$$R\mathcal{H}om(\mu\mathcal{F}, \mu\mathcal{O}_X^t) = t\mu hom(\mathcal{F}, \mathcal{O}_X).$$

Proof. The proof is similar to the proof of the formula

$$\mathrm{R}\mathcal{H}om(\mu\mathcal{F}, \mu\mathcal{G}) \simeq \mu hom(\mathcal{F}, \mathcal{G})$$

from [K5] where $\mathcal{F}, \mathcal{G} \in \mathrm{D}^b(k_X)$.

The normal deformation of the diagonal in $X \times X$ can be visualized by the following diagram

$$\begin{array}{ccccc} TX & \xrightarrow{\sim} & T_{\Delta_X}(X \times X) & \xrightarrow{s} & \widetilde{X \times X} & \xleftarrow{j} & \Omega \\ & & \downarrow \tau_X & & \downarrow p & \nearrow \tilde{p} & \\ X & \xrightarrow{\Delta_X} & X \times X & & X \times X & & \\ & \searrow id_X & \downarrow p_1 & \downarrow q_1 & \downarrow q_2 & \downarrow p_2 & \\ & & X & & X & & \end{array}$$

Note that $\tilde{p}, p_1$ and p_2 are smooth but p is not. Also, the square is not cartesian. We use the same notations as in Section 5 when we deform the diagonal in $T^*X \times T^*X$. We will need the following lemma

Lemma 9.1.4. *Let $\mathcal{F} \in \mathrm{D}_{\mathbb{R}\text{-}c}^b(\mathbb{C}_{T^*X})$ and $\mathcal{G} \in \mathrm{D}^b(I(\mathbb{C}_{T^*X}))$. Then*

$$\mathrm{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{G}) \simeq \omega_X^{-1} \left(s^{-1} \mathrm{R}\mathcal{H}om((p_2^{-1}\mathcal{F})_\Omega, p^{-1}q_1^!\mathcal{G}) \right)^\wedge.$$

Proof. Set

$$\begin{aligned} P &= \{(x; v) \in TT^*X \mid \langle v, \omega(x) \rangle \leq 0\}, \\ P' &= \{(x; v) \in TT^*X \mid \langle v, \omega(x) \rangle \geq 0\}. \end{aligned}$$

We have

$$\begin{aligned} \mathrm{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{G}) &\simeq \mathrm{R}\mathcal{H}om(\mathcal{F}, \mathrm{R}q_{1!!}(q_2^{-1}\mathcal{G} \otimes \mathrm{R}p_{!!}(\mathbb{C}_{\overline{\Omega}}) \otimes \beta(\mathbb{C}_P \otimes \mathrm{R}s_*\omega_{TT^*X|T^*X})) \\ &\simeq \mathrm{R}\mathcal{H}om(\mathcal{F}, \mathrm{R}p_{1!!}(p_2^{-1}\mathcal{G} \otimes \mathbb{C}_{\overline{\Omega}} \otimes \beta(\mathbb{C}_P \otimes \mathrm{R}s_*\omega_{TT^*X|T^*X}))) \\ &\simeq \mathrm{R}\mathcal{H}om(\mathcal{F}, \mathrm{R}p_{2!!}(p_1^{-1}\mathcal{G} \otimes \mathbb{C}_{\overline{\Omega}} \otimes \beta(\mathbb{C}_{P'} \otimes \mathrm{R}s_*\omega_{TT^*X|T^*X}))) \\ &\simeq \mathrm{R}p_{2!} \mathrm{R}\mathcal{H}om(p_2^{-1}\mathcal{F}, p_1^{-1}\mathcal{G} \otimes \mathbb{C}_{\overline{\Omega}} \otimes \beta(\mathbb{C}_{P'} \otimes \mathrm{R}s_*\omega_{TT^*X|T^*X})) \\ &\simeq \mathrm{R}p_{2!} \left(\mathrm{R}\mathcal{H}om(p_2^{-1}\mathcal{F}, p_1^{-1}\mathcal{G} \otimes \mathbb{C}_{\overline{\Omega}}) \otimes \mathbb{C}_{P'} \otimes \mathrm{R}s_*\omega_{TT^*X|T^*X} \right) \end{aligned}$$

Now note that locally on $T^*\widetilde{X} \times T^*X$ the set Ω is convex and

$$\mathrm{SS}_0(p_1^{-1}\mathcal{G}) \cap T_\Omega^*(T^*\widetilde{X} \times T^*X) \subset T_{T^*\widetilde{X} \times T^*X}^*(T^*\widetilde{X} \times T^*X).$$

Hence

$$\mathbb{C}_{\overline{\Omega}} \otimes p_1^{-1}\mathcal{G} \simeq \mathrm{R}\mathcal{H}om(\mathbb{C}_\Omega, \mathbb{C}_X) \otimes p_1^{-1}\mathcal{G} \simeq \mathrm{R}\mathcal{H}om(\mathbb{C}_\Omega, p_1^{-1}\mathcal{G}).$$

Moreover note that

$$\omega_{TT^*X|T^*X} \simeq \tau_{T^*X}^{-1} \omega_{\Delta|T^*X \times T^*X}^{\otimes -1} \simeq s^{-1} \tilde{p}^{-1} q_1^! \mathbb{C}_{T^*X}.$$

Hence

$$\begin{aligned} \mathrm{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{G}) &\simeq \mathrm{R}p_{2!} \left(\mathrm{R}\mathcal{H}om(p_2^{-1}\mathcal{F} \otimes \mathbb{C}_\Omega, p_1^{-1}\mathcal{G}) \otimes \mathbb{C}_{P'} \otimes \mathrm{R}s_*\omega_{TT^*X|T^*X} \right) \\ &\simeq \mathrm{R}p_{2!} \mathrm{R}s_! \left(s^{-1} \mathrm{R}\mathcal{H}om(p_2^{-1}\mathcal{F} \otimes \mathbb{C}_\Omega, p^{-1}q_1^!\mathcal{G}) \otimes \mathbb{C}_{P'} \right) \\ &\simeq \mathrm{R}\gamma_! \left(s^{-1} \mathrm{R}\mathcal{H}om(p_2^{-1}\mathcal{F} \otimes \mathbb{C}_\Omega, p^{-1}q_1^!\mathcal{G}) \otimes \mathbb{C}_{P'} \right) \\ &\simeq \omega_X^{-1} \left(s^{-1} \mathrm{R}\mathcal{H}om((p_2^{-1}\mathcal{F})_\Omega, p^{-1}q_1^!\mathcal{G}) \right)^\wedge. \end{aligned}$$

This shows the lemma. $\square$

Now consider the situation of the lemma but take $\mathcal{F} \in D^b(\mathbb{C}_X)$. Then a standard calculation shows that

$$R\mathcal{H}om(\pi^{-1}\mathcal{F}, \mu\mathcal{G}) = \left(s^{-1} R\mathcal{H}om((p_2^{-1}\mathcal{F})_\Omega, p^{-1}q_1^!\mathcal{G}) \right)^\wedge.$$

When we apply this result to $\mathcal{O}_X^t$ and $\mathcal{F} \in D_{\mathbb{R}\text{-c}}^b(k_X)$ we get

$$\begin{aligned} R\mathcal{H}om(\pi^{-1}\mathcal{F}, \mu\mathcal{O}^t) &\simeq \left(s^{-1} R\mathcal{H}om((p_2^{-1}\mathcal{F})_{\Omega_X}, p^{-1}q_1^!\mathcal{O}^t) \right)^\wedge \\ &\simeq \left(s^{-1} R\mathcal{H}om_{\mathcal{D}_{\widetilde{X \times X}}}(\mathcal{D}_{\widetilde{X \times X} \xrightarrow{p_1} X}, T\mathcal{H}om((p_2^{-1}\mathcal{F})_{\Omega_X}, \mathcal{O}_{\widetilde{X \times X}}) \otimes s^{-1}p^{-1}q_1^!\mathbb{C}_X) \right)^\wedge \\ &\simeq \left(s^{-1} \mathcal{D}_{X \xleftarrow{p_1} \widetilde{X \times X}} \otimes_{\mathcal{D}_{\widetilde{X \times X}}} T\mathcal{H}om((p_2^{-1}\mathcal{F})_\Omega, \mathcal{O}_{\widetilde{X \times X}})[1] \right)^\wedge \\ &\simeq t\mu hom(\mathcal{F}, \mathcal{O}_X). \end{aligned}$$

□

Following [KS4], we set for a locally free $\mathcal{O}_X$ -module $\mathcal{L}$:

$$\mathcal{L}^t = \mathcal{O}^t \underset{\beta\mathcal{O}_X}{\overset{L}{\otimes}} \beta\mathcal{L}.$$

Lemma 9.1.5. *Let $\mathcal{L}$ a locally free $\mathcal{O}_X$ -module of finite rank. Then there are natural isomorphisms*

$$\begin{aligned} \mu\mathcal{L} &\longrightarrow \mu\mathcal{O}_X \underset{\pi^{-1}\beta\mathcal{O}_X}{\overset{L}{\otimes}} \pi^{-1}\beta\mathcal{L}, \\ \mu\mathcal{L}^t &\longrightarrow \mu\mathcal{O}_X^t \underset{\pi^{-1}\beta\mathcal{O}_X}{\overset{L}{\otimes}} \pi^{-1}\beta\mathcal{L}. \end{aligned}$$

Proof. Let us show the second isomorphism, the proof of the first being similar since $\mathcal{L} \simeq \beta\mathcal{L} \underset{\beta\mathcal{O}_X}{\otimes} \mathcal{O}_X$. By definition

$$\mu(\mathcal{O}_X^t \underset{\beta\mathcal{O}_X}{\otimes} \beta\mathcal{L}) \simeq Rq_{1!!} \left(K_X \otimes q_2^{-1} \pi^{-1} (\mathcal{O}_X^t \underset{\beta\mathcal{O}_X}{\otimes} \beta\mathcal{L}) \right).$$

But since $K_X \simeq K_X \otimes \beta\mathbb{C}_\Delta$ and

$$\beta\mathbb{C}_\Delta \otimes (q_2^{-1} \pi^{-1} \mathcal{O}_X^t \underset{\beta q_2^{-1} \pi^{-1} \mathcal{O}_X}{\overset{L}{\otimes}} \beta q_2^{-1} \pi^{-1} \mathcal{L}) \simeq ((\beta\mathbb{C}_\Delta \otimes q_2^{-1} \pi^{-1} \mathcal{O}_X^t) \underset{\beta q_1^{-1} \pi^{-1} \mathcal{O}_X}{\overset{L}{\otimes}} \beta q_1^{-1} \pi^{-1} \mathcal{L})$$

we get the second isomorphism. □

9.2 $\mathcal{E}_X$ -modules

The ring $\mathcal{E}_X$ of microdifferential operators on T^*X has been defined in [SKK]. For a short introduction to the theory of $\mathcal{E}_X$ -modules we refer to [K3], for a more detailed study see [Sch]. The ring $\mathcal{E}_X$ has many “good” properties, for instance it is coherent and Noetherian. For our purpose it is convenient to consider microdifferential operators outside the zero-section, hence when we write $\mathcal{E}_X$, we consider $\mathcal{E}_X|_{T^*X}$.

We will consider two variants of this ring. In the sequel we will identify T^*X with $T_{\Delta_X}^*(X \times X)$ by the map

$$\delta^a : T^*X \xrightarrow{\sim} T_{\Delta_X}^*(X \times X) \hookrightarrow T^*X \times T^*X \quad ; \quad (x; \xi) \mapsto ((x; \xi), (x; -\xi)).$$

The functor δ^{a-1} is often omitted for complexes with support on T^*X . First one defines the ring $\mathcal{E}_X^{\mathbb{R}}$ on T^*X as

$$\mathcal{E}_X^{\mathbb{R}} \simeq H^{dx} \mu hom(\mathbb{C}_{\Delta_X}, \mathcal{O}_{X \times X}^{(dx,0)}) \simeq H^{dx} R\mathcal{H}om(\pi^{-1}\mathbb{C}_{\Delta_X}, \mu\mathcal{O}_X^{(dx,0)}).$$

In [An2] Andronikof introduced the ring $\mathcal{E}_X^{\mathbb{R},f}$ on T^*X of tempered microdifferential operators as

$$\mathcal{E}_X^{\mathbb{R},f} = H^{dx} (t\mu hom(\mathbb{C}_{\Delta}, \mathcal{O}_{X \times X}) \bigotimes_{\mathcal{O}_{X \times X}}^L \mathcal{O}_{X \times X}^{(dx,0)})$$

where we omit the functor $(\pi \times \pi)^{-1}$ in the notation.

Proposition 9.2.1. *We have*

$$\mathcal{E}_X^{\mathbb{R},f} \simeq H^{dx} R\mathcal{H}om(\pi^{-1}\mathbb{C}_{\Delta_X}, \mu\mathcal{O}_{X \times X}^{t,(dx,0)}).$$

Proof. Follows from Proposition 9.1.3 and Lemma 9.1.5. □

Recall some basic properties:

- (i) We have $\mathcal{E}_X \simeq \gamma^{-1} R\gamma_* \mathcal{E}_X^{\mathbb{R}} \simeq \gamma^{-1} R\gamma_* \mathcal{E}_X^{\mathbb{R},f}$. In particular $\gamma^{-1} R\gamma_* \mathcal{E}_X \simeq \mathcal{E}_X$. Moreover $R^i \gamma_* \mathcal{E}_X \simeq 0$ for $i \neq 0$.
- (ii) The rings $\mathcal{E}_X^{\mathbb{R}}$ and $\mathcal{E}_X^{\mathbb{R},f}$ are faithfully flat over $\mathcal{E}_X$.
- (iii) The ring $\mathcal{E}_X$ (and therefore $\mathcal{E}_X^{\mathbb{R}}, \mathcal{E}_X^{\mathbb{R},f}$) is a $\pi^{-1}\mathcal{D}_X$ -modules.

A priori, $\mathcal{E}_X$ -modules are defined on the cotangent space T^*X . But by (i) coherent $\mathcal{E}_X$ -modules (hence in particular regular holonomic $\mathcal{E}_X$ -modules) are conic objects, hence it is often convenient to work on the projective bundle P^*X or on $\mathbb{C}^\times$ -conic sets.

Let $\mathcal{M}$ be an $\mathcal{E}_X$ -module. Then its support $\text{supp}(\mathcal{M})$ is called its characteristic variety. If $\mathcal{M}$ is a $\mathcal{D}_X$ -module, then the characteristic variety of $\mathcal{E}_X \otimes_{\pi^{-1}\mathcal{D}_X} \pi^{-1}\mathcal{M}$ coincides with the characteristic variety of $\mathcal{M}$ as a $\mathcal{D}_X$ -module. The main result about the characteristic variety of $\mathcal{E}_X$ -modules is:

Proposition 9.2.2. *Let $\mathcal{M}$ be a coherent $\mathcal{E}_X$ -module. Then its characteristic variety is a closed analytic, involutive, $\mathbb{C}^\times$ -conic subset of T^*X .*

Definition 9.2.3. *Let $\mathcal{M}$ be a coherent $\mathcal{E}_X$ -module.*

One says that $\mathcal{M}$ is holonomic if its characteristic variety is Lagrangian.

Remark 9.2.4. Hence a holonomic $\mathcal{D}_X$ -module defines a holonomic $\mathcal{E}_X$ -module.

Regular holonomic $\mathcal{E}_X$ -modules (or “holonomic systems with regular singularities”) have been studied in [KK]. A $\mathcal{D}_X$ -module is regular holonomic if and only if its associated $\mathcal{E}_X$ -module is regular holonomic. We do not recall the definition here since regular holonomic systems can be characterized by the Riemann-Hilbert correspondence and refer to loc. cit.

Theorem 9.2.5. *Let $\mathcal{M}$ be a regular holonomic $\mathcal{E}_X$ -module such that its characteristic variety is in generic position at a point $p \in T^*X$. Then there exists a regular holonomic $\mathcal{D}_X$ -modules $\tilde{\mathcal{M}}$ such that*

$$\mathcal{M} \simeq \mathcal{E}_X \otimes_{\pi^{-1}\mathcal{D}_X} \tilde{\mathcal{M}}.$$

Regular holonomic $\mathcal{E}_X$ -modules form a stack of abelian categories. This stack is invariant by quantized contact transformations. Hence modulo a contact transformation a regular holonomic $\mathcal{E}_X$ -module is locally isomorphic to

$$\mathcal{E}_X \otimes_{\pi^{-1}\mathcal{D}_X} \mathrm{RH}(\mathcal{F})$$

where $\mathcal{F}$ is a perverse sheaf. For our purpose this can be taken as a definition of a regular holonomic system.

Finally recall the following theorem.

Proposition 9.2.6. ([An2], Théorème 4.2.6 and Proposition 5.6.1) *Let $\mathcal{F} \in \mathrm{D}_{\mathbb{C}-c}^b(k_X)$. Then*

$$t\mu\mathrm{hom}(\mathcal{F}, \mathcal{O}_X) \simeq \mathcal{E}_X^{\mathbb{R},f} \otimes_{\pi^{-1}\mathcal{D}_X}^L \pi^{-1} \mathrm{T}\mathcal{H}\mathrm{om}(\mathcal{F}, \mathcal{O}_X).$$

If $\mathcal{F}$ is perverse then $t\mu\mathrm{hom}(\mathcal{F}, \mathcal{O}_X)$ is concentrated in degree 0 and

$$H^0 t\mu\mathrm{hom}(\mathcal{F}, \mathcal{O}_X) \simeq \mathcal{E}_X^{\mathbb{R},f} \otimes_{\pi^{-1}\mathcal{D}_X} \pi^{-1} \mathrm{T}\mathcal{H}\mathrm{om}(\mathcal{F}, \mathcal{O}_X).$$

9.3 The microdifferential structure of $\mu\mathcal{O}_X$ and $\mu\mathcal{O}_X^t$

Recall that an object $\mathcal{A} \in \mathrm{I}(k_X)$ is called a k_X -algebra if there exist morphisms

$$k_X \rightarrow \mathcal{A} \quad \mathcal{A} \otimes \mathcal{A} \longrightarrow \mathcal{A}$$

that satisfy the usual conditions of unit and associativity (for more details, see for instance [KS2], Section 5.4). For example, If $\mathcal{A}$ is a classical k_X -algebra in $\mathrm{Mod}(k_X)$ then $\beta\mathcal{A}$ is a k_X -algebra in $\mathrm{I}(k_X)$.

Let $\mathcal{A} \in \mathrm{I}(k_X)$ be a k_X -algebra. A left $\mathcal{A}$ -module in $\mathrm{I}(k_X)$ is given by an object $\mathcal{M} \in \mathrm{I}(k_X)$ and a structure morphism

$$\mathcal{A} \otimes \mathcal{M} \longrightarrow \mathcal{M}$$

that satisfies the usual compatibility conditions with the structure morphisms of $\mathcal{A}$ (for more details, see loc. cit.).

Similarly, one defines the notion of a k_X -algebra $\mathcal{A}$ in $\mathrm{D}^b(\mathrm{I}(k_X))$ and the notion of a left $\mathcal{A}$ -module in $\mathrm{D}^b(\mathrm{I}(k_X))$. Note that if a k_X -algebra $\mathcal{A}$ in $\mathrm{D}^b(\mathrm{I}(k_X))$ is concentrated in a single degree, then it defines a k_X -algebra in $\mathrm{I}(k_X)$. However, even if $\mathcal{A}$ is concentrated in a single degree, an $\mathcal{A}$ -module in $\mathrm{D}^b(\mathrm{I}(k_X))$ is in general not well-defined in the derived category of $\mathcal{A}$ -modules in $\mathrm{I}(k_X)$. In order to avoid confusion, one often calls an $\mathcal{A}$ -module in $\mathrm{D}^b(\mathrm{I}(k_X))$ a formal $\mathcal{A}$ -module.

In this section we will show that $\mu\mathcal{O}_X$ is a $\beta\mathcal{E}_X^{\mathbb{R}}$ -module in $\mathrm{I}(k_X)$. The same strategy will show that $\mu\mathcal{O}_X^t$ is a formal $\beta\mathcal{E}_X^{\mathbb{R},f}$ -module in the derived category $\mathrm{D}^b(\mathrm{I}(k_X))$. In particular its cohomology ind-sheaves are $\beta\mathcal{E}_X^{\mathbb{R},f}$ -modules.

Let $f : X \rightarrow Y$ be a morphism of complex manifolds.

Lemma 9.3.1. *There is a natural morphism in $\mathrm{D}^b(\mathrm{I}(\mathbb{C}_X))$*

$$\varphi : \mathrm{R}f_{!!}\Omega_X[d_X] \longrightarrow \Omega_Y[d_Y]$$

such that $\alpha(\varphi)$ is the classical integration morphism.

Proof. Consider the chain of natural isomorphisms:

$$\begin{aligned} \mathrm{Hom}_{\mathrm{D}^b(\mathrm{I}(\mathbb{C}_Y))}(\mathrm{R}f_{!!}\Omega_X[d_X], \Omega_Y[d_Y]) &\simeq \mathrm{Hom}_{\mathrm{D}^b(\mathrm{I}(\mathbb{C}_X))}(\Omega_X[d_X], f^!\Omega_Y[d_Y]) \\ &\simeq \mathrm{Hom}_{\mathrm{D}^b(\mathbb{C}_X)}(\Omega_X[d_X], f^!\Omega_Y[d_Y]) \simeq \mathrm{Hom}_{\mathrm{D}^b(\mathbb{C}_Y)}(\mathrm{R}f_!\Omega_X[d_X], \Omega_Y[d_Y]) \end{aligned}$$

The inverse image of the classical integration morphism gives φ . □

There is a natural tempered version of the integration morphism:

Proposition 9.3.2. *There is a natural morphism in $D^b(I(\mathbb{C}_X))$*

$$Rf_{!!}\Omega_X^t[d_X] \longrightarrow \Omega_Y^t[d_Y].$$

Proof. This is a simple version (not respecting $\mathcal{D}$ -module structures) of the morphism established in [KS2]:

$$Rf_{!!}(\Omega_X^t \bigotimes_{\beta\mathcal{D}_X}^L \beta\mathcal{D}_{X \rightarrow Y})[d_X] \longrightarrow \Omega_Y^t[d_Y].$$

□

Now consider complex manifolds X, Y, Z of complex dimensions d_X, d_Y, d_Z and the diagram

$$\begin{array}{ccc} & X \times Y \times Z & \\ q_{12} \swarrow & \downarrow q_{13} & \searrow q_{23} \\ X \times Y & X \times Z & Y \times Z. \end{array}$$

Lemma 9.3.3. *The integration morphisms induce natural morphisms*

$$\begin{aligned} \mu\mathcal{O}_{X \times Y}^{(d_X, 0)} \circ^a \mu\mathcal{O}_{Y \times Z}^{(d_Y, 0)} &\longrightarrow \mu\mathcal{O}_{X \times Z}^{(d_X, 0)}[-d_Y] \\ \mu\mathcal{O}_{X \times Y}^{t, (d_X, 0)} \circ^a \mu\mathcal{O}_{Y \times Z}^{t, (d_Y, 0)} &\longrightarrow \mu\mathcal{O}_{X \times Z}^{t, (d_X, 0)}[-d_Y] \end{aligned}$$

Proof. The two constructions being similar (and just an ind-variant of the construction used in Lemma 11.4.3. of [KS1]) we will only show how to define the second morphism. First let us consturct the natural morphism

$$\mathcal{O}_{X \times Y}^{t, (d_X, 0)} \circ \mathcal{O}_{Y \times Z}^{t, (d_Y, 0)} \longrightarrow \mathcal{O}_{X \times Z}^{t, (d_X, 0)}[-\dim_{\mathbb{C}} Y].$$

It can be obtained as follows

$$\begin{aligned} \mathcal{O}_{X \times Y}^{t, (d_X, 0)} \circ \mathcal{O}_{Y \times Z}^{t, (d_Y, 0)} &\simeq R p_{13!!} (p_{12}^{-1} \mathcal{O}_{X \times Y}^{t, (d_X, 0)} \otimes p_{23}^{-1} \mathcal{O}_{Y \times Z}^{t, (d_Y, 0)}) \\ &\rightarrow R p_{13!!} \mathcal{O}_{X \times Y \times Z}^{t, (d_X, d_Y, 0)} \\ &\rightarrow \mathcal{O}_{X \times Z}^{t, (d_X, 0)}[-d_Y] \end{aligned}$$

where the last morphism is the integration morphism of Proposition 9.3.2. Using the microlocal composition formula we get a morphism

$$\mu\mathcal{O}_{X \times Y}^{t, (d_X, 0)} \circ^a \mu\mathcal{O}_{Y \times Z}^{t, (d_Y, 0)} \longrightarrow \mu(\mathcal{O}_{X \times Y}^{t, (d_X, 0)} \circ \mathcal{O}_{Y \times Z}^{t, (d_Y, 0)}) \longrightarrow \mu\mathcal{O}_{X \times Z}^{t, (d_X, 0)}[-d_Y].$$

Therefore we get the morphism of the proposition. □

Proposition 9.3.4. *Let $\mathcal{K}_1 \in D^b(I(\mathbb{C}_{T^*X \times T^*Y}))$ and $\mathcal{K}_2 \in D^b(I(\mathbb{C}_{T^*Y \times T^*Z}))$. There are natural morphisms*

$$\begin{aligned} R\mathcal{J}\mathcal{H}om(\mathcal{K}_1, \mu\mathcal{O}_{X \times Y}^{(d_X, 0)}) \circ^a R\mathcal{J}\mathcal{H}om(\mathcal{K}_2, \mu\mathcal{O}_{Y \times Z}^{(d_Y, 0)}) &\longrightarrow R\mathcal{J}\mathcal{H}om(\mathcal{K}_1 \circ^a \mathcal{K}_2, \mu\mathcal{O}_{X \times Z}^{(d_X, 0)})[-d_Y] \\ R\mathcal{J}\mathcal{H}om(\mathcal{K}_1, \mu\mathcal{O}_{X \times Y}^{t, (d_X, 0)}) \circ^a R\mathcal{J}\mathcal{H}om(\mathcal{K}_2, \mu\mathcal{O}_{Y \times Z}^{t, (d_Y, 0)}) &\longrightarrow R\mathcal{J}\mathcal{H}om(\mathcal{K}_1 \circ^a \mathcal{K}_2, \mu\mathcal{O}_{X \times Z}^{t, (d_X, 0)})[-d_Y] \end{aligned}$$

These morphisms satisfy the obvious associativity condition (analogous to Lemma 11.4.3 in [KS1]).

Proof. Note that there is a natural morphism

$$\mathrm{R}\mathcal{H}om(\mathcal{F}_1, \mathcal{G}_1) \overset{a}{\circ} \mathrm{R}\mathcal{H}om(\mathcal{F}_2, \mathcal{G}_2) \longrightarrow \mathrm{R}\mathcal{H}om(\mathcal{F}_1 \overset{a}{\circ} \mathcal{F}_2, \mathcal{G}_1 \overset{a}{\circ} \mathcal{G}_2).$$

Combining this morphism with the morphisms of Lemma 9.3.3, we get the desired arrows. The associativity condition is tedious to write down. It is a straightforward consequence of the corresponding associativity conditions of the morphisms involved in the construction. $\square$

Remark 9.3.5. Consider $\mathcal{K}_1 \in \mathrm{D}^b(\mathbb{C}_{X \times Y})$ and $\mathcal{K}_2 \in \mathrm{D}^b(\mathbb{C}_{Y \times Z})$. Applying the functor α to the first morphism of Proposition 9.3.4, we get a morphism

$$\mathrm{R}p_{13!} \left(p_{12}^{a-1} \mu hom(\mathcal{K}_1, \mathcal{O}_{X \times Y}^{(d_X, 0)}) \otimes p_{23}^{-1} \mu hom(\mathcal{K}_2, \mathcal{O}_{Y \times Z}^{(d_Y, 0)}) \right) \longrightarrow \mu hom(\mathcal{K}_1 \circ \mathcal{K}_2, \mathcal{O}_{X \times Z}^{(d_X, 0)})[-d_Y].$$

This is precisely the morphism of [KS1], Lemma 11.4.3.

In the situation of Proposition 9.3.4 consider $Z = \{pt\}$ and $\mathcal{K}_2 = \mu\mathcal{O}_Y$ (resp. $\mathcal{K}_2 = \mu\mathcal{O}_Y^t$). Applying α we get natural morphisms

$$\mathrm{R}p_{1!} \left(\mathrm{R}\mathcal{H}om(\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(d_X, 0)}) \otimes p_2^{a-1} \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_Y, \mu\Omega_Y) \right) \longrightarrow \mathrm{R}\mathcal{H}om(\mathcal{K} \overset{a}{\circ} \mu\mathcal{O}_Y, \mu\Omega_X)[-d_Y],$$

$$\mathrm{R}p_{1!} \left(\mathrm{R}\mathcal{H}om(\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{t, (d_X, 0)}) \otimes p_2^{a-1} \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_Y^t, \mu\Omega_Y^t) \right) \longrightarrow \mathrm{R}\mathcal{H}om(\mathcal{K} \overset{a}{\circ} \mu\mathcal{O}_Y^t, \mu\Omega_X^t)[-d_Y].$$

Hence we get

$$\mathrm{R}p_{1!} \left(\mathrm{R}\mathcal{H}om(\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(0, d_Y)}) \otimes p_2^{a-1} \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_Y, \mu\mathcal{O}_Y) \right) \longrightarrow \mathrm{R}\mathcal{H}om(\mathcal{K}[d_Y] \overset{a}{\circ} \mu\mathcal{O}_Y, \mu\mathcal{O}_X), \quad (9.3.1)$$

$$\mathrm{R}p_{1!} \left(\mathrm{R}\mathcal{H}om(\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{t, (0, d_Y)}) \otimes p_2^{a-1} \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_Y^t, \mu\mathcal{O}_Y^t) \right) \longrightarrow \mathrm{R}\mathcal{H}om(\mathcal{K}[d_Y] \overset{a}{\circ} \mu\mathcal{O}_Y^t, \mu\mathcal{O}_X^t). \quad (9.3.2)$$

Taking $X = Y$ and $\mathcal{K} = \pi^{-1}\mathbb{C}_{\Delta_X}$ we get the morphisms

$$\mathcal{E}_X^{\mathbb{R}} \otimes \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_X, \mu\mathcal{O}_X) \longrightarrow \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_X, \mu\mathcal{O}_X),$$

$$\mathcal{E}_X^{\mathbb{R}, f} \otimes \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_X^t, \mu\mathcal{O}_X^t) \longrightarrow \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_X^t, \mu\mathcal{O}_X^t).$$

Note that for any $\mathcal{F} \in \mathrm{D}^b(\mathrm{I}(\mathbb{C}_X))$ the identity of $\mathcal{F}$ defines a natural morphism $\mathbb{C}_{\Delta_X} \rightarrow \mathrm{R}\mathcal{H}om(\mathcal{F}, \mathcal{F})$. Hence we get the structure morphisms

$$\beta\mathcal{E}_X^{\mathbb{R}} \otimes \mu\mathcal{O}_X \longrightarrow \mu\mathcal{O}_X, \quad (9.3.3)$$

$$\beta\mathcal{E}_X^{\mathbb{R}, f} \otimes \mu\mathcal{O}_X^t \longrightarrow \mu\mathcal{O}_X^t. \quad (9.3.4)$$

In order to prove that the two morphisms (9.3.3) and (9.3.4) define structures of formal modules in $\mathrm{D}^b(\mathrm{I}(\mathbb{C}_X))$ one uses the associativity of the construction in Proposition (9.3.4). Therefore we get:

Proposition 9.3.6. *The object $\mu\mathcal{O}_X$ (resp. $\mu\Omega_X$) is a left (resp. right) $\beta\mathcal{E}_X^{\mathbb{R}}$ -module in $\mathrm{D}^b(\mathrm{I}(\mathbb{C}_X))$.*

The object $\mu\mathcal{O}_X^t$ (resp. $\mu\Omega_X^t$) is a left (resp. right) $\beta\mathcal{E}_X^{\mathbb{R}, f}$ -module in $\mathrm{D}^b(\mathrm{I}(\mathbb{C}_X))$.

Corollary 9.3.7. *Let $\mathcal{F} \in \mathrm{D}^b(\mathrm{I}(\mathbb{C}_{T^*X}))$. Then*

- (1) the complex $\mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{F}, \mu\mathcal{O}_X)$ (resp. $\mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{F}, \mu\mathcal{O}_X^t)$) is a left $\beta\mathcal{E}_X^{\mathbb{R}}$ -module (resp. $\beta\mathcal{E}_X^{\mathbb{R},f}$ -module) in $D^b(I(\mathbb{C}_X))$,
- (2) the complex $\mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{F}, \mu\Omega_X)$ (resp. $\mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{F}, \mu\Omega_X^t)$) is a right $\beta\mathcal{E}_X^{\mathbb{R}}$ -module (resp. $\beta\mathcal{E}_X^{\mathbb{R},f}$ -module) in $D^b(I(\mathbb{C}_X))$.

Corollary 9.3.8. (i) The object $\mu\mathcal{O}_X$ (resp. $\mu\Omega_X$) is a left (resp. right) $\beta\mathcal{E}_X^{\mathbb{R}}$ -module in $I(\mathbb{C}_{T^*X})$.

(ii) For any $\mathcal{F} \in D^b(I(\mathbb{C}_{T^*X}))$ the complex $\mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{F}, \mu\mathcal{O}_X)$ (resp. $\mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{F}, \mu\Omega_X)$) is well defined in the bounded derived category of left (resp. right) $\beta\mathcal{E}_X^{\mathbb{R}}$ -modules. In particular if $\mathcal{F} \in D^b(k_X)$ then $\mu\mathrm{hom}(\mathcal{F}, \mathcal{O}_X)$ (resp. $\mu\mathrm{hom}(\mathcal{F}, \Omega_X)$) is well defined in the bounded derived category of left (resp. right) $\mathcal{E}_X^{\mathbb{R}}$ -modules.

(iii) Let $\mathcal{F} \in D^b(\mathbb{C}_X)$. Then $\mu\mathrm{hom}(\mathcal{F}, \mathcal{O}_X)$ is well defined in the derived category of $\mathcal{E}_X^{\mathbb{R}}$ -modules.

(iv) Let $\mathcal{F} \in D^b(I(\mathbb{C}_{T^*X}))$ such that $\mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{F}, \mu\mathcal{O}_X^t)$ is concentrated in a single degree (for instance if $\mathcal{F}$ is a microlocal perverse sheaf (see Lemma 9.6.1 below)). Then the natural morphism

$$\mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{F}, \mu\mathcal{O}_X^t) \longrightarrow \mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{F}, \mu\mathcal{O}_X)$$

is well defined in the derived category of $\mathcal{E}_X$ -modules.

Proposition 9.3.9. The natural morphism

$$\begin{aligned} \mathrm{Rp}_{13!!} \left(p_{12}^{a-1} \mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{K}_1, \mu\mathcal{O}_{X \times Y}^{(d_X, 0)}) \otimes p_{23}^{-1} \mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{K}_2, \mu\mathcal{O}_{Y \times Z}^{(d_Y, 0)}) \right) \\ \longrightarrow \mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{K}_1 \overset{a}{\circ} \mathcal{K}_2, \mu\mathcal{O}_{X \times Z}^{(d_X, 0)})[-d_Y] \end{aligned}$$

factors through

$$\mathrm{Rp}_{13!!} \left(p_{12}^{a-1} \mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{K}_1, \mu\mathcal{O}_{X \times Y}^{(d_X, 0)}) \overset{\mathrm{L}}{\otimes}_{p_2^{-1}\beta\mathcal{E}_X^{\mathbb{R}}} p_{23}^{-1} \mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{K}_2, \mu\mathcal{O}_{Y \times Z}^{(d_Y, 0)}) \right).$$

Proof. This is a consequence of the associativity condition of Proposition 9.3.4. $\square$

9.4 Quantized contact transformations for $\mu\mathcal{O}_X$ and $\mu\mathcal{O}_X^t$

In this section we will adapt Section 11.4 of [KS1] to study the behaviour of $\mu\mathcal{O}_X$ and $\mu\mathcal{O}_X^t$ under complex contact transformations. We will restrict ourselves now to $\mu\mathcal{O}_X$, the study of $\mu\mathcal{O}_X^t$ being similar. We also did not include all proofs which are rather similar to Section 11.4 of loc. cit.

Our main interest is to prove that the complex

$$\mathrm{R}\mathcal{H}\mathrm{om}_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mu\mathcal{O}_X)$$

is invariant under quantized contact transformations where $\mathcal{M}$ is a coherent $\mathcal{E}_X$ -module. More precisely consider a contact transformation

$$\chi : \Omega_X \xrightarrow{\sim} \Omega_Y$$

Then locally we may find a kernel $\mathcal{K}$ such that $\Phi_{\mu\mathcal{K}}^a$ induces an equivalence of microlocal perverse sheaves and Since $\mathcal{M}$ is coherent, the natural morphism

$$\mu\mathcal{K} \overset{a}{\circ} \mathrm{R}\mathcal{H}\mathrm{om}_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mu\mathcal{O}_X) \longrightarrow \mathrm{R}\mathcal{H}\mathrm{om}_{\beta(\chi_*\mathcal{E}_X)}(\beta(\chi_*\mathcal{M}), \mu\mathcal{K} \overset{a}{\circ} \mu\mathcal{O}_X)$$

is an isomorphism. Hence we are reduced to construct an isomorphism

$$\mu\mathcal{K} \circ^a \mu\mathcal{O}_Y \longrightarrow \mu\mathcal{O}_X$$

that is compatible with the action of $\beta\mathcal{E}_X$ (resp. $\beta\mathcal{E}_Y$) on $\mu\mathcal{O}_X$ (resp. $\mu\mathcal{O}_Y$). Such a morphism has been constructed in $D^b(k_X, p)$ in Section 11.4 of [KS1].

Recall the morphism (9.3.1)

$$Rp_{1!}\left(\mathrm{R}\mathcal{H}om(\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(0, d_Y)}) \otimes p_2^{a-1} \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_Y, \mu\mathcal{O}_Y)\right) \longrightarrow \mathrm{R}\mathcal{H}om(\mathcal{K}[d_Y] \circ^a \mu\mathcal{O}_Y, \mu\mathcal{O}_X).$$

Note that if $\mathcal{K} \simeq \mu K$ and $\mathrm{supp} K \rightarrow X$ is proper, then we get the commutative diagram

$$\begin{array}{ccc} Rp_{1!}\left(\mu hom(K, \mathcal{O}_{X \times Y}^{(0, d_Y)}) \otimes p_2^{a-1} \mu hom(\mathcal{O}_Y, \mathcal{O}_Y)\right) & \longrightarrow & \mathrm{R}\mathcal{H}om(\mu K[d_Y] \circ^a \mu\mathcal{O}_Y, \mu\mathcal{O}_X) \\ & \searrow & \uparrow \\ & & \mu hom(K[d_Y] \circ \mathcal{O}_Y, \mathcal{O}_X) \end{array}$$

where the diagonal morphism is the classical morphism from [KS1].

Now suppose that $\mathrm{R}\mathcal{H}om(\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(0, d_Y)})$ is concentrated in positive degrees and that

$$\mathrm{supp} \mathrm{R}\mathcal{H}om(\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(0, d_Y)}) \rightarrow X$$

is proper. Then by taking the 0-cohomology we get a morphism

$$p_{1*}\left(H^0 \mathrm{R}\mathcal{H}om(\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(0, d_Y)}) \otimes p_2^{a-1} H^0 \mathrm{R}\mathcal{H}om(\mu\mathcal{O}_Y, \mu\mathcal{O}_Y)\right) \longrightarrow H^0 \mathrm{R}\mathcal{H}om(\mathcal{K}[d_Y] \circ^a \mu\mathcal{O}_Y, \mu\mathcal{O}_X) \quad (9.4.1)$$

Hence the identity of $\mu\mathcal{O}_Y$ and any section

$$s \in H^0 \mathrm{R}\mathcal{H}om(\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(0, d_Y)}).$$

defines a morphism

$$\varphi(s) : \mathcal{K}[d_Y] \circ^a \mu\mathcal{O}_Y \longrightarrow \mu\mathcal{O}_X$$

In the sequel we will only consider kernels $\mathcal{K} \in D^b(\mathbb{C}_{T^*X \times T^*Y})$ satisfying

- (i) $\mathcal{K}$ is $\mathbb{R}$ -constructible,
- (ii) $(\Omega_Y \times T^*X \cup T^*Y \times \Omega_X^a) \cap \mathrm{SS}(\mathcal{K}) \subset \Lambda$,
- (iii) $\mathcal{K}$ is simple¹⁴ with shift 0 along Λ .

Note that given a $\mathbb{C}^\times$ -conic Lagrangian subvariety Λ that is associated to a contact transformation, there always locally (on $P^*(X \times Y)$) exists $\mathcal{K}$, i.e. for each $p \in \Omega_X$ there exists a $\mathbb{C}^\times$ -conic open neighborhood Ω'_X of $\mathbb{C}^\times p$ such that (i), (ii) and (iii) are satisfied. Recall the morphism

$$\begin{aligned} p_{13*}^a \left(p_{12}^{a-1} H^0 \mathrm{R}\mathcal{H}om(\mathcal{K}_1, \mu\mathcal{O}_{X \times Y}^{(0, d_Y)}) \otimes_{\beta\mathcal{E}_Y^{\mathbb{R}}} p_{23}^{a-1} \mathrm{R}\mathcal{H}om(\mathcal{K}_2, \mu\mathcal{O}_{Y \times Z}^{(0, d_Z)}) \right) \\ \longrightarrow \mathrm{R}\mathcal{H}om(\mathcal{K}_1 \circ^a \mathcal{K}_2[d_Y], \mu\mathcal{O}_{X \times Z}^{(0, d_Z)}). \end{aligned}$$

Denote by $s \circ s'$ the image of $s \otimes s'$ by this morphism.

Then we get

¹⁴For the definition of simple sheaves see [KS1], Section 7.5.

Proposition 9.4.1.

$$\varphi(s \circ s') = \varphi(s) \circ (\mathcal{K}[n] \overset{a}{\circ} \alpha(s'))$$

Proof. This follows from the fact that the morphism of Proposition 9.3.4 satisfies to the obvious associativity condition. $\square$

Theorem 9.4.2. *For every $p \in \Omega_X$ there exists a $\mathbb{C}^\times$ -conic open neighborhood $\Omega'_X \subset \Omega_X$ of $\mathbb{C}^\times p$ such that if we set $\Omega'_Y = \chi(\Omega_Y)$ we can find a section*

$$s \in H^0 \operatorname{RHom}(\mu\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(d_X, 0)})|_{\Omega'_X \times \Omega'_Y}$$

such that the morphisms

$$\begin{aligned} \mathcal{E}_X^{\mathbb{R}}|_{\Omega_X} &\longrightarrow p_{1*} H^0 \operatorname{RHom}(\mu\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(d_X, 0)})|_{\Omega'_X \times \Omega'_Y} & ; \quad P \mapsto Ps \\ \mathcal{E}_Y^{\mathbb{R}}|_{\Omega_Y} &\longrightarrow p_{2*} H^0 \operatorname{RHom}(\mu\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{(d_X, 0)})|_{\Omega'_X \times \Omega'_Y} & ; \quad Q \mapsto sQ \end{aligned}$$

are isomorphisms and we get an antiisomorphism

$$\chi_* \mathcal{E}_X^{\mathbb{R}}|_{\Omega_X} \xrightarrow{\sim} \mathcal{E}_Y^{\mathbb{R}}|_{\Omega_Y} \quad ; \quad P \mapsto Q \text{ such that } Ps = sQ$$

For such a section s the morphism

$$\varphi(s) : \mathcal{K}[d_Y] \overset{a}{\circ} \mu\mathcal{O}_Y \longrightarrow \mu\mathcal{O}_X$$

is an isomorphism of $\beta\mathcal{E}_X^{\mathbb{R}}$ -modules.

Corollary 9.4.3. *Let s be a section as in Theorem 9.4.2 and $\mathcal{F}$ be a perverse sheaf. Then $\varphi(s)$ defines an isomorphism of $\mathcal{E}_X^{\mathbb{R}}$ -modules*

$$\chi_* \operatorname{RHom}(\pi^{-1} \mu\mathcal{F}, \mu\mathcal{O}_Y) \simeq \operatorname{RHom}(\mu\mathcal{K} \overset{a}{\circ} \mu\mathcal{F}, \mu\mathcal{O}_X).$$

Corollary 9.4.4. *Let s be a section as in Theorem 9.4.2 and $\mathcal{F}$ be a coherent $\mathcal{E}_Y$ -module. Then $\varphi(s)$ defines an isomorphism*

$$\operatorname{RJHom}_{\beta\mathcal{E}_Y}(\beta\mathcal{F}, \mu\mathcal{O}_Y) \simeq \operatorname{RJHom}_{\beta\mathcal{E}_X}(\chi_* \beta\mathcal{F}, \mu\mathcal{O}_X)$$

The tempered version of Theorem 9.4.2 is similar with the only difference that the microdifferential structures are formal structures in the derived categories.

Theorem 9.4.5. *For every $p \in \Omega_X$ there exists a $\mathbb{C}^\times$ -conic open neighborhood $\Omega'_X \subset \Omega_X$ of $\mathbb{C}^\times p$ such that if we set $\Omega'_Y = \chi(\Omega_Y)$ we can find a section*

$$s \in H^0 \operatorname{RHom}(\mu\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{t, (d_X, 0)})|_{\Omega'_X \times \Omega'_Y}$$

such that the morphisms

$$\begin{aligned} \mathcal{E}_X^{\mathbb{R}, f}|_{\Omega_X} &\longrightarrow p_{1*} H^0 \operatorname{RHom}(\mu\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{t, (d_X, 0)})|_{\Omega'_X \times \Omega'_Y} & ; \quad P \mapsto Ps \\ \mathcal{E}_Y^{\mathbb{R}, f}|_{\Omega_Y} &\longrightarrow p_{2*} H^0 \operatorname{RHom}(\mu\mathcal{K}, \mu\mathcal{O}_{X \times Y}^{t, (d_X, 0)})|_{\Omega'_X \times \Omega'_Y} & ; \quad Q \mapsto sQ \end{aligned}$$

are isomorphisms and we get an antiisomorphism

$$\chi_* \mathcal{E}_X^{\mathbb{R}, f}|_{\Omega_X} \xrightarrow{\sim} \mathcal{E}_Y^{\mathbb{R}, f}|_{\Omega_Y} \quad ; \quad P \mapsto Q \text{ such that } Ps = sQ$$

For such a section s the morphism

$$\varphi^t(s) : \mathcal{K}[d_Y] \overset{a}{\circ} \mu\mathcal{O}_Y^t \longrightarrow \mu\mathcal{O}_X^t$$

is an isomorphism of formal $\beta\mathcal{E}_X^{\mathbb{R}, f}$ -modules in $D^b(I(k_X))$.

Corollary 9.4.6. *Let s be a section as in Theorem 9.4.5 and $\mathcal{F}$ be a perverse sheaf. Then $\varphi(s)$ defines an isomorphism of $\mathcal{E}_X^{\mathbb{R}, f}$ -modules*

$$\chi_* \operatorname{RHom}(\pi^{-1} \mu\mathcal{F}, \mu\mathcal{O}_Y^t) \simeq \operatorname{RHom}(\mu\mathcal{K} \overset{a}{\circ} \mu\mathcal{F}, \mu\mathcal{O}_X^t).$$

9.5 Classical Riemann-Hilbert Theorem and Ind-sheaves

Using the functor μ we can reformulate the isomorphisms of the classical Riemann-Hilbert Theorem from the microlocal point of view.

Lemma 9.5.1. (1) Let $\mathcal{F} \in D_{\mathbb{C}-c}^b(k_X)$. Then

$$\begin{aligned} \mathrm{R}\mathcal{H}om(\mu\mathcal{F}, \mu\mathcal{O}_X^t) &\simeq \mathcal{E}_X^{\mathbb{R},f} \underset{\pi^{-1}\mathcal{D}_X}{\overset{\mathrm{L}}{\otimes}} \pi^{-1} \mathrm{RH}(\mathcal{F}), \\ \gamma^{-1} \mathrm{R}\gamma_* \mathrm{R}\mathcal{H}om(\mu\mathcal{F}, \mu\mathcal{O}_X^t) &\simeq \mathcal{E}_X \underset{\pi^{-1}\mathcal{D}_X}{\overset{\mathrm{L}}{\otimes}} \pi^{-1} \mathrm{RH}(\mathcal{F}). \end{aligned}$$

(2) Let $\mathcal{M}$ be a coherent $\mathcal{D}_X$ -module. Then

$$\begin{aligned} \mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X^{\mathbb{R},f}}(\beta\mathcal{E}_X^{\mathbb{R},f} \underset{\pi^{-1}\beta\mathcal{D}_X}{\overset{\mathrm{L}}{\otimes}} \beta(\pi^{-1}\mathcal{M}), \mu\mathcal{O}_X) &\simeq \mu(\mathrm{Sol}(\mathcal{M})), \\ \mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\mathcal{E}_X \underset{\pi^{-1}\beta\mathcal{D}_X}{\overset{\mathrm{L}}{\otimes}} \beta(\pi^{-1}\mathcal{M}), \mu\mathcal{O}_X) &\simeq \mu(\mathrm{Sol}(\mathcal{M})). \end{aligned}$$

Proof. (1) is a reformulation of Theorem 4.2.6 of [An2] in terms of ind-sheaves using the fact that $\gamma^{-1} \mathrm{R}\gamma_* \mathcal{E}_X^{\mathbb{R},f} \simeq \mathcal{E}_X$ and (2) follows directly from the fact that $\mathcal{M}$ is a coherent $\mathcal{D}_X$ -module. $\square$

Therefore we can now formulate Riemann-Hilbert Theorem in terms of ind-sheaves:

Proposition 9.5.2. (1) Let $\mathcal{F}$ be a perverse sheaf on X . Then

$$\begin{aligned} \mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X^{\mathbb{R},f}}(\beta \mathrm{R}\mathcal{H}om(\mu\mathcal{F}, \mu\mathcal{O}_X^t), \mu\mathcal{O}_X) &\simeq \mu\mathcal{F} \\ \mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\gamma^{-1} \mathrm{R}\gamma_* \mathrm{R}\mathcal{H}om(\mu\mathcal{F}, \mu\mathcal{O}_X^t), \mu\mathcal{O}_X) &\simeq \mu\mathcal{F} \end{aligned}$$

(2) Let $\mathcal{M}$ be a regular holonomic $\mathcal{D}_X$ -module. Then

$$\begin{aligned} \mathrm{R}\mathcal{H}om(\mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X^{\mathbb{R},f}}(\beta(\mathcal{E}_X^{\mathbb{R},f} \underset{\pi^{-1}\mathcal{D}_X}{\overset{\mathrm{L}}{\otimes}} \pi^{-1}\mathcal{M}), \mu\mathcal{O}_X), \mu\mathcal{O}_X^t) &\simeq \mathcal{E}_X^{\mathbb{R},f} \underset{\pi^{-1}\mathcal{D}_X}{\overset{\mathrm{L}}{\otimes}} \pi^{-1}\mathcal{M} \\ \gamma^{-1} \mathrm{R}\gamma_* \mathrm{R}\mathcal{H}om(\mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta(\mathcal{E}_X \underset{\pi^{-1}\mathcal{D}_X}{\overset{\mathrm{L}}{\otimes}} \pi^{-1}\mathcal{M}), \mu\mathcal{O}_X), \mu\mathcal{O}_X^t) &\simeq \mathcal{E}_X \underset{\pi^{-1}\mathcal{D}_X}{\overset{\mathrm{L}}{\otimes}} \pi^{-1}\mathcal{M} \end{aligned}$$

Now let us formulate the comparison theorem for regular holonomic $\mathcal{E}_X$ -modules in terms of ind-sheaves. The classical version ([An1], Proposition 5.6.3) states

Proposition 9.5.3. Let $\mathcal{M}$ be a regular holonomic $\mathcal{E}_X$ -module and $\mathcal{G} \in D_{\mathbb{R}-c}^b(k_X)$ such that $t\mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X)$ is concentrated in a single degree. Then the natural morphism

$$\mathrm{R}\mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, t\mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X)) \longrightarrow \mathrm{R}\mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, \mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X))$$

is an isomorphism.

Remark 9.5.4. Note that we do not have to assume that $\mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X)$ is concentrated in degree zero because we now know that $\mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X)$ is well defined in the derived category of microdifferential modules.

It is slightly more complicated to deal with $\mu\mathcal{O}_X^t$ since we do not know if $\mu\mathcal{O}_X^t$ is well defined in the derived category of microdifferential modules. The analog of the last proposition should be the formula

$$\mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mu\mathcal{O}_X) \xrightarrow{\sim} \mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mu\mathcal{O}_X^t)$$

for any regular holonomic $\mathcal{E}_X$ -module $\mathcal{M}$. However the second term of this isomorphism is unfortunately not (yet) well-defined. Therefore we only get the following weaker statement:

Proposition 9.5.5. *Let $\mathcal{M}$ be a regular holonomic $\mathcal{E}_X$ -module. Then there is a natural morphism*

$$\mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mu\mathcal{O}_X) \longrightarrow \mathrm{R}\mathcal{H}om(\beta\mathcal{M}, \mu\mathcal{O}_X^t).$$

Proof. By the comparison theorem if $\mathcal{G} \in \mathrm{D}_{\mathrm{perv}}^b(\mathbb{C}_X, U)$ we have a natural morphism

$$\mathrm{R}\mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, \mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X)) \simeq \mathrm{R}\mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, t\mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X)) \longrightarrow \mathrm{R}\mathcal{H}om(\mathcal{M}, t\mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X)).$$

Since we have

$$\begin{aligned} \mathrm{R}\mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, \mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X)) &\simeq \mathrm{R}\mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, \mathrm{R}\mathcal{H}om(\mu\mathcal{G}, \mu\mathcal{O}_X)) \\ &\simeq \mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mathrm{R}\mathcal{H}om(\mu\mathcal{G}, \mu\mathcal{O}_X)) \\ &\simeq \mathrm{R}\mathcal{H}om(\mu\mathcal{G}, \mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mu\mathcal{O}_X)) \end{aligned}$$

and

$$\begin{aligned} \mathrm{R}\mathcal{H}om(\mathcal{M}, t\mu\mathrm{hom}(\mathcal{G}, \mathcal{O}_X)) &\simeq \mathrm{R}\mathcal{H}om(\mathcal{M}, \mathrm{R}\mathcal{H}om(\mu\mathcal{G}, \mu\mathcal{O}_X^t)) \\ &\simeq \mathrm{R}\mathcal{H}om(\beta\mathcal{M}, \mathrm{R}\mathcal{H}om(\mu\mathcal{G}, \mu\mathcal{O}_X^t)) \\ &\simeq \mathrm{R}\mathcal{H}om(\mu\mathcal{G}, \mathrm{R}\mathcal{H}om(\beta\mathcal{M}, \mu\mathcal{O}_X^t)) \end{aligned}$$

we get a morphism

$$\mathrm{R}\mathcal{H}om(\mu\mathcal{G}, \mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mu\mathcal{O}_X)) \longrightarrow \mathrm{R}\mathcal{H}om(\mu\mathcal{G}, \mathrm{R}\mathcal{H}om(\beta\mathcal{M}, \mu\mathcal{O}_X^t)).$$

Now recall that $\mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\mathcal{M}, \mu\mathcal{O}_X)$ is a microlocal perverse sheaf. Hence locally it is of the form $\mu\mathcal{G}$ for some object $\mathcal{G} \in \mathrm{D}_{\mathrm{perv}}^b(\mathbb{C}_X, U)$. Thus locally the identity morphism of $\mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\mathcal{M}, \mu\mathcal{O}_X)$ defines the desired morphism and we may patch it because microlocal perverse sheaves form a stack. $\square$

Remark 9.5.6. If $\mu\mathcal{O}_X^t$ was well-defined in the derived category of $\beta\mathcal{E}_X$ -modules then the proof of the last Proposition would establish the isomorphism

$$\mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mu\mathcal{O}_X) \xrightarrow{\sim} \mathrm{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\mathcal{M}, \mu\mathcal{O}_X^t).$$

9.6 Microlocal Riemann-Hilbert morphism

Lemma 9.6.1. *Let $\mathcal{F} \in \mu\mathrm{Perv}(\Omega)$ and set $U = \gamma^{-1}(\Omega)$. Then*

$$\mathrm{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t|_U)$$

is an $\mathcal{E}_X^{\mathbb{R},f}|_U$ -module. Moreover

$$\gamma^{-1}\mathrm{R}\gamma_*\mathrm{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t|_U)$$

is a regular holonomic $\mathcal{E}_X|_U$ -module.

Proof. First we will show that $\mathrm{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t|_U)$ is a well-defined $\mathcal{E}_X^{\mathbb{R},f}|_U$ -module.

By Proposition 9.3.8 it is enough to prove that the complex

$$\mathrm{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t|_U)$$

is concentrated in a single degree. This is a local problem. Thus we may assume that $\mathcal{F} \simeq \mu\tilde{\mathcal{F}}$ where $\tilde{\mathcal{F}}$ is an object of $\mathrm{D}_{\mathrm{perv}}^b(k_X, U)$. Therefore

$$\mathrm{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t|_U) \simeq \mathrm{R}\mathcal{H}om(\mu\tilde{\mathcal{F}}|_U, \mu\mathcal{O}_X^t|_U) \simeq t\mu\mathrm{hom}(\tilde{\mathcal{F}}, \mathcal{O}_X)|_U$$

Since $t\mu\text{hom}(\tilde{\mathcal{F}}, \mathcal{O}_X)|_U$ is invariant under quantized contact transformations, we can suppose that $\tilde{\mathcal{F}}$ is a perverse sheaf. Then the complex $t\mu\text{hom}(\tilde{\mathcal{F}}, \mathcal{O}_X)|_U$ is concentrated in degree 0. Hence $\text{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t|_U)$ is a well-defined $\mathcal{E}_X^{\mathbb{R},f}|_U$ -module. Hence $\gamma^{-1}\text{R}\gamma_*\text{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t|_U)$ is an $\mathcal{E}_X$ -module. Let us show that it is regular holonomic. This is again a local question, invariant by quantized contact transformations. Therefore we may assume that $\mathcal{F} \simeq \mu\tilde{\mathcal{F}}$ for a perverse sheaf $\tilde{\mathcal{F}}$. Recall that

$$t\mu\text{hom}(\tilde{\mathcal{F}}, \mathcal{O}_X)|_U \simeq (\mathcal{E}_X^{\mathbb{R},f} \otimes_{\pi^{-1}\mathcal{D}_X} \pi^{-1}\text{T}\mathcal{H}om(\tilde{\mathcal{F}}, \mathcal{O}_X))|_U.$$

By the Riemann-Hilbert theorem $\text{T}\mathcal{H}om(\tilde{\mathcal{F}}, \mathcal{O}_X)$ is a regular holonomic $\mathcal{D}_X$ -module. Hence $\gamma^{-1}\text{R}\gamma_*\text{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t|_U)$ is regular holonomic. $\square$

Lemma 9.6.2. *Let $\mathcal{M} \in \mathcal{H}ol\mathcal{R}eg(U)$. Then*

$$\text{R}\mathcal{H}om_{\beta\mathcal{E}_X|_U}(\beta\mathcal{M}, \mu\mathcal{O}_X|_U)$$

is an object of $\mu\text{Perv}(\Omega)$.

Proof. This is a local problem, invariant by quantized contact transformations. Therefore we may assume that $\mathcal{M}$ is isomorphic to $\mathcal{E}_X \otimes_{\pi^{-1}\mathcal{D}_X} \pi^{-1}\tilde{\mathcal{M}}$ on U where $\tilde{\mathcal{M}}$ is a regular holonomic $\mathcal{D}_X$ -module. Then

$$\text{R}\mathcal{H}om_{\beta\mathcal{E}_X|_U}(\beta\mathcal{M}, \mu\mathcal{O}_X|_U) \simeq \text{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\mathcal{E}_X \otimes_{\pi^{-1}\mathcal{D}_X} \pi^{-1}\tilde{\mathcal{M}}, \mu\mathcal{O}_X)|_U \simeq \mu\text{R}\mathcal{H}om_{\mathcal{D}_X}(\tilde{\mathcal{M}}, \mathcal{O}_X)|_U$$

By the Riemann-Hilbert Theorem $\text{R}\mathcal{H}om_{\mathcal{D}_X}(\tilde{\mathcal{M}}, \mathcal{O}_X)$ is a perverse sheaf. $\square$

For any open subset $\Omega \subset P^*X$ let us define the microlocal Riemann-Hilbert correspondence:

$$\mu\text{Perv}(\Omega) \begin{array}{c} \xrightarrow{\mu\text{RH}} \\ \xleftarrow{\mu\text{Sol}} \end{array} \mathcal{H}ol\mathcal{R}eg(\mathcal{E}_X|_{\gamma^{-1}\Omega}).$$

by the formulas

$$\mu\text{Sol}(\mathcal{M}) = \text{R}\mathcal{H}om_{\beta(\mathcal{E}_X|_{\gamma^{-1}\Omega})}(\beta(\mathcal{M}), \mu\mathcal{O}_X|_{\gamma^{-1}\Omega})$$

$$\mu\text{RH}(\mathcal{F}) = \gamma_\Omega^{-1}\text{R}\gamma_{\Omega*}(\text{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t|_{\gamma^{-1}(\Omega)}))$$

where γ_Ω is the restriction of γ to $\gamma^{-1}\Omega$.

The functors μSol and μRH are obviously functors of stacks.

Lemma 9.6.3. *There is a natural morphism*

$$\text{Id} \longrightarrow \mu\text{Sol} \circ \mu\text{RH}. \quad (9.6.1)$$

Proof. Let $\mathcal{F}$ be a microlocal perverse sheaf. We will define the morphism (9.6.1) by a natural element of

$$\text{Hom}_{\text{D}^b(\text{I}(k_X))}(\mathcal{F}, \text{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\gamma^{-1}\text{R}\gamma_*\text{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t), \mu\mathcal{O}_X)).$$

Note that

$$\begin{aligned} \text{Hom}_{\text{D}^b(\text{I}(k_X))}(\mathcal{F}, \text{R}\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\gamma^{-1}\text{R}\gamma_*\text{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t), \mu\mathcal{O}_X)) \\ \simeq \text{Hom}_{\text{D}^b(\text{I}(\beta\mathcal{E}_X))}(\mathcal{F} \otimes \beta\gamma^{-1}\text{R}\gamma_*\text{R}\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t), \mu\mathcal{O}_X). \end{aligned}$$

Now recall that

$$\beta\gamma^{-1} R\gamma_* R\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X^t) \longrightarrow \beta\gamma^{-1} R\gamma_* R\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X)$$

is $\beta\mathcal{E}_X$ -linear. Hence the natural morphism in $D^b(I(\beta\mathcal{E}_X))$

$$\mathcal{F} \otimes \beta\gamma^{-1} R\gamma_* R\mathcal{H}om(\mathcal{F}, \mu\mathcal{O}_X) \longrightarrow \mu\mathcal{O}_X$$

defines the morphism of the lemma. $\square$

Theorem 9.6.4. *The functors μSol and μRH define quasi-inverse equivalences of stacks*

$$\mu\text{Perv} \xrightleftharpoons[\mu\text{Sol}]{\mu\text{RH}} \gamma_* \mathcal{H}ol\mathcal{R}eg(\mathcal{E}_X).$$

Proof. First let us show that the morphism of Lemma 9.6.3 is an isomorphisms.

Let $\mathcal{F}$ be a microlocal perverse sheaf defined in a neighborhood at p . Then there exists $\tilde{\mathcal{F}} \in D_{\text{perv}}^b(k_X, \mathbb{C}^\times p)$ such that $\mathcal{F} \simeq \mu\tilde{\mathcal{F}}$. Let χ be a contact transformation such that $\chi(\text{SS}(\tilde{\mathcal{F}}))$ is in generic position at $\chi(p)$. Then

$$\Phi_{\mathcal{K}} R\mathcal{H}om_{\beta\mathcal{E}_X}(\beta R\mathcal{H}om(\mu\tilde{\mathcal{F}}, \mu\mathcal{O}_X^t), \mu\mathcal{O}_X) \simeq R\mathcal{H}om_{\beta\mathcal{E}_X}(\beta R\mathcal{H}om(\Phi_{\mathcal{K}}\tilde{\mathcal{F}}, \mu\mathcal{O}_X^t), \mu\mathcal{O}_X)$$

Since $\Phi_{\mathcal{K}}(\tilde{\mathcal{F}})$ is isomorphic to a perverse sheaf in a neighborhood of $\pi(p)$ the isomorphism follows from the second part of Proposition 9.5.2.

Now let us show that the functor μRH is an equivalence of stacks. It is sufficient to prove this locally. In order to prove that μRH is essentially surjective, we show that μRH and μSol are inverse to each other on the level of objects. We have already seen that $\mu\text{Sol} \circ \mu\text{RH}(\mathcal{F}) \simeq \mathcal{F}$ for any microlocal perverse sheaf $\mathcal{F}$.

Let $\mathcal{M}$ be a regular holonomic $\mathcal{E}_X$ -module defined in a neighborhood of $p \in \dot{T}^*X$. Let χ be a contact transformation such that $\chi_*\mathcal{M}$ is in generic position at $\chi(p)$. Then

$$\begin{aligned} \chi_* R\gamma^{-1} R\gamma_* R\mathcal{H}om(R\mathcal{H}om_{\beta\mathcal{E}_X}(\beta(\mathcal{M}), \mu\mathcal{O}_X), \mu\mathcal{O}_X^t) \\ \simeq \gamma^{-1} R\gamma_* R\mathcal{H}om(\Phi_{\mathcal{K}}^\mu R\mathcal{H}om_{\beta\mathcal{E}_X}(\beta(\mathcal{M}), \mu\mathcal{O}_X), \mu\mathcal{O}_X^t) \\ \simeq \gamma^{-1} R\gamma_* R\mathcal{H}om(R\mathcal{H}om_{\beta\mathcal{E}_X}(\beta\chi_*\mathcal{M}, \mu\mathcal{O}_X), \mu\mathcal{O}_X^t) \end{aligned}$$

But since $\chi_*\mathcal{M}$ is in generic position there exists a regular holonomic $\mathcal{D}_X$ -module $\tilde{\mathcal{M}}$ such that

$$\mathcal{M} \simeq \mathcal{E}_X \bigotimes_{\pi^{-1}\mathcal{D}_X}^{\text{L}} \pi^{-1}\tilde{\mathcal{M}}.$$

Hence μRH is essentially surjective.

Let us show that μRH is fully faithful. Let $\mathcal{F}, \mathcal{G}$ be microlocal perverse sheaves. By invariance under quantized contact transformations we may assume that there exists perverse sheaves $\tilde{\mathcal{F}}, \tilde{\mathcal{G}}$ such that $\mathcal{F} \simeq \mu\tilde{\mathcal{F}}$ and $\mathcal{G} \simeq \mu\tilde{\mathcal{G}}$. Then the fact that μRH is fully faithful follows from the well-known formula

$$\mu\text{hom}(\mathcal{F}, \mathcal{G}) \simeq R\mathcal{H}om(\text{RH}(\mathcal{G}), \text{RH}(\mathcal{F})).$$

$\square$

A Stacks on Topological Spaces

In this section we will make use of the language of 2-functors and 2-colimits. But since we only work with the 2-category $\mathcal{CAT}$ of all (small) categories, we will not recall the abstract theory and refer to [St],[McL].

A.1 Prestacks

A prestack is a “presheaf of categories up to equivalence”. More precisely, let X be a topological space and denote by $\mathcal{T}(X)$ the category of open sets of X . A prestack on X is just a 2-functor $\mathcal{T}(X)^\circ \rightarrow \mathcal{CAT}$. Therefore many authors call a 2-functor with domain $\mathcal{T}$ a prestack on $\mathcal{T}$.

Since the category $\mathcal{T}(X)$ is particularly simple, we will recall here a detailed description.

Definition A.1.1. A *prestack* $(\mathcal{C}, \rho, \Phi)$ on X consists of

- (1) a small category $\mathcal{C}(U)$ for every open set $U \subset X$,
- (2) a functor $\rho_{VU} : \mathcal{C}(U) \rightarrow \mathcal{C}(V)$ for any two open sets $V \subset U \subset X$, called *restriction functor* or just *restriction*,
- (3) a natural equivalence $\Phi_{WVU} : \rho_{WU} \xrightarrow{\sim} \rho_{WV} \rho_{VU}$ for every three open sets $W \subset V \subset U \subset X$.

This data should satisfy

(P2F1) $\rho_{UU} = Id_{\mathcal{C}(U)}$ for every open set $U \subset X$.

(P2F2) $\Phi_{UUU} = Id_{Id_{\mathcal{C}(U)}}$ for every open set $U \subset X$.

(P2F3) For every four open sets $T \subset W \subset V \subset U \subset X$ the equation

$$(\rho_{TW} \bullet \Phi_{WVU}) \circ \Phi_{TWU} = (\Phi_{TWV} \bullet \rho_{VU}) \circ \Phi_{TVU},$$

holds, i. e. the following diagram commutes:

$$\begin{array}{ccc} \rho_{TU} & \xrightarrow[\sim]{\Phi_{TWU}} & \rho_{TW} \rho_{WU} \\ \Phi_{TVU} \downarrow \sim & & \sim \downarrow \rho_{TW} \bullet \Phi_{WVU} \\ \rho_{TV} \rho_{VU} & \xrightarrow[\sim]{\Phi_{TWV} \bullet \rho_{VU}} & \rho_{TW} \rho_{WV} \rho_{VU} \end{array}$$

We shall mostly denote a prestack $(\mathcal{C}, \rho, \Phi)$ by $\mathcal{C}$ for short. Moreover it is often convenient to denote the restriction functor ρ_{VU} by i_{VU}^{-1} .

Hence a prestack is just a contravariant 2-functor $\mathcal{C} : \mathcal{T}(X) \rightarrow \mathcal{CAT}$ with strict identities. We therefore immediately get the notion of a functor of prestacks (being a 2-natural transformation of the underlying 2-functors) and the notion of a natural transformation of functors of prestacks (being a modification of the underlying 2-natural transformations). In particular we get the concept of an equivalence of prestacks and we may define the (2-)category $\mathcal{PST}(X)$ of prestacks on X .

Remark A.1.2. Let $\mathcal{C}$ be a prestack on X , $U \subset X$ an open subset and $A, B \in \text{Ob } \mathcal{C}(U)$. For $V \subset U$, we set

$$\mathcal{H}om_{e|_U}(A, B)(V) = \text{Hom}_{\mathcal{C}(V)}(A|_V, B|_V)$$

If $W \subset V \subset U$ the restriction functor ρ_{WV} and the natural equivalence Φ_{WVU} define a restriction map and one easily verifies that $\mathcal{H}om_{e|_U}(A, B)$ is a presheaf.

Let us add some notations

Definition A.1.3. (1) A prestack $\mathcal{C}$ is called additive if for any $U \subset X$ the category $\mathcal{C}(U)$ is additive and the restriction functors are additive.

(2) An additive prestack $\mathcal{C}$ is called triangulated if for any $U \subset X$ the category $\mathcal{C}(U)$ is triangulated and the restriction functors are exact.

(3) An additive prestack $\mathcal{C}$ is called abelian if for any $U \subset X$ the category $\mathcal{C}(U)$ is abelian and the restriction functors are exact.

We then get the obvious concept of an additive (resp. exact) functor between additive (resp. triangulated or abelian) prestacks.

A.2 Operations on prestacks

Let $f : X \rightarrow Y$ be a continuous map.

Proposition A.2.1. Let $\mathcal{C}$ be a prestack on X . Then there is a natural prestack $f_*\mathcal{C}$ on Y such that for any open set $V \subset Y$ we have a canonical equivalence $f_*\mathcal{C}(V) \simeq \mathcal{C}(f^{-1}(V))$.

Proposition A.2.2. Let $\mathcal{C}$ be a prestack on Y . Then there is a natural prestack $f_p^{-1}\mathcal{C}$ on X such that for any open set $U \subset X$ we have a canonical equivalence $f_p^{-1}\mathcal{C}(U) \simeq \varinjlim_{f(U) \subset V} \mathcal{C}(V)$.

Proposition A.2.3. The operations f_* and f_p^{-1} are (2-)adjoint to each other, i.e. there is a (2-)natural equivalence of categories

$$\underline{\mathrm{Hom}}_{\mathcal{PST}(X)}(f_p^{-1}\mathcal{C}, \mathcal{D}) \simeq \underline{\mathrm{Hom}}_{\mathcal{PST}(Y)}(\mathcal{C}, f_*\mathcal{D}).$$

A.3 Stalks

Since prestacks (and stacks) are often treated in the more general framework of sites, we will describe here in detail the notion of a stalk of a prestack on a topological space. Of course if $p \in X$ and $i : \{p\} \hookrightarrow X$ is the inclusion, then the stalk $\mathcal{C}_p$ of a prestack $\mathcal{C}$ at p is nothing but $i_p^{-1}\mathcal{C}$.

Definition A.3.1. Let $\mathcal{C}$ be a prestack on X and $p \in X$ a point.

Consider the category $\mathcal{T}_p(X)$ of open sets that contain the point p . Note that since the set of open sets containing p is stable by union and intersection the category $\mathcal{T}_p(X)$ is filtered and cofiltered.

The prestack $\mathcal{C}$ induces a 2-functor $\alpha_p : \mathcal{T}_p(X)^\circ \rightarrow \mathcal{CAT}$.

We set

$$\mathcal{C}_p = \varinjlim_{U \ni p} \mathcal{C}(U) = \varinjlim_{U \in \mathcal{T}_p(X)} \alpha_p(U)$$

and call $\mathcal{C}_p$ the **stalk** of $\mathcal{C}$ at p or the **category of germs** of $\mathcal{C}$ at p .

Hence the stalk of a prestack is defined up to canonical equivalence of categories. It can easily be described using the explicit construction of 2-colimits. We get

$$\mathrm{Ob} \mathcal{C}_p = \left\{ (U, A) \mid p \in U \subset X \text{ open and } A \in \mathrm{Ob} \mathcal{C}(U) \right\} = \bigsqcup_{p \in U \subset X} \mathrm{Ob} \mathcal{C}(U).$$

Let $(U, A), (V, B)$ be two objects of $\mathcal{C}_p$. Then

$$\mathrm{Hom}_{\mathcal{C}_p}((U, A), (V, B)) = \varinjlim_{p \in W \subset U \cap V} \mathrm{Hom}_{\mathcal{C}(W)}(A|_W, B|_W).$$

Hence a morphism $f : (U, A) \rightarrow (V, B)$ is defined on a small neighborhood $W \subset U \cap V$ of p . In particular we get

Proposition A.3.2. *Let $\mathcal{C}$ be a prestack on X , $p \in X$ a point, $U \subset X$ an open set containing p and $A, B \in \text{Ob } \mathcal{C}(U)$ two objects. Then we have a canonical isomorphism*

$$\mathcal{H}om_{e|_U}(A, B)_p \xrightarrow{\sim} \text{Hom}_{e_p}(A, B).$$

This isomorphism is compatible with the composition maps in $\mathcal{C}$ in the following way:

Let $U \subset X$ be an open set containing p and $A, B, C \in \text{Ob } \mathcal{C}(U)$.

Then the morphism of sheaves

$$\mathcal{H}om_{e|_U}(A, B) \times \mathcal{H}om_{e|_U}(B, C) \longrightarrow \mathcal{H}om_{e|_U}(A, C)$$

induces in its stalks the composition in the stalk:

$$\text{Hom}_{e_p}(A, B) \times \text{Hom}_{e_p}(B, C) \longrightarrow \text{Hom}_{e_p}(A, C).$$

Hence germs of morphisms may be seen as morphisms in the category of germs. The compatibility with the composition map has an obvious corollary:

Corollary A.3.3. *Let $\mathcal{C}$ be a prestack on X , $p \in X$ a point, $U, V \subset X$ two open sets containing p and $A \in \text{Ob } \mathcal{C}(U)$, $B \in \text{Ob } \mathcal{C}(V)$ two objects. Then A and B are isomorphic in $\mathcal{C}_p$ if and only if they are isomorphic on an open neighborhood of p .*

Let us observe that an object $A \in \text{Ob } \mathcal{C}(U)$ (with $p \in U$) is isomorphic to all its restrictions to sets $V \ni p$ but there is no equivalence relation imposed on the objects. If $A \in \text{Ob } \mathcal{C}(U)$ we will still denote by A its image in $\mathcal{C}_p$. If $f : A \rightarrow B$ is a morphism in $\mathcal{C}(U)$ then we note $f_p : A \rightarrow B$ its image in $\mathcal{C}_p$. The reason why we do not write A_p is given by the following remark.

Remark A.3.4. Consider a sheaf of rings $\mathcal{A}$ and the stack $\mathcal{MOD}(\mathcal{A})$. One shall beware that the natural functor

$$\mathcal{MOD}(\mathcal{A})_p \longrightarrow \mathcal{MOD}(\mathcal{A}_p)$$

is not an equivalence of categories because the morphism

$$\text{Hom}(\mathcal{F}, \mathcal{G})_p \longrightarrow \text{Hom}_{\mathcal{A}_p}(\mathcal{F}_p, \mathcal{G}_p)$$

is not an isomorphism in general.

Proposition A.3.5. (1) *If $\mathcal{C}$ is additive then its stalks are additive categories and the natural functors into the stalks are additive.*

(2) *If $\mathcal{C}$ is triangulated then its stalks are triangulated categories and the natural functors into the stalks are exact.*

(3) *If $\mathcal{C}$ is abelian then its stalks are abelian categories and the natural functors into the stalks are exact.*

A.4 Stacks

Definition A.4.1. *A prestack $\mathcal{C}$ on X is **separated** if for all open subsets $U \subset X$ and all objects $A, B \in \text{Ob } \mathcal{C}(U)$ the presheaf*

$$\mathcal{H}om_{e|_U}(A, B)$$

is a sheaf.

Definition A.4.2. A prestack $\mathcal{C}$ on X is a **stack** if the following two conditions are satisfied

- (i) The prestack $\mathcal{C}$ is separated
- (ii) Let $U = \bigcup_{i \in I} U_i$ be an open covering of an open subset $U \subset X$ and suppose that we are given the following data
 - (a) for every $i \in I$ an object $A_i \in \mathcal{C}(U_i)$
 - (b) for every $i, j \in I$ an isomorphism $\sigma_{ij} : A_j|_{U_{ij}} \xrightarrow{\sim} A_i|_{U_{ij}}$ such that for any $i, j, k \in I$ the equation $\sigma_{jk} \circ \sigma_{ij} = \sigma_{ik}$ holds on U_{ijk} .

Then there exists an object $A \in \mathcal{C}(U)$ and isomorphisms $\rho_i : A|_{U_i} \rightarrow A_i$ such that $\sigma_{ij} \circ \rho_j = \rho_i$.

Proposition A.4.3. Let $\mathcal{C}, \mathcal{C}'$ be two stacks on X . Consider a functor $F : \mathcal{C} \rightarrow \mathcal{C}'$. Then we have

- (1) F is faithful if and only if F_p is faithful for all $p \in X$.
- (2) F is fully faithful if and only if F_p is fully faithful for all $p \in X$.
- (3) F is an equivalence of stacks if and only if F_p is an equivalence of categories for all $p \in X$.

Proof. We know that F is faithful (resp. fully faithful) if and only if the morphisms of sheaves

$$\mathcal{H}om_{\mathcal{C}|_U}(A, B) \longrightarrow \mathcal{H}om_{\mathcal{C}'|_U}(F(A), F(B))$$

are monomorphisms (resp. isomorphisms). Since these two properties are verified in the stalks, we immediately get (1) and (2).

Let's prove (3). Note that the condition is clearly necessary.

Now suppose that F_p is an equivalence of categories for all $p \in X$.

By (2) we know that F is fully faithful. Hence it is sufficient to show that F is essentially surjective for any open set $U \subset X$.

Let $A' \in \text{Ob } \mathcal{C}'(U)$. For any point $p \in U$ there is an object $A_p \in \mathcal{C}_p$ such that $F_p(A_p) \simeq A'_p$. Hence by corollary (A.3.3) there is an open neighborhood $V(p)$ of p , an object $A(p) \in \text{Ob } (\mathcal{C}(V(p)))$ and an isomorphism $\vartheta(p) : F(A(p)) \simeq A'|_{V(p)}$.

These isomorphisms define a cocycle that patches together the objects $F(A(p))$ to an object isomorphic to A' . Since F is fully faithful this cocycle can be lifted to a cocycle in $\mathcal{C}'$ where the $A(p)$ patch together to an object A such that $F(A)$ is isomorphic to A' . $\square$

A.5 The stack associated to a prestack

In this paragraph we will describe the stack associated to a prestack on a topological space. As is the case of the sheaf associated to a presheaf this can be done explicitly and is less complicated than on an arbitrary site.

Proposition A.5.1. Let $\mathcal{C}$ be a prestack on X . Then there exists a separated prestack $\mathcal{C}^\dagger$ on X and a canonical functor $\eta_{\mathcal{C}}^\dagger : \mathcal{C} \rightarrow \mathcal{C}^\dagger$ that induces an equivalence of categories on the stalks, such that any morphism $\mathcal{C} \rightarrow \mathcal{D}$ into a separated prestack $\mathcal{D}$ factors uniquely through $\mathcal{C}^\dagger$ (up to unique equivalence).

Moreover for any functor $F : \mathcal{C} \rightarrow \mathcal{D}$ there exists a functor $F^\dagger : \mathcal{C}^\dagger \rightarrow \mathcal{D}^\dagger$ such that the diagram commutes

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ \downarrow & & \downarrow \\ \mathcal{C}^\dagger & \xrightarrow{F^\dagger} & \mathcal{D}^\dagger. \end{array}$$

Proof.

Let $U \subset X$ be an open subset. Let us define a category $\mathcal{C}^\dagger(U)$. Set

$$\text{Ob } \mathcal{C}^\dagger(U) = \text{Ob } \mathcal{C}(U)$$

now let $A, B \in \text{Ob } \mathcal{C}(U)$ be two objects. Put

$$\text{Hom}_{\mathcal{C}^\dagger(U)}(A, B) = \Gamma(U, \mathcal{H}om_{\mathcal{C}|_U}(A, B)^\dagger),$$

where $\mathcal{H}om_{\mathcal{C}|_U}(A, B)^\dagger$ is the sheaf associated to $\mathcal{H}om_{\mathcal{C}|_U}(A, B)$. Note that we have a canonical map $\text{Hom}_{\mathcal{C}(U)}(A, B) \rightarrow \text{Hom}_{\mathcal{C}^\dagger(U)}(A, B)$ which is just the natural morphism from the presheaf $\text{Hom}_{\mathcal{C}(U)}(A, B)$ into its associated sheaf.

The map

$$\mathcal{H}om_{\mathcal{C}|_U}(A, B) \times \mathcal{H}om_{\mathcal{C}|_U}(B, C) \longrightarrow \mathcal{H}om_{\mathcal{C}|_U}(A, C)$$

induces the composition in $\mathcal{C}^\dagger$:

$$\mathcal{H}om_{\mathcal{C}^\dagger|_U}(A, B) \times \mathcal{H}om_{\mathcal{C}^\dagger|_U}(B, C) \longrightarrow \mathcal{H}om_{\mathcal{C}^\dagger|_U}(A, C).$$

The restriction functors of $\mathcal{C}^\dagger$ and the equivalences can easily be constructed by the universal property of the sheaf associated to a presheaf which also implies that all the axioms are verified.

The universal property of the separated prestack associated to a prestack also follows from the universal property of the sheaf associated to a presheaf. $\square$

Theorem A.5.2. *Let $\mathcal{C}$ be a prestack on X . Then there exists a stack $\mathcal{C}^\dagger$ on X together with a canonical functor of prestacks $\eta_{\mathcal{C}}^\dagger : \mathcal{C} \rightarrow \mathcal{C}^\dagger$ such that any morphism $\mathcal{C} \rightarrow \mathcal{D}$ into some stack $\mathcal{D}$ factors uniquely through $\mathcal{C}^\dagger$.*

Moreover for any stack $\mathcal{D}$ and any morphism of prestacks $F : \mathcal{C} \rightarrow \mathcal{D}$ there exists a canonical functor of stacks $F^\dagger$ such that the following diagram is commutative

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ \eta_{\mathcal{C}}^\dagger \downarrow & & \downarrow \eta_{\mathcal{D}}^\dagger \\ \mathcal{C}^\dagger & \xrightarrow{F^\dagger} & \mathcal{D}^\dagger. \end{array}$$

Finally the functor $\eta_{\mathcal{C}}^\dagger : \mathcal{C} \rightarrow \mathcal{C}^\dagger$ induces equivalences of categories on the stalks at every point of X .

Proof. By the proposition we may assume that $\mathcal{C}$ is a separated prestack, i.e. all associated presheaves are actually sheaves.

Let $U \subset X$ be an open subset. We have to define a category $\mathcal{C}^\dagger(U)$.

Consider families $A = \{(A_p, U_p^A)\}_{p \in U}$ where U_p^A is an open neighborhood of p with $A_p \in \mathcal{C}(U_p^A)$ and families of morphisms $\theta^A = \{\theta_{pq}^A\}_{p, q \in U}$ where $\theta_{pq}^A : A_q|_{U_{pq}^A} \xrightarrow{\sim} A_p|_{U_{pq}^A}$ is an

isomorphism for all $p, q \in U$ (here $U_{pq}^A = U_p^A \cap U_q^A$) satisfying the cocycle condition. During the proof let us call a pair (A, θ^A) a cocycle on U .

We shall now define morphisms of cocycles. A morphism $f : (A, \theta^A) \rightarrow (B, \theta^B)$ consists of a family of germs morphisms $f_p : (A_p, U_p^A) \rightarrow (B_p, U_p^B)$ of $\text{Mor } \mathcal{C}_p$ such that for any point $p \in U$ there is an open set U_p^f on which f_p is represented as a morphism $f_p : A_p|_{U_p^f} \rightarrow B_p|_{U_p^f}$ (where $U_p^f \subset U_p^{AB} = U_p^A \cap U_p^B$ is an open neighborhood of p) satisfying the following compatibility condition: the diagram

$$\begin{array}{ccc} A_q|_{U_{pq}^f} & \xrightarrow{\theta_{pq}^A|_{U_{pq}^f}} & A_p|_{U_{pq}^f} \\ f_q|_{U_{pq}^f} \downarrow & & \downarrow f_p|_{U_{pq}^f} \\ B_q|_{U_{pq}^f} & \xrightarrow{\theta_{pq}^B|_{U_{pq}^f}} & B_p|_{U_{pq}^f} \end{array}$$

should be commutative for all $p, q \in U$.

Now define $\mathcal{C}^\dagger(U)$ to be the category of cocycles. The obvious restriction maps define a prestack $\mathcal{C}^\dagger$ on U (which is actually a presheaf). It is now tedious but straightforward, that $\mathcal{C}^\dagger$ is a stack that satisfies the universal property. Note that we need the assumption that $\mathcal{C}$ is separated when proving the patching condition. $\square$

Corollary A.5.3. *Let $\mathcal{C}$ be a stack. Then there exists a stack $\mathcal{C}'$, canonically isomorphic to $\mathcal{C}$, which is also a presheaf of categories.*

Corollary A.5.4. *Let $\mathcal{C}$ be a prestack and $F : \mathcal{C} \rightarrow \mathcal{D}$ be a morphism into a stack $\mathcal{D}$. Suppose that F induces equivalences of categories in the stalks. Then $\mathcal{D}$ is equivalent to the stack associated to $\mathcal{C}$.*

A.6 Patching of stacks

Theorem A.6.1 (Patching Theorem).

Let $X = \bigcup_{i \in I} U_i$ be an open covering. Suppose that for any $i \in I$ we are given a stack $\mathcal{C}_i$ and for any $i, j \in I$ an equivalence of stacks $\theta_{ij} : \mathcal{C}_j|_{U_{ij}} \xrightarrow{\sim} \mathcal{C}_i|_{U_{ij}}$ satisfying the cocycle relation $\theta_{ij}|_{U_{ijk}} \circ \theta_{jk}|_{U_{ijk}} = \theta_{ik}|_{U_{ijk}}$.

Then there exists a stack $\mathcal{C}$ on X , unique up to equivalence of stacks and equivalences $\theta_i : \mathcal{C}|_{U_i} \xrightarrow{\sim} \mathcal{C}_i$ staisfying $\theta_{ij} \circ \theta_j|_{U_{ij}} = \theta_i|_{U_{ij}}$.

B The functor of ind-micolocalization

In this appendix, we first recall some definitions and statements of the theory of analytic ind-sheaves from [KS2]. Then we define Kashiwara's functor of ind-micolocalization and give (without proof) some basic properties that we used in Sections 8 and 9.

B.1 Ind-sheaves

If $\mathcal{C}$ is a category, one embeds $\mathcal{C}$ into the category of presheaves (of sets) on $\mathcal{C}$ by the fully faithful Yoneda-functor:

$$\mathcal{C} \longrightarrow \widehat{\mathcal{C}} \quad ; \quad A \mapsto \text{Hom}_{\mathcal{C}}(\cdot, A)$$

where $\widehat{\mathcal{C}}$ is the category of contravariant functors $\mathcal{C} \rightarrow \mathcal{Set}$. An object in the essential image of the Yoneda-functor is called representable.

Note that $\widehat{\mathcal{C}}$ admits all small colimits since the category $\mathcal{Set}$ does but even if $\mathcal{C}$ admits colimits the Yoneda-functor does not commute with them.

One denotes by $\text{Ind}\mathcal{C}$ the full subcategory of $\widehat{\mathcal{C}}$ formed by small filtered colimits of representable objects and calls it the category of ind-objects of $\mathcal{C}$. Then $\text{Ind}\mathcal{C}$ admits all small filtered colimits.

If $\mathcal{C}$ is abelian then $\text{Ind}\mathcal{C}$ is abelian and the Yoneda-functor induces an exact fully faithful functor $\mathcal{C} \rightarrow \text{Ind}\mathcal{C}$.

Now let X be a locally compact topological space and fix a field k . One sets

$$\mathbf{I}(k_X) = \text{Ind } \mathcal{M}od^c(k_X)$$

where $\mathcal{M}od^c(k_X)$ denotes the full subcategory of $\mathcal{M}od(k_X)$ formed by sheaves with compact support. We call $\mathbf{I}(k_X)$ the category of ind-sheaves (of k -vector spaces). One can show that the prestack $X \supset U \mapsto \mathbf{I}(k_U)$ is a proper stack, in particular it is an abelian stack.

There are three important basic functors for ind-sheaves

$$\begin{aligned} \iota & : \mathcal{M}od(k_X) \longrightarrow \mathbf{I}(k_X) \quad ; \quad \mathcal{F} \mapsto \varinjlim_{U \subset \subset X} \mathcal{F}_U \\ \alpha & : \mathbf{I}(k_X) \longrightarrow \mathcal{M}od(k_X) \quad ; \quad \varinjlim_{i \in I} \mathcal{F}_i \mapsto \varinjlim_{i \in I} \mathcal{F}_i \\ \beta & : \mathcal{M}od(k_X) \longrightarrow \mathbf{I}(k_X) \quad \text{left adjoint to } \alpha \end{aligned}$$

where we write “ $\varinjlim$ ” for colimits in the category $\mathbf{I}(k_X)$. All three functors induce functors of stacks.

Proposition B.1.1. *(i) The functor ι is fully faithful and exact.*

(ii) The functor α is exact.

(iii) The functor β is fully faithful and exact.

(iv) The triple (β, α, ι) is a triple of adjoint functors, i.e. β is left adjoint to α and α is left adjoint to ι .

Note that since the functors ι, α, β are exact they are well-defined in the derived categories, guard the adjoint properties and α, ι are still fully faithful. An object $\mathcal{F} \in \mathbf{D}^b(k_X)$ is identified with $\iota\mathcal{F}$ in $\mathbf{D}^b(\mathbf{I}(k_X))$.

There are internal operations on ind-sheaves

$$(\cdot) \otimes (\cdot) \quad \text{and} \quad \mathcal{H}om(\cdot, \cdot)$$

and an external

$$\mathcal{H}om(\cdot, \cdot) : \mathbf{I}(k_X) \times \mathbf{I}(k_X) \longrightarrow \mathcal{M}od(k_X).$$

Moreover for any continuous map $f : X \rightarrow Y$ between locally compact spaces we get the external operations

$$f^{-1}, f_*, f_{!!},$$

where the notation $f_{!!}$ indicates that $\iota f_! \neq f_{!!} \iota$.

While $\otimes$ and f^{-1} are exact the other functors have a right derived functor and pass to the derived category where we can define Poincaré-Verdier duality, i.e. we have a right adjoint $f^!$ to $\mathbf{R}f_{!!}$ and we get the usual formalism of Grothendieck’s six operations. We will not recall here the various natural isomorphisms relating these functors and refer to [KS2] but let us summarize the commutation properties with ι, α, β :

Proposition B.1.2. (i) The functor ι commutes to $\otimes, f^{-1}, f^!, Rf_*$.

(ii) The functor α commutes to $\otimes, f^{-1}, Rf_*, Rf_!$ and $\alpha R\mathcal{H}om(\cdot, \cdot) \simeq R\mathcal{H}om(\cdot, \cdot)$.

(iii) The functor β commutes to $\otimes, f^{-1}$.

Finally let us state the following Proposition which has no counterpart in classical sheaf theory:

Proposition B.1.3. Let $\mathcal{F}, \mathcal{G} \in D^b(k_X)$ and $\mathcal{M} \in D^b(I(k_X))$. Then there is a natural isomorphism

$$R\mathcal{H}om(\mathcal{F}, \mathcal{M}) \otimes \beta \mathcal{G} \xrightarrow{\sim} R\mathcal{H}om(\mathcal{F}, \mathcal{M} \otimes \beta \mathcal{G}).$$

B.2 Microlocalization of ind-sheaves

In [K5], Kashiwara establishes the following theorem

Theorem B.2.1. There is a functor

$$\mu : D^b(I(k_X)) \longrightarrow D^b(I(k_{T^*X}))$$

such that for any $\mathcal{F}, \mathcal{G} \in D^b(k_X)$ we have a natural isomorphism

$$R\mathcal{H}om(\mu \mathcal{F}, \mu \mathcal{G}) \simeq R\mathcal{H}om(\pi^{-1} \mathcal{F}, \mu \mathcal{G}) \simeq \mu hom(\mathcal{F}, \mathcal{G}).$$

Remark B.2.2. Note that if $\mathcal{F} \in D^b(k_X)$, then

$$\text{supp}(\mu \mathcal{F}) = \text{supp}(R\mathcal{H}om(\mu \mathcal{F}, \mu \mathcal{F})) = \text{supp}(\mu hom(\mathcal{F}, \mathcal{F})) = \text{SS}(\mathcal{F}).$$

The construction of μ is rather straight-forward if we want to have the property of the Theorem (cf. Proposition 9.1.3 in Section 9.1). Let us recall the definition. The normal deformation of the diagonal in $T^*X \times T^*X$ can be visualized by the following diagram

$$\begin{array}{ccccc} TT^*X & \xrightarrow{\sim} & T_{\Delta_{T^*X}}(T^*X \times T^*X) & \xhookrightarrow{s} & \widetilde{T^*X \times T^*X} & \xleftarrow{j} & \Omega \\ & & \downarrow \tau_{T^*X} & & \downarrow p & \nearrow \tilde{p} & \\ & & T^*X & \xhookrightarrow{\Delta_{T^*X}} & T^*X \times T^*X & & \end{array}$$

Note that $\tilde{p}$ is smooth but p is not. Also, the square is not cartesian. Set

$$K_X = R p_{!!} (k_{\overline{\Omega}} \otimes \beta(k_P)) \otimes \beta(\omega_{\Delta_{T^*X}|T^*X \times T^*X}^{\otimes -1})$$

where the set $P \subset TT^*X$ is defined by

$$P = \left\{ (x, \xi; v_x, v_\xi) \mid \langle v_x, \xi \rangle \geq 0 \right\}.$$

Definition B.2.3. Kashiwara's functor of microlocalisation is defined on T^*X as

$$\mu : D^b(I(k_X)) \longrightarrow D^b(I(k_{T^*X})) ; \mathcal{F} \mapsto \mu \mathcal{F} = K_X \circ \pi^{-1} \mathcal{F}.$$

Lemma B.2.4. Let $S \subset T^*X$ be an arbitrary subset. Then μ defines functors

$$\mu : D^b(k_X, S) \longrightarrow D^b(I(k_S)).$$

If one considers these functors for open subsets $U \subset T^*X$, they define functors of prestacks.

Proof. It is enough to show the existence of the first functor. If $\mu(\mathcal{F})|_S \simeq 0$, then $\text{supp}(\mu(\mathcal{F})) \cap S = \emptyset$, hence $\text{SS}(\mathcal{F}) \cap S = \emptyset$ and $\mu(\cdot)|_S$ factors through $D^b(k_X, S)$. $\square$

The following Proposition is used in Section 9.1.

Proposition B.2.5. *Let $\mathcal{F} \in D_{\mathbb{R}\text{-}c}^b(k_X)$ and $\mathcal{G} \in D^b(I(k_X))$ and assume that*

$$\text{SS}(\mathcal{F}) \cap \text{supp}(\mu\mathcal{G}) \subset T_X^*X$$

Then there is a natural isomorphism

$$\text{R}\mathcal{H}om(\mathcal{F}, k_X) \otimes \mathcal{G} \xrightarrow{\sim} \text{R}\mathcal{H}om(\mathcal{F}, \mathcal{G}).$$

The main theorem of [K5] is the microlocal composition theorem that we included in Section 8.3 (Theorem 8.3.1).

References

- [An1] E. Andronikof, *A microlocal version of the Riemann-Hilbert correspondance*, Topol. Methods in Nonlinear Anal. 4, 1994.
- [An2] E. Andronikof, *Microlocalisation tempérée*, Bull. d. l. SMF 122, 1994.
- [BBD] B. Beilinson, J. Bernstein, and P. Deligne, *Faisceaux pervers*, Analyse et Topologie sur les Espaces Singuliers, Astérisque 100, Soc. Math. France, 1982, pp. 1-171.
- [D'A] A. D'Agnolo, *On the microlocal cutoff of sheaves*, Topol. Methods in Nonlinear Anal. 8, 1996.
- [GMV] S. Gelfand, R. MacPherson and K. Vilonen, *Microlocal Perverse Sheaves*, announced.
- [K1] M. Kashiwara, *Systems of microdifferential equations*, Progress in Math. 34, Birkhäuser, 1983.
- [K2] M. Kashiwara, *The Riemann-Hilbert Problem for Holonomic Systems*, Publ. RIMS, Kyoto Univ. 20, 1984, pp 319-365.
- [K3] M. Kashiwara, *Introduction to microlocal analysis*, L'Enseignement Mathématique, Monographie 32, 1986.
- [K4] M. Kashiwara, *Quantization of Contact Manifolds*, Publ. RIMS, Kyoto Univ. 32, 1996.
- [K5] M. Kashiwara, *Ind-Microlocalization*, written by F. Ivorra and I. Waschkie following a manuscript of M. Kashiwara, to appear 2002.
- [KK] M. Kashiwara and T. Kawai, *On Holonomic Systems of Microdifferential Equations III*, Publ. RIMS, Kyoto Univ. 17, 1981.
- [KS1] M. Kashiwara and P. Schapira, *Sheaves on Manifolds*, Springer, 1990.
- [KS2] M. Kashiwara and P. Schapira, *Ind-Sheaves*, Astérisque 271, Soc. Math. France, 2001.
- [KS3] M. Kashiwara and P. Schapira, *Microlocal Study of Ind-Sheaves I: Micro-Support and Regularity*, preprint math.AG/0108065, 2001.

- [KS4] M. Kashiwara and P. Schapira, *Categories and Sheaves*, to appear 2003.
- [McL] S. MacLane, *Categories for the working mathematician*, Springer.
- [Sch] P. Schapira, *Microdifferential Systems in the Complex Domain*, Grundlehren der mathematischen Wissenschaften 269, Springer, 1985.
- [St] R. Street, *Categorical structures*, Handbook of algebra, Vol. 1, pp. 529–577, North-Holland, 1996.
- [SKK] M. Sato, T. Kawai and M. Kashiwara, *Hyperfunctions and pseudo-differential equations*, Proceedings Katata 1971, H. Komatsu (ed.), Lecture Notes in Math., vol. 287, Springer, 1973, pp. 265-529.
- [Th] R. W. Thomason, *Homotopy colimits in the category of small categories*, Math. Proc. Camb. Phil. Soc. 85, pp. 91–109, 1979.

Ingo Waschkes
 Université Pierre et Marie Curie
 Institut de Mathématiques
 175, rue du Chevaleret, 75013 Paris, France
 E-mail: ingo@math.jussieu.fr